

# Domination, packing and excluded minors

Thomas Böhme<sup>\*†</sup>

Institut für Mathematik  
Technische Universität Ilmenau  
Ilmenau, Germany

Bojan Mohar<sup>‡§</sup>

Department of Mathematics  
University of Ljubljana  
Ljubljana, Slovenia

April 19, 2001

## Abstract

Let  $\gamma(G)$  be the domination number of a graph  $G$ , and let  $\alpha_k(G)$  be the maximum number of vertices in  $G$ , no two of which are at distance  $\leq k$  in  $G$ . It is easy to see that  $\gamma(G) \geq \alpha_2(G)$ . In this note it is proved that  $\gamma(G)$  is bounded from above by a linear function in  $\alpha_2(G)$  if  $G$  has no large complete bipartite graph minors. Extensions to other parameters  $\alpha_k(G)$  are also derived.

## 1 Introduction and main results

Let  $G$  be a finite undirected graph. A graph  $H$  is a *minor* of  $G$  if it can be obtained from a subgraph of  $G$  by contracting edges. The *distance*  $\text{dist}_G(x, y)$  in  $G$  of two vertices  $x, y \in V(G)$  is the length of a shortest  $(x, y)$ -path in  $G$ . The distance of a vertex  $x$  from a set  $A \subseteq V(G)$  is  $\min\{\text{dist}_G(x, a) \mid a \in A\}$ .

For a set  $A \subseteq V(G)$ ,  $G(A)$  denotes the subgraph of  $G$  induced by  $A$ . If  $k$  is a nonnegative integer, we denote by  $\overline{N}_k(A)$  the set of all vertices of  $G$  which are at distance  $\leq k$  from  $A$ . The set  $A$  is a  $k$ -*dominating set* in  $G$  if  $\overline{N}_k(A) = V(G)$ . The cardinality of a smallest  $k$ -dominating set of  $G$  is denoted by  $\gamma_k(G)$ . A vertex set  $X_0 \subseteq V(G)$  is an  $\alpha_k$ -*set* if no two vertices in  $X_0$  are at distance  $\leq k$  in  $G$ . Let  $\alpha_k(G)$  denote the cardinality of a largest  $\alpha_k$ -set of  $G$ . Observe that  $\gamma(G) = \gamma_1(G)$  and  $\alpha(G) = \alpha_1(G)$  are the usual

---

<sup>\*</sup>Supported by SLO-German grant SVN 99/003.

<sup>†</sup>E-mail address: [tboehme@theoinf.tu-ilmenau.de](mailto:tboehme@theoinf.tu-ilmenau.de)

<sup>‡</sup>Supported in part by SLO-German grant SVN 99/003.

<sup>§</sup>E-mail address: [bojan.mohar@uni-lj.si](mailto:bojan.mohar@uni-lj.si)

*domination number* and the *independence* (or *stability*) *number* of  $G$ . We refer to [1] for further details on domination in graphs.

It is clear that  $\gamma_k(G) \geq \alpha_{2k}(G)$ . On the other hand, for any  $r$  there is a graph such that  $\alpha_{k+1}(G) = 1$  and  $\gamma_k(G) \geq r$ . In order to see this, let  $H_n$  be the Cartesian product of  $k+1$  copies of the complete graph  $K_n$ . Then any two vertices of  $H_n$  have distance at most  $k+1$  in  $H_n$ . Therefore,  $\alpha_{k+1}(H_n) = 1$ . Since  $\deg_{H_n}(x) = (k+1)(n-1)$  and  $|V(H_n)| = n^{k+1}$ , it follows that  $\gamma_k(H_n) \geq n/(k+1)^k$ .

The main result of the present note is the following theorem which gives a linear upper bound on  $\gamma_k(G)$  in terms of  $\alpha_m(G)$ ,  $k \leq m < \frac{3}{2}(k+1)$ , in any set of graphs with a fixed excluded minor.

**Theorem 1.1** *Let  $k \geq 0$  and  $m \geq 1$  be integers such that  $k \leq m < \frac{3}{2}(k+1)$ . If  $\gamma_k(G) \geq (2mr + (q-1)(mr - r + 1))\alpha_m(G) - 2mr + r + 1$ , then  $G$  has a  $K_{q,r}$ -minor.*

Our original motivation was the case when  $k = 1$  and  $m = 2$ .

**Corollary 1.2** *If  $\gamma(G) \geq (4r + (q-1)(r+1))\alpha_2(G) - 3r + 1$ , then  $G$  has a  $K_{q,r}$ -minor.*

By excluding  $K_{3,3}$ -minors, we get:

**Corollary 1.3** *If  $G$  is a planar graph, then  $\gamma(G) \leq 20\alpha_2(G) - 9$ .*

The existence of a linear bound  $\gamma(G) \leq c_1\alpha_2(G) + c_2$  for planar graphs was conjectured by F. Göring (private communication) who proved such a bound for plane triangulations.

Corollary 1.3 can be generalized to graphs on any surface. Since the graph  $K_{3,k}$  cannot be embedded in a surface of Euler genus  $g \leq (k-3)/2$  the following bound holds:

**Corollary 1.4** *Suppose that  $G$  is a graph embedded in a surface of Euler genus  $g$ . Then  $\gamma(G) \leq 4(2g+5)\alpha_2(G) - 9$ .*

The special case of Theorem 1.1 when  $k = 0$  and  $m = 1$  is also interesting. The proof of Theorem 1.1 in this special case yields an even stronger statement since the sets  $A_1, \dots, A_r$  in that proof are mutually at distance 1 and hence, in the constructed minor  $K_{q,r}$ , any two of the  $r$  vertices in the second bipartition class are adjacent. Since  $\gamma_0(G) = |V(G)|$ , the following result is obtained:

**Corollary 1.5** *Let  $K_{q,r}^+$  be the graph obtained from  $K_{q,r}$  by adding the  $r$ -clique on the vertex set of the bipartition class of cardinality  $r$ . Suppose that  $K_{q,r}^+$  is not a minor of  $G$ . Then*

$$\alpha(G) \geq \frac{|V(G)| + r}{2r + q - 1}.$$

Duchet and Meyniel [2] obtained a special case of Corollary 1.5 when  $q \leq 1$ . (Note that  $K_{1,r-1}^+ = K_{0,r}^+ = K_r$ .) They proved that in a graph  $G$  without  $K_r$  minor

$$\alpha(G) \geq \frac{|V(G)| + r - 1}{2r - 2}. \quad (1)$$

As it turns out, our proof of Theorem 1.1 restricted to this special case is quite similar to Duchet and Meyniel's proof.

Altohough Theorem 1.1 does not work for the case  $k = 1$  and  $m = 3$ , the following result can be used to get such an extension:

**Corollary 1.6** *Let  $k \geq 0$  be an integer and let  $G$  be a graph. Let  $r$  be the largest integer such that  $K_r$  is a minor of  $G$ . Then*

$$\alpha_{2k}(G) \leq r(2\alpha_{2k+1}(G) - 1).$$

**Proof.** Let  $S$  be a maximum  $\alpha_{2k}$ -set in  $G$ . Define a graph  $H$  with  $V(H) = S$  in which two vertices  $x, y$  are adjacent if and only if  $\text{dist}_G(x, y) = 2k + 1$ . Suppose that  $K$  is a subgraph of  $H$ . Let  $K'$  be a subgraph of  $G$  obtained by taking vertices in  $V(K)$  and, for each edge  $xy$  of  $K$ , adding a path of length  $2k + 1$  in  $G$  joining  $x$  and  $y$ . Since all such paths are geodesics of odd length  $2k + 1$ , they cannot intersect each other. This implies that  $K'$  is a subdivision of  $K$ . In particular, if  $H$  has  $K_r$  minor, so does  $G$ .

Clearly,  $\alpha(H) \leq \alpha_{2k+1}(G)$ . Since  $|V(H)| = \alpha_{2k}(G)$ , (1) implies that  $H$  contains  $K_r$  minor, where  $r \geq \alpha_{2k}(G)/(2\alpha_{2k+1}(G) - 1)$ . Then also  $G$  contains a  $K_r$  minor, and this completes the proof.  $\square$

The relation between  $\alpha_{2k}$  and  $\alpha_{2k+1}$  in Corollary 1.6 cannot be extended to  $\alpha_{2k+1}$  and  $\alpha_{2k+2}$  as shown by the following examples (which are all planar and hence  $K_{3,3}$  minor free). Let  $T_k$  be the tree obtained from the star  $K_{1,p}$  ( $p \geq 1$ ) by replacing each edge by a path of length  $k + 1$ . Then  $\gamma_k(T_k) = p$  (if  $k \geq 1$ ),  $\alpha_{2k+1}(T_k) = p$ , and  $\alpha_{2k+2}(T_k) = 1$ . This example also shows that Theorem 1.1 cannot be extended to the value  $m = 2k + 2$  if  $k \geq 1$ .

## 2 Proof of Theorem 1.1

In this section,  $k$  and  $m$  will denote fixed nonnegative integers such that  $k \leq m \leq 2k + 1$ . Let  $G$  be a graph, and  $A \subseteq V(G)$ . Let  $Q = Q_k^m(A)$  be the subgraph of  $G$  which is obtained from the vertex set  $U = U_k(A) := V(G) \setminus \bar{N}_k(A)$  by adding vertices and edges of all paths of length  $\leq m$  in  $G$  which connect two vertices in  $U$ . Since  $m \leq 2k + 1$ ,  $V(Q) \cap A = \emptyset$ . Observe that  $U = \emptyset$  if and only if  $A$  is a  $k$ -dominating set of  $G$ .

An *extended  $\alpha_m$ -pair* with respect to  $A$  and  $k$  is a pair  $(X, X_0)$  where  $X_0 \subseteq X \subseteq V(G)$  such that:

- (a)  $X_0 \subseteq U_k(A)$  is an  $\alpha_m$ -set in  $G$  and every vertex in  $U_k(A)$  is at distance  $\leq m$  from  $X_0$ .
- (b) Every vertex of  $X \setminus X_0$  lies on an  $(X_0, X_0)$ -path in  $Q = Q_k^m(A)$  which is of length  $\leq 2m$ .
- (c) Every component of  $Q$  contains precisely one connected component of  $Q(X)$ .

Observe that by (a),  $X_0 \neq \emptyset$  if  $A$  is not  $k$ -dominating.

**Lemma 2.1** *If  $k \leq m \leq 2k + 1$  and  $A \subseteq V(G)$ , then there exists an extended  $\alpha_m$ -pair  $(X, X_0)$  with respect to  $A$  and  $k$ . If  $m \geq 1$  and  $A$  is not  $k$ -dominating, then  $|X| \leq 2m|X_0| - 2m + 1$ .*

**Proof.** If  $A$  is  $k$ -dominating, then  $X_0 = X = \emptyset$  will do. If  $m = 0$ , then  $X_0 = X = U_k(A)$ . Suppose now that  $A$  is not  $k$ -dominating and that  $m \geq 1$ . Let  $B$  be a component of  $Q$ . Let  $B_0 = B \cap G(U)$  and  $V_0 = V(B_0)$ . Let us build a set  $X \subseteq V(B)$  and the corresponding  $\alpha_m$ -set  $X_0 \subseteq V_0$  as follows. Start with  $X = X_0 = \{v\}$ , where  $v \in V_0$ . If there exists a vertex of  $V_0$  at distance in  $B$  at least  $m + 1$  from the current set  $X_0$ , let  $u$  be such a vertex which is as close as possible to  $X_0$  in  $B$ . Observe that  $\text{dist}_G(u, X_0) \geq m + 1$  although the distance in  $G$  may be smaller than the distance in  $B$ .

Let  $u_0u_1\dots u_r$  be a shortest path in  $B$  from  $X_0$  (so  $u_0 \in X_0$ ) to  $u = u_r \in V_0$ . Then  $\text{dist}_B(u_i, X_0) = i$  for  $i = 0, \dots, r$ . Suppose that  $r > 2m$ . The vertices  $u_{m+1}, \dots, u_{r-1}$  do not belong to  $V_0$  since their distance from  $X_0$  is  $\geq m + 1$  but smaller than the distance between  $u$  and  $X_0$ . Let  $p = r - \lfloor \frac{m}{2} \rfloor - 1$ . By the definition of  $B$ , the edge  $u_pu_{p+1}$  lies on a path of length  $\leq m$  joining two vertices of  $V_0$ . In particular, an end  $u'$  of this edge is at distance  $\leq \lceil \frac{m}{2} \rceil - 1$  from a vertex  $u'' \in V_0$ . If  $\text{dist}_B(u'', X_0) \leq m$ , then  $\text{dist}_B(u, X_0) \leq \text{dist}_B(u, u') + \text{dist}_B(u', u'') + \text{dist}_B(u'', X_0) \leq (\lfloor \frac{m}{2} \rfloor + 1) +$

$(\lceil \frac{m}{2} \rceil - 1) + m < r$ . This contradiction shows that  $\text{dist}_B(u'', X_0) \geq m + 1$ . However,  $\text{dist}_B(u'', X_0) \leq \text{dist}_B(u'', u') + \text{dist}_B(u', X_0)$ . If  $m$  is even, this implies that  $\text{dist}_B(u'', X_0) < r$ . If  $m$  is odd, then we may assume that  $u' = u_p$ , and then the same conclusion holds. This contradiction to the choice of  $u$  implies that  $\text{dist}_B(u, X_0) = r \leq 2m$ .

Let us add  $u$  into  $X_0$  and add the vertices  $u_0, u_1, \dots, u_r$  into the set  $X$ . This procedure gives rise to an extended  $\alpha_m$ -pair inside  $B$ . Clearly,  $|X| \leq 2m|X_0| - 2m + 1$ .

By taking the union of such sets constructed in all components of  $Q$ , an appropriate extended  $\alpha_m$ -pair is obtained.  $\square$

**Proof of Theorem 1.1.** By Lemma 2.1, there are pairwise disjoint vertex sets  $A_1, A_2, \dots, A_r$  such that  $(A_1, A_1^0)$  is an extended  $\alpha_m$ -pair with respect to  $k$  and  $A^{(1)} = \emptyset$ , and  $(A_i, A_i^0)$  is an extended  $\alpha_m$ -pair with respect to  $k$  and the set  $A^{(i)} := A_1 \cup \dots \cup A_{i-1}$ , for  $i = 2, \dots, r$ . Moreover,  $|A_i| \leq 2m\alpha_m - 2m + 1$ , where  $\alpha_m = \alpha_m(G)$ . Suppose that  $\gamma_k(G) \geq (2mr + (q-1)(mr - r + 1))\alpha_m - 2mr + r + 1$ . Then  $\gamma_k(G) > (2m\alpha_m - 2m + 1)(r - 1)$ , so  $A^{(r)}$  is not a  $k$ -dominating set. Therefore,  $A_1, \dots, A_r$  are all nonempty.

For  $i = 1, \dots, r$ , let  $H_i = Q_k^m(A^{(i)})$ . Let  $H_r^1, \dots, H_r^t$  be the connected components of  $H_r$ . If  $i \geq 2$ , then  $H_i \subseteq H_{i-1}$ . This implies that each component of  $H_i$  is contained in some component of  $H_{i-1}$ . For  $j = 1, \dots, t$ , let  $H_i^j$  be the component of  $H_i$  containing  $H_r^j$ . By (c), each  $H_i^j$  contains a component  $C_r^j$  of  $H_r(A_r)$ . Each  $C_r^j$  contains at least one vertex from the  $\alpha_m$ -set  $A_r^0$ . Therefore,  $t \leq \alpha_m$ .

Let  $B_1 = A_1 \cup \dots \cup A_r$ . Since  $\gamma_k(G) > r(2m\alpha_m - 2m + 1)$ ,  $B_1$  is not  $k$ -dominating. Hence, there is a vertex  $v_1 \in U_k(B_1)$ . By (a),  $v_1$  is at distance  $\leq m$  from some component  $C_r^j$  ( $1 \leq j \leq t$ ) of  $H_r(A_r)$ . Then  $H_r^j, H_{r-1}^j, \dots, H_1^j$  are the components of  $H_r, H_{r-1}, \dots, H_1$  (respectively) containing  $C_r^j$ . For any of the components  $H_i^j$  ( $1 \leq i \leq r$ ), there is a path  $P_i^1$  in  $G$  of length  $\leq m$  connecting  $v_1$  with  $C_i^j \subseteq H_i^j$ . Let  $B_2$  be the union of  $B_1$  with  $\{v_1\}$  and the internal vertices of the paths  $P_1^1, P_2^1, \dots, P_r^1$ . Let us repeat the process with  $B_2$  instead of  $B_1$  to obtain a vertex  $v_2 \in U_k(B_2)$  and linking paths  $P_1^2, P_2^2, \dots, P_r^2$  of length  $\leq m$  joining  $v_2$  with  $A_1, A_2, \dots, A_r$ , respectively.

Now, repeat the process by constructing  $B_3$ , obtaining  $v_3$  and paths  $P_1^3, P_2^3, \dots, P_r^3$ , and so on, as long as possible. This way we get a sequence of vertices  $v_1, v_2, \dots, v_s$  and paths of length  $\leq m$  joining these vertices with  $A_1, \dots, A_r$ . The only requirement which guarantees the existence of  $v_1, \dots, v_s$  and the corresponding paths is that  $\gamma_k(G) > r(2m\alpha_m - 2m + 1) + (s-1)(1+r(m-1))$ . Since  $\gamma_k(G) > (2mr + (q-1)(mr - r + 1))\alpha_m - 2mr + r$ ,

we may take  $s > (q-1)\alpha_m \geq (q-1)t$ . Then  $q$  of the vertices among  $v_1, \dots, v_s$  correspond to the same component  $C_r^j$ , say to  $C_r^1$ . Suppose that these vertices are  $v_1, \dots, v_q$ .

Let us now consider two vertices  $v_i, v_j$  ( $1 \leq i < j \leq q$ ) and two of their paths  $P_a^i$  and  $P_b^j$  where  $a \neq b$ . Suppose that they intersect in a vertex  $v$ . Denote by  $y = \text{dist}_G(v, A_a)$ ,  $z = \text{dist}_G(v_j, v)$ , and  $w = \text{dist}_G(v, A_b)$ . Then  $z + y \leq m$ ,  $z + w \leq m$ ,  $z \geq k + 1$ , and  $y + w \geq \text{dist}_G(A_a, A_b) \geq k + 1$ . This implies that  $k \leq y + w - 1 \leq 2m - 2z - 1 \leq 2m - 2k - 3$ . Consequently,  $P_a^i$  and  $P_b^j$  cannot intersect if  $2m < 3(k + 1)$ . In such a case it is easy to verify that vertices  $v_1, \dots, v_q$ , the connected subgraphs  $C_1^1, C_2^1, \dots, C_r^1$  and the linking paths  $P_a^i$  ( $1 \leq i \leq q$ ,  $1 \leq a \leq r$ ) give rise to a  $K_{q,r}$ -minor in  $G$ . This completes the proof of Theorem 1.1.  $\square$

## References

- [1] T. W. Haynes, S. T. Hedetniemi, P. J. Slater, *Fundamentals of Domination in Graphs*, Marcel Dekker, Inc., New York, 1998.
- [2] P. Duchet, H. Meyniel, On Hadwiger's number and the stability number, in "Graph theory (Cambridge, 1981)," pp. 71–73, North-Holland Math. Stud. 62, North-Holland, 1982.