



ELSEVIER

Available online at www.sciencedirect.com



Molecular Phylogenetics and Evolution xxx (2004) xxx–xxx

MOLECULAR
PHYLOGENETICS
AND
EVOLUTION

www.elsevier.com/locate/ympev

Short communication

Mitochondrial DNA phylogeny and rates of larval evolution in *Macrophiothrix* brittlestars

Michael W. Hart^{a,1}, Robert D. Podolsky^{b,*}^a Department of Biology, Dalhousie University, 1355 Oxford Street, Halifax, NS, Canada B3H 4J1^b Department of Biology, University of North Carolina, Chapel Hill, NC 27599, USA

Received 6 September 2004; revised 23 September 2004

Abstract

Phylogenetic analysis has led to significant insights into the evolution of early life-history stages of marine invertebrates. Although echinoderms have been a major focus, developmental and phylogenetic information are relatively poor for ophiuroids, the most species-rich echinoderm class. We used DNA sequences from two mitochondrial genes to develop a phylogenetic hypothesis for 14 brittlestar species in the genus *Macrophiothrix* (Family Ophiotrichidae). Species are similar in adult form and ecology, but have diverse egg sizes and modes of larval development. In particular, two species have rare larval forms with characteristics that are intermediate between more common modes of feeding and non-feeding development. We use the phylogeny to address whether intermediate larval forms are rare because the evolution of a simplified morphology is rapid once food is no longer required for development. In support of this hypothesis, branch lengths for intermediate forms were short relative to those for species with highly derived non-feeding forms. The absolute rarity of such forms makes robust tests of the hypothesis difficult.

© 2004 Published by Elsevier Inc.

Keywords: Ophiuroid; Brittlestar; Facultative feeding; Pluteus; Mode of development; Life-history transition; Mitochondrial DNA

1. Introduction

The evolutionary analysis of early life cycle diversity in marine invertebrates has advanced significantly over the last decade from the use of phylogenetically based comparative methods (Cunningham, 1999; Hadfield et al., 1995; Hart et al., 1997; Jeffery and Emler, 2003; Levitan, 2000; Rouse and Fitzhugh, 1994; Villinski et al., 2002). These analyses have confirmed or challenged a number of assumptions about the order, timing, and frequency of

change in developmental and life-history characters (reviewed in Hart, 2000).

Certain echinoderm taxa feature prominently in this work (Arndt et al., 1996; Emler, 1990; Hart et al., 1997; Jeffery and Emler, 2003; Smith et al., 1995; Wada et al., 1996; Wray, 1996). Gametes and embryos of sea stars (Asterozoa) and sea urchins (Echinozoa) are easy to obtain and culture (Strathmann, 1987), and some larval traits can be inferred indirectly from egg size (Emler et al., 1987) or adult traits (Emler, 1989). In contrast, the study of development and larval biology in brittlestars (Ophiurozoa), the most-species rich echinoderm class, has been hindered by the cryptic lifestyles of adults, a lack of reliable methods for inducing spawning and oocyte maturation (Selvakumaraswamy and Byrne, 2000; Strathmann, 1987), and the absence of adult characters that can diagnose larval traits (Emler, 1989; Hendler, 1978). As in better documented classes, egg size in ophiuroids is corre-

* Corresponding author. Fax: +1 919 962 1625.

E-mail addresses: mike_hart@sfu.ca (M.W. Hart), podolsky@unc.edu (R.D. Podolsky).

¹ Present address: Department of Biological Sciences, Simon Fraser University, 8888 University Drive, Burnaby, British Columbia, Canada V5A 1S6.

Table 1
Modes of larval development and maximum likelihood branch lengths for *Macrophiothrix* species with obligate non-feeding (group 1) or intermediate (group 2) larval forms and feeding modes

Species	Egg volume (nl)	Larval morphology	Can feed?	Must feed?	Branch length					
					No clock			With clock		
					16S	COI	Both	16S	COI	Both
<i>M. belli</i>	35.1	Non-pluteus	No	No	.129	.239	.208	.123	.286	.214
<i>M. lampra</i>	12.6	(non-pluteus)	(No)	(No)	.110	na	na	.111	na	na
<i>M. nereidina</i>	9.7	Non-pluteus	No	No	.154	na	na	.123	na	na
<i>M. caenosa</i>	7.4	Pluteus	No	No	.082	.097	.086	.071	.148	.115
<i>M. rhabdota</i> ^a (to <i>lorioli</i>)	6.3	Pluteus	Yes	No	.023	na	na	.018	na	na
<i>M. lorioli</i>	2.4	Pluteus	Yes	Yes	.062	.089	.084	.042	.081	.062
<i>M. longipeda</i>	1.9	Pluteus	Yes	Yes	na	na	na	na	na	na
<i>M. koehleri</i>	1.6	Pluteus	Yes	Yes	na	na	na	na	na	na
<i>M. paucispina</i>	<1.0	(pluteus)	(Yes)	(Yes)	na	na	na	na	na	na

Also shown are obligate feeding species (group 3) that are important to the analysis of branch lengths. Larval character states are known from larval culture (R. Podolsky, unpublished data) except for those of *M. lampra* and *M. paucispina*, which are inferred from egg size relative to other species. Of nine additional *Macrophiothrix* species for which we have egg size data, all are likely to be obligately feeding plutei. Because COI data were missing for some species that are relevant to the branch length analysis (*M. lampra*, *M. nereidina*) and COI genetic distances were saturated for the two outgroups, branch lengths in units of expected change per nucleotide site are given for 16S (TVM + G model, 18 taxa), COI (GTR + G + I, 12 ophiotrichid taxa only), and both genes analysed together (GTR + G, 12 ophiotrichids plus *Ophionereis*) under maximum likelihood substitution models estimated independently for each analysis. na, not available or applicable.

^a The relevant branch length for estimating persistence time of the intermediate larval form of *M. rhabdota* depends on whether the closest relative with feeding larval development is *M. paucispina* (inferred from egg size, first line) or *M. lorioli* (known from larval culture, second line); see Fig. 1.

lated with mode of development (Hendler, 1991) and developmental patterns are diverse (sensu McEdward and Janies, 1997). However, no species-level phylogenies have been published for ophiuroids that would aid the analysis of early life cycle evolution (McEdward and Miner, 2001).

The brittlestar genus *Macrophiothrix* (Clark; F. Ophiotrichidae) is common to certain Indo-west Pacific coral reef habitats and especially diverse (21 species) in tropical Australia (Hoggett, 1990). Despite the occurrence of exceptionally high local diversity, adults of different *Macrophiothrix* species are similar in size, form, and ecological habits (Hoggett, 1991). The genus, however, exhibits remarkable variation and specificity in early life cycle characters, involving species-specific sperm chemotaxis (Miller, 1997) and extensive variation in egg size and development mode. At Lizard Island, Australia, 12 *Macrophiothrix* congeners vary more than 60-fold in average egg volume (Podolsky, unpublished data) and produce larval forms that include a typical and morphologically complex feeding pluteus, a morphologically simple non-feeding larval form, and two forms that are functionally intermediate (Table 1). Larvae of *M. rhabdota* are facultative planktotrophs that do not require food to complete development, whereas larvae of *M. caenosa* do not feed but retain the pluteus morphology (R. Podolsky, unpublished data). The rarity of these intermediate larval forms—relative to the large number of evolutionary transitions from feeding to

non-feeding larval forms (Duda and Palumbi, 1999; Hart et al., 1997; Jeffery and Emler, 2003; Lieberman et al., 1993; Wray, 1996)—led Wray (1996) and others to propose that intermediate forms are evolutionarily unstable and short-lived. Phylogenetic information can be used to address this hypothesis (Hart, 1996).

We use information from mitochondrial DNA sequences to resolve relationships among 14 species classified in the genus *Macrophiothrix*, two species in the genus *Ophiotrix* not recently referred to *Macrophiothrix* (Hoggett, 1990), and two outgroups from related families. We use the phylogeny to infer relationships between species with functionally intermediate, putatively transitional larval forms and species with non-feeding larval morphologies, and to test the hypothesis that the former species are more recently derived (Hart, 1996).

2. Materials and methods

We obtained new mtDNA sequences from the large subunit ribosomal RNA (16S) and cytochrome *c* oxidase subunit I (COI) genes for 16 ophiotrichid brittlestar species in the genera *Macrophiothrix* and *Ophiotrix* and for two outgroup taxa (the ophiocomid *Ophiarthrum pictum* and the ophionereid *Ophionereis porrecta*; see Appendix A for details of laboratory methods). We analyzed phylogenetic information from both genes 166

167 together by maximum likelihood (ML) methods, and
 168 used bootstrapping and posterior probabilities as esti-
 169 mates of clade support (see Appendix B for details of
 170 analytical methods).

171 To illustrate the utility of the phylogeny for testing
 172 hypotheses about life history evolution, we inferred and
 173 compared times of persistence for species with highly
 174 derived, non-feeding larval forms and for species with
 175 more intermediate modes of larval development. The
 176 intermediate forms develop from relatively large eggs and
 177 do not require food, but retain phenotypic traits of sus-
 178 pension feeding larvae that are absent from the more
 179 derived forms. If branch lengths leading to intermediate
 180 forms are short relative to those leading to highly derived
 181 non-feeding forms, this pattern would support the
 182 hypothesis that intermediate phenotypes are short-lived
 183 steps in a transition from obligate feeding to simplified
 184 non-feeding development, rather than stable develop-
 185 mental modes (Hart, 1996). (This hypothesis assumes
 186 that rates of molecular evolution are similar across lin-
 187 eages with different derived modes of development.)

188 We asked whether the ML branch lengths leading to
 189 two ophiotrichids with intermediate larval phenotypes
 190 (*M. caenosa* and *M. rhabdota*) are shorter than those
 191 leading to three species that are known (*M. nereidina*, *M.*
 192 *belli*) or inferred (*M. lampra*) to have obligate non-feed-
 193 ing larvae. Other species (*M. longipeda*, *M. koehleri*, *M.*
 194 *paucispina*, and *M. lorioli*) that are known or inferred to
 195 produce obligately feeding plutei (R. Podolsky, unpub-
 196 lished data) were used to establish the earliest possible
 197 times of divergence from species with obligate feeding
 198 larval forms (Table 1).

201 3. Results

202
 203 General sequence characteristics (including missing
 204 COI sequence data for some species) are summarized in
 205 Appendix A. Within- and among-species genetic dis-
 206 tances are summarized in Appendix B.

208 3.1. Tree topology

209
 210 We found a single ML tree with a $-\ln(L)$ score of
 211 7788.9 (Fig. 1). Seven clades were moderately or strongly
 212 supported by high bootstrapping percentages and pos-
 213 terior probabilities: *M. leucosticha* + *M. megapoma*; this
 214 species pair in a larger clade with *M. caenosa*, *M. koeh-*
 215 *leri*, and *M. longipeda*; *M. belli* + *M. nereidina*; *M.*
 216 *demessa* + *O. trilineata*; and the clade (*M. lampra*) (*M.*
 217 *lorioli* (*M. rhabdota* + *M. paucispina*)). The three most
 218 basal nodes, and some other interior branches in the tree,
 219 were short and poorly supported, which could reflect low
 220 resolving power of these genes for these taxa or rapid
 221 divergence of clades. We were unable to identify the sis-
 222 ter groups of three species: *M. robillardi*, *M. propinqua*,

223 *O. caespitosa*. Only one of these (*M. robillardi*) lacks
 224 COI data (see Appendix A), which suggests that includ-
 225 ing these taxa with missing data did not greatly impede
 226 our ability to find some well-supported relationships
 227 based on 16S only (see also Appendix B).

228 *Ophiotrix* and *Macrophiotrix* species did not appear
 229 to form reciprocally monophyletic groups: *O. trilineata*
 230 was strongly inferred to be sister species to *M. demessa*
 231 (Fig. 1). The unresolved relationship between *O. caespit-*
 232 *osa* and other ophiotrichids is consistent with some
 233 *Ophiotrix* as sister group to a paraphyletic *Macrophi-*
 234 *otrix*. However, the low resolution of basal relationships
 235 prevents us from drawing a firm conclusion about
 236 monophyly for these two genera: a tree in which *O.*
 237 *caespitosa* + *O. trilineata* was the sister group to *Macro-*
 238 *phiotrix* species (with other relationships among *Macro-*
 239 *phiotrix* as shown in Fig. 1) had a $-\ln(L)$ score of 7799.5
 240 and was not significantly less likely than the ML tree
 241 (7788.9) by the Shimodaira–Hasegawa test in PAUP*
 242 (Swofford, 2002, $P=0.177$). We therefore cannot reject the
 243 hypothesis that *Macrophiotrix* (sensu Hoggett, 1991) is a
 244 monophyletic clade with respect to *Ophiotrix*.

245 3.2. Branch lengths of derived larval forms

246
 247
 248 The terminal branches leading to species with inter-
 249 mediate larval forms (*M. rhabdota*, *M. caenosa*) were rel-
 250 atively short (e.g., 0.023, 0.082, for 16S without a clock)
 251 compared to terminal branches for species with simpli-
 252 fied, non-feeding larval forms. We obtained qualitatively
 253 the same result when we allowed sister lineages to differ
 254 in rates of molecular evolution and when we enforced a
 255 molecular clock (Table 1). Patterns were similar for 16S
 256 sequences alone, for COI, and for both genes analyzed
 257 together. Both species with intermediate forms were
 258 close relatives of species known to develop as feeding
 259 larvae (*M. lorioli*; *M. longipeda*, and *M. koehleri*), so the
 260 intermediate form is assumed to have arisen along the
 261 short terminal branches (Table 1). If *M. paucispina*
 262 (whose mode of larval development is uncertain) were
 263 also a transitional non-feeding pluteus larva, then the
 264 persistence time for that transitional form could be as
 265 long as the sum of the terminal and internal branches
 266 separating *M. rhabdota* and *M. paucispina* from their
 267 common ancestor with *M. lorioli*. Measurements of egg
 268 size from a single museum specimen of *M. paucispina*
 269 (Australian Museum J22076)—which was dried before
 270 ethanol preservation, and could have experienced a
 271 reduction in egg dimensions—indicated an egg volume
 272 (<1 nl) characteristic of species with planktonic, feeding
 273 pluteus larvae (Table 1). This observation suggests that
 274 the transitional larval form of *M. rhabdota* evolved after
 275 its recent divergence from *M. paucispina*, and supports
 276 the inference of the shorter branch lengths in Table 1.

277 In contrast, the terminal branches leading to sister spe-
 278 cies (*M. belli*, *M. nereidina*) with obligate non-feeding lar-



Fig. 1. Phylogram for 16 *Macrothrix* and *Ophiothrix* brittlestar species and an outgroup (*Ophionereis*) based on maximum likelihood analysis of combined 16S rDNA and COI sequences (six-rate GTR + G model). Numbers beside branches are bootstrap percentages (bold) and Bayesian posterior probabilities (italics); only values >50% are shown. Heavy lines show branch lengths relevant to the analysis of modes of larval development (shown as abbreviations to the right of some taxon names): feeding planktonic larvae (P), non-feeding lecithotrophic larvae (L), transitional pluteus larvae that cannot feed (T_L), transitional pluteus larvae that can feed (T_P) (see Table 1). The mode of larval development for *M. paucispina* is inferred from egg size in a museum specimen and is considered uncertain but probably a feeding planktonic larva (P?).

val development were longer (e.g., 0.129, 0.154 for 16S without a clock). The terminal branch lengths for these sister species give a minimum estimate of persistence time for species with obligate non-feeding development if this larval form and life history evolved once in their common ancestor. A third, independent terminal branch for a species assumed to have non-feeding larvae (*M. lampra*) gives a maximum estimate of this persistence time that is also longer (0.110) than branches leading to species with intermediate larval forms. Branch lengths for *M. belli* based on COI (which were missing for *M. lampra* and *M. nereidina*) and the combined sequences were also longer than the comparable branch lengths for species with intermediate forms. Of course, any of these derived larval forms could have evolved at any time since the divergence of the species from an ancestor with obligate feeding larvae, and the number of such lineages is small. In addition, some long

terminal branches might be shortened by the discovery of sister group relationships with other *Macrothrix* species that we were not able to sample.

4. Discussion

4.1. Taxonomic and phylogenetic resolution in *Macrothrix*

Analysis of mtDNA sequences from *Macrothrix* and *Ophiothrix* species identified some sister-group relationships that are statistically well-supported, but left other relationships unresolved. Improved taxonomic sampling of more *Macrothrix* species using more slowly evolving molecular characters is needed to fully resolve these basal relationships. Nevertheless, our

391 current results provide a useful context for comparative
392 analyses of developmental evolution among ophiotri-
393 chids.

394 Our mtDNA sequence comparisons add to earlier
395 efforts of Hoggett (1991) to resolve generic boundaries
396 involving species in *Macrophiothrix* and related sub-
397 genera of *Ophiiothrix*. Among the 16 species for which
398 we generated mtDNA sequences, we found some well-
399 supported clades of species that Hoggett had assigned
400 to *Macrophiothrix*. However, two other *Macrophio-*
401 *thrix* were not reliably grouped with other members of
402 the genus. First, morphological characters reliably
403 placed *M. propinqua* and *M. robillardi* with other
404 *Macrophiothrix* (Hoggett, 1990), but 16S and COI
405 characters could not reliably identify the sister group
406 of either. Second, analyses of mtDNA tend to group
407 *M. demessa* with *O. trilineata*, though Hoggett (1990)
408 assigned each species with confidence to its respective
409 genus on the basis of adult morphological and allo-
410 zyme evidence.

411 We also found examples of strong conflict between
412 our mtDNA phylogeny and cladograms based on mor-
413 phology or allozymes (Hoggett, 1990). Some species that
414 share numerous adult morphological apomorphies, such
415 as *M. belli*, *M. caenosa*, and *M. paucispina* (Hoggett,
416 1990; Fig. 2.17), are distant relatives in the mtDNA trees.
417 Other species with highly divergent allozyme alleles,
418 such as *M. leucosticha* and *M. megapoma* (Hoggett,
419 1990; Fig. 4.3), are strongly supported as sister species
420 in the mtDNA trees. In general, the mtDNA data more
421 finely and robustly resolved species relationships for the
422 clades identified in common by the three analyses.

423 Our analyses of mtDNA and life-history variation
424 support other taxonomic revisions. For example, Hogg-
425 ett (1990) reversed a long history of classification (Clark,
426 1938) by distinguishing *M. caenosa* (sp. nov.) from *M.*
427 *longipeda*. The main diagnostic character (shape of
428 papillae on the adult dental plate) is difficult to resolve
429 with live specimens. In contrast, the species are easily
430 distinguished by a 3.8-fold difference in egg volume and
431 qualitative differences in mode of development (R. Pod-
432 olsky, unpublished data). Mitochondrial DNA
433 sequences extend this conclusion by suggesting that
434 these taxa might not even be sister species (Fig. 1). Simi-
435 lar surveys of reproductive traits could be useful for
436 identifying cryptic species diversity in putative morpho-
437 species such as *M. demessa* that have broad geographic
438 ranges (Philippines through Australia and several oce-
439 anic islands), long mtDNA terminal branches (Fig. 1),
440 and high within-species mtDNA sequence variation
441 (Appendix B).

442 4.2. Evolutionary analysis of early life cycle characters

445 Efforts to understand the origins of developmental
446 diversity have focused on processes that underlie

transitions between modes of development (Wray, 447
1995a). In marine invertebrates, one widespread transi- 448
tion involves the loss of a complex feeding larval mor- 449
phology in association with the evolution of large, 450
nutrient-rich eggs (Emlet et al., 1987; Strathmann, 451
1985). Intermediate phenotypes, such as facultative 452
planktotrophy, could represent short-lived steps on the 453
path toward non-feeding development and morphol- 454
ogy (Wray and Raff, 1991) or advantageous and evolu- 455
tionarily stable modes of larval development (Emlet, 456
1986; McEdward, 1997; Levitan, 2000). Estimates of 457
relative persistence times for these larval forms are a 458
powerful and novel way to evaluate these two hypothe- 459
ses (Hart, 1996; Wray, 1996). 460

Intermediate larval forms are taxonomically rare and 461
phylogenetically scattered (Alatalo et al., 1984; Crump, 462
1989; Emlet, 1986; Hart, 1996; Kempf and Hadfield, 463
1985; Kempf and Todd, 1989; Perron, 1981). As the first 464
genus documented to include more than one species 465
with an intermediate larval form—in addition to several 466
species with more derived, non-feeding forms—*Macro-* 467
phiothrix offered a unique opportunity to apply phylo- 468
genetic information to analysis of this transition. 469
Branch lengths from our phylogeny provided estimates 470
of maximum persistence times for intermediate forms. 471
All of these estimates (from two species, and two genes 472
analyzed separately or in combination, with and with- 473
out rate variation among lineages) were shorter than the 474
relevant branch lengths for species or clades with 475
obligate non-feeding development and simple larval 476
morphology. 477

These data are consistent with predictions of the 478
rapid transitions hypothesis (Hart, 1996; Wray, 1995b). 479
Notably, comparison between the two transitional lar- 480
val forms is also consistent with this hypothesis: the spe- 481
cies (*M. rhabdota*) with larvae that retain the capacity to 482
feed occurs on a shorter branch than the species (*M.* 483
caenosa) with larvae that retain a feeding morphology 484
but have lost the capacity to feed (e.g., 0.081 versus 485
0.148 for COI with a molecular clock). This observation 486
provides indirect support—because these two species are 487
part of separate lineages—for the concept of an ordered 488
transformation series (Wray, 1996) in which feeding 489
capacity and feeding morphology are lost in sequential 490
steps (R. Podolsky, unpublished data). Whether persis- 491
tence times for these intermediate forms are transient 492
enough to explain their absence in most well-sampled 493
taxa that include both feeding and simplified non-feed- 494
ing forms is uncertain, but we have failed to reject this 495
hypothesis for the most extensive phylogenetic example 496
available. As the first species-level phylogeny to address 497
early life cycle evolution in ophiuroids, the analysis 498
presented here is a starting point for improved under- 499
standing of a large, diverse, and ecologically important 500
class for which developmental and life-history data are 501
relatively scarce. 502

503 **Acknowledgment**

504
 505 We thank J. Allen and J. McAlister for discussions
 506 and contributions to larval culture and measurement, J.
 507 Mader and K. Kemp for collection of samples from
 508 museum collections, J. Addison for assistance with
 509 sequencing, P. Berents (Australian Museum) and J. Fro-
 510 mont (Western Australian Museum) for access to and
 511 assistance with collections, D. Posada for software assis-
 512 tance, K. Beckenbach and M. J. Smith for sequence data,
 513 and B. Schierwater and four anonymous reviewers for
 514 constructive criticism. A. Hoggett and L. Vail provided
 515 research space and logistical support at Lizard Island
 516 Research Station. Supported by National Science Founda-
 517 tion grant OCE-9811121, the Canada Foundation for
 518 Innovation, and the Natural Sciences and Engineering
 519 Research Council.

520

521

522 **Appendix A. Laboratory methods and sequence**
523 **characteristics**

524

525 The table below shows reference information for the
 526 mtDNA data. Australian Museum accession numbers for
 527 tissue samples are in parentheses after the species name;
 528 other samples were collected from fresh specimens at Liz-
 529 ard Island, Australia, and fixed in 70% ethanol. Identifi-
 530 cations of live material were based on Hoggett (1990) and
 531 verified by A. Hoggett (personnal communication). Two
 532 GenBank numbers are given for species in which 16S
 533 sequences were obtained from two individuals. We used
 534 standard digestion and extraction methods (e.g., Gros-
 535 berg et al., 1996) to obtain genomic DNA from gonads,
 536 gametes, or arms. From serial dilutions of these DNA
 537 samples, we amplified part of each of the 16S and COI
 538 genes using ‘universal’ PCR primers (Folmer et al., 1994;
 539 Palumbi, 1996). For six species (*M. caenosa*, *M. demessa*,
 540 *M. koehlerii*, *M. longipeda*, *M. lorioli*, and *M. nereidina*)
 541 we obtained 16S sequences from two individuals. Only
 542 four species (*M. belli*, *M. demessa*, *M. longipeda*, and
 543 *Ophionereis*) could be reliably amplified and sequenced
 544 using the universal COI primers; all other COI amplifica-
 545 tions required some variable combination of ten other
 546 oligonucleotide sequences (available from the authors on
 547 request) that we designed from preliminary sequence data
 548 and comparison with complete COI sequences for two
 549 other brittlestars (Scouras et al., 2004). We were unable to
 550 amplify and sequence COI from *M. lampra*, *M. nereidina*,
 551 *M. paucispina*, and *M. robillardii* using the universal or
 552 modified primers. We sequenced these PCR products
 553 directly using the PCR primers, Thermosequenase (USB),
 554 and infrared dye-labeled dideoxy terminators (Li-Cor),
 555 and we visualized the fragments on a Li-Cor 4200L-2
 556 DNA sequencer. We used the manufacturers’ standard
 557 cycle sequencing and electrophoresis conditions. Comple-
 558 mentary sequences were compared and edited in AlignIR

(Li-Cor). Most PCR products were sequenced from both 559
 strands through the end of the opposite primer, but in 560
 some cases we obtained only partial sequences of the 561
 product or sequences from just one strand, so that the 562
 lengths of sequences for each gene varied among taxa. 563

The 24 aligned 16S sequences were 596bp long; 184 of 564
 these sites were parsimony-informative (with gap sites 565
 coded as missing). Many alignment gaps corresponded to 566
 insertions in the *Ophiarthrum* sequence relative to *Ophi-* 567
onereis and the ophiotrichids. Some gaps were large (e.g., a 568
 37-base insertion near the 3’ end of the 16S sequence of 569
M. propinqua). The 14 aligned COI sequences were 654 570
 bp; 225 of these sites were parsimony-informative. We 571
 found one amino acid deletion: a Gly/Gln codon near the 572
 5’ end of the *Ophiarthrum* sequence. 573

574

Taxon names, sequence lengths, and GenBank accession 575
 numbers 576

Species	16S	COI	
Ophiotrichidae			577
<i>Macrophiothrix belli</i>	492 (AY365143)	571 (AY365144)	580
<i>M. caenosa</i>	492 (AY365145)	654 (AY365146)	581
	491 (AY365147)		582
<i>M. demessa</i>	483 (AY365148)	654 (AY365149)	583
	508 (AY365150)		584
<i>M. koehlerii</i>	490 (AY365151)	654 (AY365152)	585
	489 (AY365153)		586
<i>M. lampra</i>	494 (AY365154)		587
(J14001)			588
<i>M. leucosticha</i>	511 (AY365155)	565 (AY365156)	589
<i>M. longipeda</i>	491 (AY365158)	654 (AY365159)	590
	489 (AY365160)		591
<i>M. lorioli</i>	511 (AY365161)	654 (AY365162)	592
	493 (AY365163)		593
<i>M. megapoma</i>	494 (AY365165)	654 (AY365166)	594
<i>M. nereidina</i>	494 (AY365167)		595
	492 (AY365169)		596
<i>M. paucispina</i>	498 (AY365170)		597
<i>M. propinqua</i>	542 (AY365172)	654 (AY365173)	598
<i>M. rhabdota</i>	493 (AY365174)	654 (AY365175)	599
<i>M. robillardii</i>	491 (AY365176)		600
(J19445)			601
<i>Ophiothrix</i>	442 (AY365179)	654 (AY365180)	602
<i>caespitosa</i> (J16397)			603
<i>O. trilineata</i>	491 (AY365182)	654 (AY365183)	604
(J19233)			605
Ophionereidae			606
<i>Ophionereis porrecta</i>	507 (AY365184)	571 (AY365185)	607
			608
Ophiocomidae			609
<i>Ophiarthrum pictum</i>	523 (AY365186)	651 (AY365187)	610
			611
			612
			613
			614

615 Appendix B. Phylogenetic methods and genetic distance 616 results

617
618 The table below shows pairwise genetic distances
619 based on maximum likelihood (ML) models. 16S nucleo-
620 tide sequences were aligned in ClustalX using the default
621 multiple alignment parameters (Jeanmougin et al., 1998).
622 COI sequences were translated using the echinoderm
623 mtDNA translation table in GeneJockey ([http://](http://www.biosoft.com)
624 www.biosoft.com), checked to confirm that they con-
625 tained no frame shifts, nonsense codons, or stop codons,
626 and aligned as amino acid sequences in ClustalX.

627 We analyzed phylogenetic information from both
628 genes together (total evidence; Kluge, 1989). We favored
629 this approach because we preferred to include some data
630 for taxa where we lacked COI sequences (see Appendix
631 A) rather than to leave these taxa out of some analyses
632 entirely. Other recent studies show that adding taxa with
633 missing sequence characters tends to improve the quality
634 of total evidence analyses (Hughes and Vogler, 2004;
635 Wiens, 1998). As a quantitative check on this approach,
636 we used the incongruence length difference (ILD) test in
637 PAUP* (Swofford, 2002). We inferred a preliminary
638 maximum parsimony (MP) tree in PAUP* with equal
639 weighting for transitions and transversions, then esti-
640 mated likelihood scores and the Ti/Tv ratio for sequence
641 data mapped onto that MP tree under a two-parameter
642 ML substitution model. We estimated the appropriate
643 transversion weighting separately for three classes of
644 characters: 16S, COI 3rd codon positions, and all other
645 COI nucleotides. We applied these estimated transver-
646 sion weightings in the ILD test by use of step matrices
647 with accelerated character transformations, heuristic
648 searching, TBR branch swapping, and 1000 iterations of
649 the test. Although we are aware of the uncertainty sur-
650 rounding use of the ILD test for identifying significant
651 conflict between data partitions (e.g., Darlu and Lecointre,
652 2002; Dowton and Austin, 2002; Hipp et al., 2004;
653 Yoder et al., 2001), we found no indication that the 16S
654 and COI sequences gave conflicting information
655 ($P=0.722$), and in subsequent analyses we used both
656 genes together with COI characters coded as missing for
657 *M. lampra*, *M. nereidina*, *M. paucispina*, and *M. robil-*
658 *lardi*.

659 Our analyses of phylogenetic relationships among
660 ophiotrichids emphasized ML methods rather than par-
661 simony methods because our specific goal was to infer
662 branch lengths leading to species with different larval
663 phenotypes under a biologically realistic model of DNA
664 sequence evolution. We used hierarchical likelihood
665 ratio testing in Modeltest (Posada and Crandall, 1998)

to choose the appropriate substitution model (six-rate 666
GTR + G, $\alpha=0.2136$), then estimated the tree topology 667
using this model in PAUP* with stepwise addition of 668
taxa, TBR branch swapping, and no molecular clock. 669
We estimated nodal support for the ML tree as boot- 670
strap percentages in PAUP* (100 replicates) and as the 671
posterior probabilities of clades under the same likeli- 672
hood model in coalescent simulations using MrBAYES 673
(Huelsenbeck and Ronquist, 2001). We ran this ML sim- 674
ulation for 110,000 generations, and discarded the first 675
10,000 generations as a burn-in (in preliminary analyses 676
this number was sufficient to avoid sampling trees and 677
parameter values in the non-stationary part of the simu- 678
lation). We set the model parameter values (six substitu- 679
tion rates, alpha value for among-site rate variation, and 680
nucleotide frequencies) as priors using the “prset” com- 681
mand. We used default values for other Bayesian param- 682
eters, including number of chains (4), temperature (0.2), 683
and sampling frequency (100). 684

685 Because some branch length comparisons involved 685
species for which we lacked COI sequence data, we esti- 686
mated these branch lengths (with and without a molecular 687
clock enforced) under the appropriate substitution model 688
identified for 16S alone (five-rate TVM + G, $\alpha=0.2823$), 689
COI alone (six-rate GTR + G + I, $\alpha=0.5544$, $I=0.4301$), 690
and for the combined data (parameters above). 691

692 Genetic distances between 16S sequences of conspe- 692
cific pairs (bold values below) were smallest in *M. lorioli* 693
(0.017) and largest in *M. demessa* (0.042). These dis- 694
tances are less than or comparable to conspecific genetic 695
distances based on 16S sequences of other ophiotrichids 696
(Baric and Sturmbauer, 1999) or other brittlestar fami- 697
lies (Sponer et al., 2001). Most among-species distances 698
were larger, but two sister species pairs were separated 699
by genetic distances within this range: *M. leucosticha* 700
and *M. megapoma* (0.030); *M. paucispina* and *M. rhab-* 701
dota (0.032). Genetic distances between COI sequences 702
were substantially larger, and reflected more rapid evo- 703
lution of this gene relative to 16S. COI distances 704
between the outgroup taxa and the ophiotrichids were 705
very large or undefined and are not reported in the 706
table. 707

708 In some preliminary phylogenetic analyses, we found 708
very long terminal branch lengths leading to *Ophiar-* 709
thrum in comparison to all other internal and terminal 710
branches. This result reflects the unexpectedly large 711
genetic distances between this species and all other 712
sequences. To avoid artifacts associated with this excep- 713
tionally long branch, we removed *Ophiarthrum* from the 714
analyses reported in the main text, and we used *Ophione-* 715
reis as the sole outgroup. 716

Genetic distances among brittlestar mtDNA sequences for 16S rDNA (below diagonal) and COI (above diagonal). Undefined distances are indicated with an asterisk

	<i>bel</i>	<i>cae1</i>	<i>cae2</i>	<i>dem1</i>	<i>dem2</i>	<i>koe1</i>	<i>koe2</i>	<i>lam</i>	<i>leu</i>	<i>lon1</i>	<i>lon2</i>	<i>lor1</i>	<i>lor2</i>	<i>meg</i>	<i>ner1</i>	<i>ner2</i>	<i>pau</i>	<i>pro</i>	<i>rha</i>	<i>rob</i>	<i>csp</i>	<i>tri</i>	<i>por</i>	<i>pic</i>
<i>belli</i>		0.658	*	0.308	*	0.654	*	*	0.544	0.529	*	0.572	*	0.564	*	*	*	0.467	0.585	*	0.697	0.863	*	*
<i>caenosa1</i>	0.236		*	0.695	*	0.388	*	*	0.259	0.262	*	0.591	*	0.253	*	*	*	0.638	0.523	*	0.980	0.859	*	*
<i>caenosa2</i> ^a	0.240	0.028		*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>demessa1</i>	0.263	0.321	0.307		*	0.842	*	*	0.612	0.592	*	0.571	*	0.659	*	*	*	0.593	0.569	*	0.684	0.655	*	*
<i>demessa2</i> ^a	0.265	0.316	0.293	0.042		*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>koehler1</i>	0.269	0.172	0.163	0.371	0.322		*	*	0.429	0.369	*	0.707	*	0.443	*	*	*	0.809	0.620	*	1.018	1.072	*	*
<i>koehler2</i> ^a	0.257	0.146	0.137	0.334	0.331	0.041		*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>lampra</i> ^a	0.297	0.277	0.276	0.291	0.311	0.341	0.286		*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
<i>leucosticha</i>	0.291	0.141	0.149	0.415	0.361	0.185	0.162	0.362		0.268	*	0.495	*	0.017	*	*	*	0.647	0.576	*	0.741	0.759	*	*
<i>longipeda1</i>	0.260	0.113	0.119	0.321	0.317	0.128	0.110	0.325	0.148		*	0.485	*	0.258	*	*	*	0.612	0.457	*	0.671	0.872	*	*
<i>longipeda2</i> ^a	0.273	0.124	0.114	0.321	0.297	0.145	0.121	0.316	0.157	0.020		*	*	*	*	*	*	*	*	*	*	*	*	*
<i>loriol1</i>	0.245	0.214	0.226	0.319	0.333	0.259	0.234	0.149	0.267	0.258	0.247		*	0.465	*	*	*	0.644	0.157	*	0.622	0.727	*	*
<i>loriol2</i> ^a	0.249	0.239	0.234	0.334	0.351	0.244	0.217	0.148	0.269	0.255	0.247	0.017		*	*	*	*	*	*	*	*	*	*	*
<i>megapoma</i>	0.251	0.106	0.126	0.356	0.350	0.158	0.126	0.309	0.030	0.113	0.120	0.224	0.233		*	*	*	0.643	0.564	*	0.688	0.832	*	*
<i>neraidina1</i> ^a	0.214	0.259	0.259	0.326	0.340	0.280	0.237	0.344	0.282	0.283	0.244	0.311	0.302	0.230		*	*	*	*	*	*	*	*	*
<i>neraidina2</i> ^a	0.223	0.272	0.270	0.362	0.377	0.269	0.236	0.314	0.288	0.249	0.208	0.268	0.267	0.235	0.030		*	*	*	*	*	*	*	*
<i>paucispina</i> ^a	0.324	0.287	0.277	0.400	0.400	0.261	0.241	0.185	0.340	0.319	0.295	0.068	0.057	0.296	0.370	0.344		*	*	*	*	*	*	*
<i>propinqua</i>	0.410	0.416	0.345	0.408	0.408	0.424	0.422	0.356	0.527	0.458	0.450	0.418	0.386	0.500	0.519	0.487	0.385		0.609	*	0.541	0.720	*	*
<i>rhabdota</i>	0.308	0.316	0.285	0.393	0.356	0.260	0.245	0.180	0.344	0.325	0.318	0.073	0.063	0.307	0.379	0.352	0.032	0.314		*	0.709	0.773	*	*
<i>robillard</i> ^a	0.220	0.223	0.191	0.287	0.249	0.190	0.185	0.269	0.247	0.220	0.204	0.227	0.207	0.225	0.284	0.289	0.272	0.351	0.271		*	*	*	*
<i>caespitosa</i>	0.342	0.410	0.375	0.382	0.360	0.299	0.270	0.390	0.407	0.328	0.301	0.315	0.303	0.372	0.359	0.362	0.311	0.572	0.320	0.365		0.835	*	*
<i>trilineata</i>	0.409	0.417	0.371	0.381	0.344	0.404	0.418	0.435	0.508	0.467	0.467	0.393	0.435	0.450	0.461	0.518	0.425	0.524	0.463	0.381	0.360		*	*
<i>ophionereis</i>	0.797	0.764	0.774	0.817	0.774	0.788	0.757	0.725	0.981	0.802	0.701	0.637	0.648	0.869	0.863	0.809	0.705	0.954	0.715	0.818	0.867	0.751		*
<i>ophiarthrum</i>	1.817	1.807	1.808	1.994	2.048	1.773	1.814	2.019	1.930	1.822	1.687	1.922	1.997	1.732	1.768	1.987	1.973	2.284	1.957	1.782	2.262	2.326	1.863	

Distances between conspecifics are shown in bold face.

^a 16S sequence only

717 References

- 718 Alatalo, P., Berg, C.J.J., D'Asaro, C.N., (1984). Reproduction and
719 development in the lucinid clam *Codakia orbicularis*. *Bull. Mar. Sci.*
720 34, 424–434.
- 721 Arndt, A., Marquez, C., Lambert, P., Smith, M.J., (1996). Molecular
722 phylogeny of eastern Pacific sea cucumbers (Echinodermata: Holo-
723 thuroidea) based on mitochondrial DNA sequences. *Mol. Phyloge-
724 net. Evol.* 6, 425–437.
- 725 Baric, S., Sturmbauer, C., (1999). Ecological parallelism and cryptic
726 species in the genus *Ophiothrix* derived from mitochondrial DNA
727 sequences. *Mol. Phylogenet. Evol.* 11, 157–162.
- 728 Clark, H.L., 1938. Echinoderms from Australia. *Memoirs of the*
729 *Museum of Comparative Zoology, Harvard University* 55, 1–
730 596.
- 731 Crump, M.L., (1989). Life history consequences of feeding versus non-
732 feeding in a facultatively non-feeding toad larva. *Oecologia* 78,
733 486–489.
- 734 Cunningham, C.W., (1999). Some limitations of ancestral character-
735 state reconstruction when testing evolutionary hypotheses. *Syst.*
736 *Biol.* 48, 665–674.
- 737 Darlu, P., Lecointre, G., (2002). When does the incongruence length
738 difference test fail?. *Mol. Biol. Evol.* 19, 432–437.
- 739 Dowton, M., Austin, A.D., (2002). Increased congruence does not neces-
740 sarily indicate increased phylogenetic accuracy—the behavior of the
741 incongruence length difference test in mixed-model analyses.
742 *Syst. Biol.* 51, 19–31.
- 743 Duda, T.F., Palumbi, S.R., (1999). Developmental shifts and species
744 selection in gastropods. *Proc. Natl. Acad. Sci. USA* 96, 10272–
745 10277.
- 746 Emlet, R.B., (1986). Facultative planktotrophy in the tropical echinoid
747 *Clypeaster rosaceus* and a comparison with obligate planktotrophy
748 in *Clypeaster subdepressus* (Clypeasteroidea Echinoidea). *J. Exp.*
749 *Mar. Biol. Ecol.* 95, 183–202.
- 750 Emlet, R.B., (1989). Apical skeletons of sea urchins (Echinodermata
751 Echinoidea): two methods for inferring mode of larval develop-
752 ment. *Paleobiology* 15, 223–254.
- 753 Emlet, R.B., (1990). World patterns of developmental mode in echinoid
754 echinoderms. In: Hoshi, M., Yamashita, O. (Eds.), *In: Advances in*
755 *Invertebrate Development*, vol. 5. Elsevier, New York.
- 756 Emlet, R.B., McEdward, L.R., Strathmann, R.R., (1987). Echinoderm
757 larval ecology viewed from the egg. In: Jangoux, M., Lawrence,
758 J.M. (Eds.), *In: Echinoderm Studies*, vol. 2 A.A.. Balkema, Rotter-
759 dam, Netherlands, pp. 55–136.
- 760 Folmer, O., Black, M., Hoeh, W., Lutz, R., Vrijenhoek, R., (1994).
761 DNA primers for amplification of mitochondrial cytochrome c oxi-
762 dase subunit I from diverse metazoan invertebrates. *Mol. Mar. Biol.*
763 *Biotechnol.* 3, 294–297.
- 764 Grosberg, R.K., Levitan, D.R., Cameron, B.B., (1996). Characterization
765 of genetic structure and genealogies using RAPD-PCR markers: a
766 random primer for the novice and nervous. In: Ferraris, J.D.,
767 Palumbi, S.R. (Eds.), *Molecular Zoology: Advances, Strategies, and*
768 *Protocols*. Wiley-Liss, New York, pp. 67–100.
- 769 Hadfield, K.A., Swalla, B.J., Jeffery, W.R., (1995). Multiple origins of
770 anural development in ascidians inferred from rDNA sequences. *J.*
771 *Mol. Evol.* 40, 413–427.
- 772 Hart, M.W., (1996). Evolutionary loss of larval feeding: development,
773 form, and function in a facultatively feeding larva, *Brisaster lati-*
774 *frons*. *Evolution* 50, 174–187.
- 775 Hart, M.W., (2000). Phylogenetic analyses of mode of larval develop-
776 ment. *Sem. Cell Dev. Biol.* 11, 411–418.
- 777 Hart, M.W., Byrne, M., Smith, M.J., (1997). Molecular phylogenetic
778 analysis of life-history evolution in asterinid starfish. *Evolution* 51,
779 1848–1861.
- 780 Hendler, G., (1978). Development of *Amphiplus abditus* (Echinoder-
781 mata Ophiuroidea). Part 2. Description and discussion of ophiuroid
782 skeletal ontogeny and homologies. *Biol. Bull.* 154, 79–95.
- Hendler, G., (1991). Echinodermata: Ophiuroidea. In: Giese, A.C., 783
Pearse, J.S., Pearse, V.B. (Eds.), *Reproduction of Marine Inverte-*
784 *brates*, vol. VI Echinoderms and Lophophorates. Boxwood Press,
785 Pacific Grove, CA. 786
- Hipp, A.L., Hall, J.C., Sytsma, K.J., (2004). Congruence versus phyloge- 787
netic accuracy: revisiting the incongruence length difference test. 788
Syst. Biol. 53, 81–89. 789
- Hoggett, A., 1990. Taxonomy and systematic position of the brittlestar 790
genus *Macrophiothrix* H.L. Clark (Echinodermata: Ophiuroidea). 791
Dissertation, University of Queensland, Brisbane, Australia.. 792
- Hoggett, A.K., (1991). The genus *Macrophiothrix* (Ophiuroidea: 793
Ophiotrichidae) in Australian waters. *Invertebr. Taxon.* 4, 1077–
794 1146. 795
- Huelsenbeck, J.P., Ronquist, F., (2001). MRBAYES: Bayesian infer- 796
ence of phylogenetic trees. *Bioinformatics* 17, 754–755. 797
- Hughes, J., Vogler, A.P., (2004). The phylogeny of acorn weevils (genus 798
Curculio) from mitochondrial and nuclear DNA sequences: the 799
problem of incomplete data. *Mol. Phylogenet. Evol.* 32, 601–615. 800
- Jeanmougin, F., Thompson, J.D., Gouy, M., Higgins, D.G., Gibson, 801
T.J., (1998). Multiple sequence alignment with Clustal X. *Trends* 802
Biochem. Sci. 23, 403–405. 803
- Jeffery, C.H., Emlet, R.B., (2003). Macroevolutionary consequences of 804
developmental mode in temnopleurid echinoids from the tertiary of 805
southern Australia. *Evolution* 57, 1031–1048. 806
- Kempf, S.C., Hadfield, M.G., (1985). Planktotrophy by the lecitho- 807
trophic larvae of a nudibranch *Phestilla sibogae* Gastropoda. *Biol.* 808
Bull. 169, 119–130. 809
- Kempf, S.C., Todd, C.D., (1989). Feeding potential in the lecitho- 810
trophic larvae of *Adalaria proxima* and *Tritonia hombergi*: an evo- 811
lutionary perspective. *J. Mar. Biol. Assoc. U.K.* 69, 659–682. 812
- Kluge, A., (1989). A concern for evidence and a phylogenetic hypothe- 813
sis for relationships among *Epicrates* (Boidae, Serpentes). *Syst.* 814
Zool. 38, 1–25. 815
- Levitan, D.R., (2000). Optimal egg size in marine invertebrates: the- 816
ory and phylogenetic analysis of the critical relationship between 817
egg size and development time in echinoids. *Am. Nat.* 156, 175–
818 192. 819
- Lieberman, B.S., Allmon, W.D., Eldredge, N., (1993). Levels of selec- 820
tion and macroevolutionary patterns in the turritellid gastropods. 821
Paleobiology 19, 205–215. 822
- McEdward, L.R., (1997). Reproductive strategies of marine benthic 823
invertebrates revisited: facultative feeding by planktotrophic lar- 824
vae. *Am. Nat.* 150, 48–72. 825
- McEdward, L.R., Janies, D.A., (1997). Relationships among develop- 826
ment, ecology, and morphology in the evolution of echinoderm lar- 827
vae and life cycles. *Biol. J. Linn. Soc.* 60, 381–400. 828
- McEdward, L.R., Miner, B.G., (2001). Larval and life-cycle patterns in 829
echinoderms. *Can. J. Zool.* 79, 1125–1170. 830
- Miller, R.L., (1997). Specificity of sperm chemotaxis among Great Bar- 831
rier Reef shallow-water holothurians and ophiuroids. *J. Exp. Zool.* 832
279, 189–200. 833
- Palumbi, S.R., (1996). Nucleic acids II: the polymerase chain reaction. 834
In: Hillis, D.M., Moritz, C., Mable, B.K. (Eds.), *Molecular System-*
835 *atics*. Sinauer Associates, Sunderland, MA, pp. 205–247. 836
- Perron, F.E., (1981). The partitioning of reproductive energy between 837
ova and protective capsules in marine gastropods of the genus 838
Comus. *Am. Nat.* 118, 110–118. 839
- Posada, D., Crandall, K.A., (1998). Modeltest: testing the model of 840
DNA substitution. *Bioinformatics* 14, 817–818. 841
- Rouse, G., Fitzhugh, K., (1994). Broadcasting fables: Is external fertil- 842
ization really primitive. Sex, size, and larvae in sabellid polychaetes. 843
Zool. Scripta 23, 271–312. 844
- Scouras, A., Beckenbach, K., Arndt, A., Smith, M.J., (2004). Complete 845
mitochondrial genome DNA sequence for two ophiuroids and a 846
holothuroid: the utility of protein gene sequence and gene maps in 847
the analyses of deep deuterostome phylogeny. *Mol. Phylogenet. Evol.* 848
31, 50–65. 849

- 850 Selvakumaraswamy, P., Byrne, M., (2000). Reproduction, spawning, 870
851 and development of 5 ophiuroids from Australia and New Zealand. 871
852 *Invertebr. Biol.* 119, 394–402. 872
- 853 Smith, A.B., Littlewood, D.T.J., Wray, G.A., (1995). Comparing pat- 873
854 terns of evolution: Larval and adult life history stages and ribo- 874
855 somal RNA of post-Palaeozoic echinoids. *Phil. Trans. R. Soc.*
856 *Lond. B* 349, 11–18. 875
- 857 Sponer, R., Deheyn, D., Roy, M.S., (2001). Large genetic distances 876
858 within a population of *Amphipholis squamata* (Echinodermata; 877
859 Ophiuroidea) do not support colour varieties as sibling species. 878
860 *Mar. Ecol. Prog. Ser.* 219, 169–175. 879
- 861 Strathmann, M.F., (1987). *Reproduction and Development of Marine* 880
862 *Invertebrates of the Northern Coast.* University of Washington 881
863 Press, Seattle. 1115–1116. 882
- 864 Strathmann, R.R., (1985). Feeding and nonfeeding larval development 883
865 and life-history evolution in marine invertebrates. *Ann. Rev. Ecol.* 884
866 *Syst.* 16, 339–361. 885
- 867 Swofford, D.L., 2002. PAUP*: Phylogenetic analysis using parsim- 886
868 ony (* and other methods). Version 4.0b10. Sinauer, Sunder- 887
869 land, MA. 50, 408–424. 888 889
- Villinski, J.T., Villinski, J.C., Byrne, M., Raff, R.A., (2002). Convergent 870
maternal provisioning and life-history evolution in echinoderms. 871
Evolution 56, 1764–1775. 872
- Wada, H., Komatsu, M., Satoh, N., (1996). Mitochondrial rDNA phy- 873
logeny of the Asteroidea suggests the primitiveness of the Paxillo- 874
sida. *Mol. Phylogenet. Evol.* 6, 97–106. 875
- Wiens, J.J., (1998). Does adding characters with missing data increase 876
or decrease phylogenetic accuracy?. *Syst. Biol.* 47 (4), 625–640. 877
- Wray, G.A., (1995a). Evolution of larvae and developmental modes. In: 878
McEdward, L.R. (Ed.), *Ecology of Marine Invertebrate Larvae.* 879
CRC Press, New York, pp. 413–447. 880
- Wray, G.A., (1995b). Punctuated evolution of embryos. *Science* 267, 881
1115–1116. 882
- Wray, G.A., (1996). Parallel evolution of nonfeeding larvae in echi- 883
noids. *Syst. Biol.* 45, 308–322. 884
- Wray, G.A., Raff, R.A., (1991). The evolution of developmental strat- 885
egy in marine invertebrates. *Trends Ecol. Evol.* 6, 45–50. 886
- Yoder, A.D., Irwin, J.A., Payseur, B.A., (2001). Failure of the ILD to 887
determine data combinability for slow loris phylogeny. *Syst. Biol.* 888
50, 408–424. 889