

RESEARCH ARTICLE

Control of Movement

## Using asymmetry to your advantage: learning to acquire and accept external assistance during prolonged split-belt walking

Natalia Sánchez,<sup>1</sup>  Surabhi N. Simha,<sup>2</sup> J. Maxwell Donelan,<sup>2</sup> and James M. Finley<sup>1,3,4</sup>

<sup>1</sup>Division of Biokinesiology and Physical Therapy, University of Southern California, Los Angeles, California; <sup>2</sup>Department of Biomedical Physiology and Kinesiology, Simon Fraser University, Burnaby, British Columbia, Canada; <sup>3</sup>Department of Biomedical Engineering, University of Southern California, Los Angeles, California; and <sup>4</sup>Neuroscience Graduate Program, University of Southern California, Los Angeles, California

### Abstract

People can learn to exploit external assistance during walking to reduce energetic cost. For example, walking on a split-belt treadmill affords the opportunity for people to redistribute the mechanical work performed by the legs to gain assistance from the difference in belts' speed and reduce energetic cost. Though we know what people should do to acquire this assistance, this strategy is not observed during typical adaptation studies. We hypothesized that extending the time allotted for adaptation would result in participants adopting asymmetric step lengths to increase the assistance they can acquire from the treadmill. Here, participants walked on a split-belt treadmill for 45 min while we measured spatiotemporal gait variables, metabolic cost, and mechanical work. We show that when people are given sufficient time to adapt, they naturally learn to step further forward on the fast belt, acquire positive mechanical work from the treadmill, and reduce the positive work performed by the legs. We also show that spatiotemporal adaptation and energy optimization operate over different timescales: people continue to reduce energetic cost even after spatiotemporal changes have plateaued. Our findings support the idea that walking with symmetric step lengths, which is traditionally thought of as the endpoint of adaptation, is only a point in the process by which people learn to take advantage of the assistance provided by the treadmill. These results provide further evidence that reducing energetic cost is central in shaping adaptive locomotion, but this process occurs over more extended timescales than those used in typical studies.

**NEW & NOTEWORTHY** Split-belt treadmill adaptation can be seen as a process where people learn to acquire positive work from the treadmill to reduce energetic cost. Though we know what people should do to reduce energetic cost, this strategy is not observed during adaptation studies. We extended the duration of adaptation and show that people continuously adapt their gait to acquire positive work from the treadmill to reduce energetic cost. This process requires longer exposure than traditionally allotted.

*locomotion; mechanical work; metabolic cost; motor adaptation; split-belt walking*

### INTRODUCTION

Humans frequently adapt their locomotor patterns to take advantage of potential sources of external assistance. For example, we use the pull on the leash from our dog to propel us as we walk down the street, and we use gravity when walking downhill to reduce the mechanical work performed by our muscles and reduce metabolic cost (1). In other settings, we actively adjust our position in the environment to gain external assistance. During surfing, we kick and paddle to position our bodies and the surfboard to allow the waves

to propel us. During sailing, we adjust the alignment of the sails relative to the wind to propel us through the water. Based on this tendency for the neuromotor system to take advantage of external assistance, several devices have been used to provide a source of external assistance and give the user the opportunity to reduce the work performed by the legs and the energetic cost of walking (2–9).

The success of assistive devices for reducing energetic cost depends greatly on the neuromotor system's ability to adapt coordination to take advantage of assistance. First, the person must sense the points during the gait cycle when the



device is providing assistance. Then, in some cases, the person must adjust their coordination patterns to acquire the assistance (8). Here, we define acquire assistance as letting the device perform work on the body. However, acquiring the assistance does not necessarily mean that the person will accept and use this assistance. The person must accept the assistance provided by the external system by adapting the amplitude and timing of muscle activity at specific points in the gait cycle to reduce muscular work that may be unnecessary given the assistance from the device. This reduction in muscle work will then lead to reductions in metabolic cost. These adjustments must take place while continuing to achieve high-level objectives, such as maintaining a steady walking pace when walking on a treadmill. Understanding how people learn to take advantage of assistance can provide novel insights into the process by which people optimize energetic cost during locomotion.

We recently demonstrated that a common task used to study locomotor learning, split-belt treadmill adaptation (10, 11), provides an opportunity where people can learn to take advantage of external assistance provided by the treadmill. Specifically, people can *acquire* assistance in the form of net positive work from the treadmill if they redistribute braking and propulsive forces between the two belts such that the leg on the fast belt produces most of the braking and the leg on the slow belt produces most of the propulsion (12). The person can then *accept* the assistance from the treadmill and reduce the positive work performed by the person's muscles, either by redistributing the propulsive forces between the two legs, by reducing total propulsive force, or by applying the propulsive force at different stages in the gait cycle, to ultimately reduce metabolic cost. We empirically tested these predictions and showed that when people use guided feedback to increase braking by stepping further forward with the leg on the fast belt, the treadmill performs net positive work on the person. This strategy results in asymmetric step lengths, where fast and slow step lengths are defined as the fore-aft distance between the feet at the respective foot strike. Thus, we observed that walking with asymmetric step lengths, where the leg on the fast belt takes steps that are longer than the leg on the slow belt, was accompanied by a reduction in the positive work generated by the legs and an associated reduction in metabolic cost (12). These findings complement previous studies showing that people reduce the mechanical work generated by the fast leg (13, 14) and the metabolic cost of walking (15, 16) as they adapt to the split-belt treadmill.

Although people can take advantage of the positive work performed by the treadmill by adopting asymmetric step lengths, the vast majority of studies of split-belt adaptation have reported that adaptation ends with people taking steps of nearly equal length (11, 17–25). This idea that the goal of adaptation is reaching symmetric step lengths is consistent with the theory that in response to a perturbation, individuals adapt their movement to minimize their sensory prediction error (26, 27), or the discrepancy between the expected and the actual sensory consequences of a movement. Thus, reduction of step length asymmetry is a proxy for the error minimization that occurs during adaptation (11, 23). Another theory is that individuals adapt their gait pattern to minimize energetic cost (15, 16, 28, 29). However, as we previously demonstrated, energetic cost is lowest when walking with

asymmetric step lengths (12, 30). Given that the vast majority of research has reported that individuals adapt toward symmetric step lengths, this would seemingly suggest that people prioritize symmetry over energetic cost.

We recently showed that after people are provided with guided experience of less energetically costly step length asymmetries, they adopt an asymmetric gait pattern when they are allowed to adapt freely. This pattern consists of longer step lengths with the leg on the fast belt (12), which is consistent with the gait that minimizes energetic cost during split-belt walking. It remains to be determined why people do not typically walk with longer steps with the leg on the fast belt during typical adaptation experiments. One potential explanation is that conventional adaptation studies may simply be too short in duration to allow self-exploration to solve the problem of learning to take advantage of external assistance. Consistent with this idea, people took longer steps with the leg on the fast belt after 45 min of guided experience walking on the treadmill (12), which is three to four times longer than the length of conventional adaptation studies. People also take longer steps with the leg on the fast belt after multiple sessions of adaptation (31), and even over 5 days (32), totaling approximately 1 h and 15 min of experience. Thus, the adoption of a gait with longer steps with the leg on the fast belt, which is less energetically costly, seems to occur after combined exposure times that are several multiples of the traditional 10–15 min allotted for adaptation.

Energy optimization during motor learning is a complex problem, which often requires extended experience. Although we can reduce energetic cost over short timescales commonly used in studies of adaptation (15, 16, 33–36), energetic cost can be further reduced over multiple sessions of experience (8, 37, 38). Adjusting movement patterns to reduce energetic cost poses a complex problem for the neuromotor system. Specifically, during split-belt walking, people can reduce energy by learning to acquire external assistance in the form of positive work from the treadmill and then accept this assistance by reducing the positive work by the legs. This process requires adjusting interlimb coordination to redistribute the amount of work generated by each leg as well as adjusting the time during the gait cycle when the legs generate work (13, 14). To remain in place on the treadmill, the net mechanical work on the person must be zero on average. Therefore, if the person adopts a gait pattern that leads to an increase in the amount of positive work by the treadmill on the body, this must be balanced in one of three ways. The person can either dissipate the energy transferred to the body by increasing the negative work by the legs, reduce the positive work generated by the legs, or use a combination of both strategies. The person can fulfill this requirement with one leg or with both legs, and they could use these strategies at different points of the gait cycle. The challenge of both *acquiring* and *accepting* assistance in a way that leads to a net energetic benefit might explain why previous studies have not observed longer steps with the leg on the fast belt—in these studies, learning may simply have been incomplete.

In this study, we have two goals. The first goal is to understand whether extending the duration allowed for adaptation leads people to adopt asymmetric step lengths during split-belt walking. The second goal is to study the time

course of how people learn to acquire positive work from the treadmill and the time course of how they learn to accept that work by reducing the positive work done by the legs. We hypothesized that extending the duration of experience on a split-belt treadmill would result in participants adopting asymmetric step lengths to increase the assistance they can acquire from the treadmill. During the extended exposure time, participants would also learn to accept this assistance and reduce the positive work generated by the legs. We also hypothesized that continued reductions in positive work by the legs would lead to continued reductions in metabolic cost beyond the cost of walking with steps of equal length. If our findings are consistent with these predictions, this will support the idea that walking with symmetric step lengths is only a point in the process by which people learn to take advantage of the assistance provided by the treadmill during split-belt adaptation. Importantly, this would suggest that locomotor adaptation requires significantly longer than 10–15 min of experience, likely because of the challenge associated with learning to take advantage of assistance. Finally, if participants can adjust their coordination to take advantage of external assistance and reduce the work generated by the legs, this would provide additional support for the role of energy optimization in shaping adaptation of locomotor behaviors.


## MATERIALS AND METHODS

### Experiment Design

A convenience sample of 15 healthy young adults participated in our study. Exclusion criteria included a history of lower extremity surgery or orthopedic injury within the past 2 years. All participants were naïve to the split-belt protocol, and right leg dominant, as assessed by self-reports. The University of Southern California Institutional Review Board approved all experimental procedures, and each participant provided written informed consent before testing began. All aspects of the study conformed to the principles described in the Declaration of Helsinki. All data were collected at the Locomotor Control Lab at the University of Southern California.

We calibrated the motion capture system immediately before each experiment per manufacturer specifications (Qualisys AB, Goteborg, Sweden), including zeroing of the force plates embedded in the dual belt treadmill (Fully Instrumented Treadmill, Bertec Corporation, Ohio). We allowed the metabolic cart used to measure metabolic cost to warm up for 30 min and then we calibrated it per manufacturer specifications (Parvomedics, Utah).

Participants walked under three different speed conditions (Fig. 1): tied belts baseline trial with both belts moving at 1.0 m/s for 6 min, a split-belt adaptation trial for 45 min, and a tied-belts washout trial with both belts moving at 1.0 m/s for 10 min. During the split-belt trial, we set the speed of the left belt at 1.5 m/s and the speed of the right leg at 0.5 m/s in a 3:1 ratio (Fig. 1C). We set the duration of the split-belt trial to 45 min to match the cumulative duration of our previous study (12), which would allow us to determine whether the less energetically costly gait pattern with asymmetric step lengths we previously observed was a consequence only of guided experience with a more economical



**Figure 1.** Experimental design. **A:** experimental setup. Participants walked on an instrumented, dual-belt treadmill with reflective markers placed bilaterally on the lateral malleoli and greater trochanters. **B:** top view of the experimental setup and the leading and trailing limb placement relative to the body that make up individual step lengths. **C:** experimental protocol and treadmill speeds for each trial. Participants rested for at least 4 min between all walking trials, to ensure metabolic cost was at standing baseline conditions before starting a new trial. **D:** raw values of step lengths calculated using the distance between markers on the lateral malleoli for a single participant during all walking trials. Vertical dashed lines indicate different walking trials. Orange: step lengths at heel strike on the fast belt. Blue: step lengths at heel strike on the slow belt. **E:** raw values of net metabolic power for a single participant. Vertical dashed lines indicate different walking trials.

gait or due to prolonged walking time. We collected data continuously for the entirety of the 45-min trial. Participants did not take any breaks during the adaptation trial.

During all walking trials, participants wore a harness designed to prevent falls while providing no body weight

support. No handrails were accessible during the experiment, and participants were instructed to avoid holding on to the harness. After each walking trial, participants rested for at least 4 min, and we used real-time measures of metabolic cost to ensure that participants' metabolic cost returned to resting levels before beginning the next walking trial.

### Data Acquisition

We recorded the positions of reflective markers located bilaterally on the lateral malleoli and greater trochanters at 100 Hz (Fig. 1A). We also recorded ground reaction forces generated by each leg at 1000 Hz to calculate mechanical power and work. We assessed metabolic cost by determining the rates of oxygen consumption ( $\dot{V}O_2$ ) and carbon dioxide production ( $\dot{V}CO_2$ ) using a TrueOne 2400 metabolic cart (Parvomedics, Utah; Fig. 1E). The metabolic cart sampled gas concentrations on a breath-by-breath basis.

### Data Processing and Analysis

We used custom-written code in MATLAB R2019b (Mathworks, Natick, MA) for all data processing and analyses. A fourth-order low-pass digital Butterworth filter smoothed marker data and ground reaction force data. The filter cutoff frequency for marker data was set to 10 Hz and the filter cutoff frequency for force data was set to 20 Hz.

We used the positions of markers on the lateral malleoli (Fig. 1A) to estimate step length asymmetry as follows (11, 23, 39):

$$\text{Step Length Asymmetry} = \frac{SL_{fast} - SL_{slow}}{SL_{fast} + SL_{slow}} \quad (1)$$

Here,  $SL_{fast}$  is the step length at leading leg heel strike on the fast belt and  $SL_{slow}$  is the step length at leading leg heel strike on the slow belt (Fig. 1, B–D). Negative values correspond to longer steps with the slow (right) leg and positive values correspond to longer steps with the fast (left) leg. Step lengths are the product of adjusting the leading and trailing limb placement at heel strike and toe off, respectively. Limb placement will affect the fore-aft component of the ground reaction force and the amount of work the legs must generate. Therefore, we calculated leading and trailing limb placement for each limb as the fore-aft distance relative to the midpoint between markers placed bilaterally on the greater trochanters (Fig. 1B), to determine whether individuals adjusted limb placement in a manner consistent with taking advantage of the work by the treadmill.

We also calculated stance, swing, and double support times from marker data (40). Swing time corresponds to the time between toe off, which was estimated as the most posterior location of the ankle markers, and heel strike on the same side, which was estimated as the most anterior location of the ankle marker. Stance time corresponds to the time between heel strike and toe off on the same side. Finally, double support time for a given limb corresponds to the time from contralateral heel strike to ipsilateral toe off.

We estimated the mechanical work performed by the legs using an extension of the individual limbs method (12–14, 41, 42). This method considers a person as a point mass body with legs that are massless pistons. The legs can generate

forces on the ground while simultaneously generating equal but opposite forces on the point mass, referred to as the center of mass. Thus, we assume that all the work performed by the person is being performed by the legs. To calculate mechanical work in our experiments, we segmented force data into strides using a vertical ground reaction force threshold of 32 N to identify foot strike (13) and then calculated the center of mass acceleration in the fore-aft, vertical, and medio-lateral directions, as the sum of the forces within a stride normalized by body mass. We then calculated the center of mass velocities in each direction as the time integral of the accelerations. We determined the integration constant for each stride by forcing the average velocity over the stride to be zero in each direction. All velocities and accelerations were expressed relative to a reference frame attached to the fixed ground. Next, we calculated the instantaneous power generated by each leg for each stride as the instantaneous sum of the dot product of the ground reaction force and the center of mass velocity and the dot product of the force applied to the respective belt and the belt speed. Finally, we calculated the total positive and total negative work performed by each leg as the time integral of the positive or negative portion of the total instantaneous power over the stride cycle. To have a comparable interpretation with metabolic power, we express all measures of work as work rate by dividing each measure by stride duration.

We estimated the energy consumed using the standard Brockway equation (43) as follows:

$$E_{met,gross} = \left( 16.48 \frac{J}{ml O_2} V_{O_2} \cdot 1000 \right) + \left( 4.48 \frac{J}{ml CO_2} V_{CO_2} \cdot 1000 \right) \quad (2)$$

From here, we divided  $E_{met,gross}$  by the exact duration (T) over which it was calculated to obtain an estimate of the gross metabolic rate  $P_{met,gross}$  measured in Watts. Finally, we subtracted each participants' standing metabolic rate from each walking trial (Fig. 1E). Thus, all metabolic rate values presented here are net metabolic rate. For each participant, we estimated metabolic cost as the average net metabolic rate of the 3 min preceding the timepoint of interest. This corresponded to the last 3 min of the standing baseline and baseline walking trials, and *min 12–15* and *min 42–45* of adaptation.

Changes in step length asymmetry during conventional studies of split-belt adaptation occur over two distinct time-scales (23, 25, 44). However, it is unclear if two distinct time-scales are necessary to model adaptation when the adaptation period is extended to 45 min. Therefore, we modeled step length asymmetry (SLA) as a function of stride number (s) for the adaptation and washout trials using either a single exponential (Eq. 3) or the sum of two exponentials (Eq. 4).

$$SLA(s) = a \times e^{-s/b} + c \quad (3)$$

$$SLA(s) = a_{fast} \times e^{-s/b_{fast}} + a_{slow} \times e^{-s/b_{slow}} + c \quad (4)$$

In the single exponential model,  $a$  corresponds to the step length asymmetry at the onset of the perturbation ( $s = 0$ ), and  $b$  is the rate constant parameter. Based on our hypothesis,

SLA would plateau at positive values; therefore, we added a term  $c$  that corresponds to the step length asymmetry as  $s \rightarrow \infty$ . For the two-exponential model,  $a_{fast}$  and  $a_{slow}$  are the initial values of the fast and slow exponentials, respectively, and  $b_{fast}$  and  $b_{slow}$  are the fast and slow rate constants, respectively. We fit one and two exponential models to step length asymmetry data for the group as a whole by concatenating step length asymmetry vectors for all participants. We then used the Akaike information criterion (AIC) (45) to compare whether the more complex two-exponential model provided a better fit to the data than a single exponential model. We selected the model with the lowest AIC.

We used the exponential model selected above and ran bootstrap analyses to obtain the 95% confidence intervals (CIs) of the adaptation model parameters. We created 10,000 bootstrapped samples of 15 participants by sampling participants with replacement. For each bootstrapped sample, we concatenated the SLA vectors for all participants and fit the data to the selected model using `fitnlm` in MATLAB. This process was repeated 10,000 times. We sorted all model parameters in ascending order and identified the 500th and 9,500th values as the limits of the 95% CI.

We modeled positive work rate by the fast leg as a function of stride number ( $s$ ) for the adaptation trials using either a single exponential (Eq. 3) or the sum of two exponentials (Eq. 4). We used the AIC to select the exponential model that fit the individual data and ran bootstrap analyses to obtain the distribution and 95% confidence intervals for the model parameters and  $R^2$  using the same approach for bootstrap analyses described in the previous paragraph for step length asymmetry.

We also determined how the duration of adaptation influenced the magnitude and washout of aftereffects associated with adaptation. We compared step length asymmetry for the first five strides of washout, which corresponds to early postadaptation (11). We analyzed washout data from our study and a previous study of 12 participants where they adapted for 15 min (46, 47). We implemented the models in Eqs. 3 and 4 and the procedure described for fitting step length asymmetry data during adaptation to determine whether a one or two exponential model provided a better fit to the washout data. We also ran bootstrap analysis for the data from the washout period to obtain the distribution of model parameters.

## Statistical Analyses

Our goal was to test the hypothesis that prolonged exposure to split-belt walking leads individuals to take longer steps with the leg on the fast belt, resulting in more positive step length asymmetries, increased positive work performed by the treadmill, reductions in positive work rate by the legs, and reductions in metabolic cost beyond those observed in traditional 15-min adaptation experiments. Based on the Kolmogorov–Smirnov test, all data were normally distributed. Therefore, we used paired samples, one-tailed  $t$  tests for step lengths, step length asymmetry, foot placement relative to the body, stance, swing and double support times, and the positive and negative mechanical work performed by the legs to test for differences between 15 and at 45 min of adaptation. For each variable, we averaged the values over the last 100 strides and the 100 strides before the 15-min time-point to obtain values for statistical analyses.

We hypothesized that individuals would adopt positive step length asymmetries to gain assistance from the treadmill, which would allow them to reduce the positive work by the legs and reduce metabolic cost. Therefore, we computed correlations between changes in step length asymmetry ( $\Delta SLA_{45-0}$ ), changes in foot placement and heel strike, and changes in positive work rate by the fast leg ( $\Delta W_{45-0}^+$ ) during adaptation to determine whether people decreased positive work by the fast leg as step length asymmetry becomes more positive. Similarly, we computed the correlation between the change in work ( $\Delta W_{45-0}^+$ ) and the change in metabolic cost ( $\Delta MetCost_{45-0}$ ) during adaptation because we hypothesized that people would reduce energetic cost as they reduced positive work. We also computed the correlation between changes in step length asymmetry ( $\Delta SLA_{45-0}$ ) and the change in metabolic cost ( $\Delta MetCost_{45-0}$ ). Given the rapid reductions in step length asymmetry that occur at initial exposure to the split-belt treadmill, for correlation analyses, we calculated average step length asymmetry and work only for the first and last five strides (11). For metabolic cost, we used data averaged over 1 min for the early and late adaptation periods. The metabolic cost for early adaptation was averaged over *min* 3 and *min* 4 of split-belt walking to account for transport lag and temporal dynamics of the change in metabolic cost in response to changes in exercise intensity (16, 48). We used the Pearson correlation coefficient, as data were normally distributed. We computed the effect size of the difference between rate constants of the exponential fits for step length asymmetry and positive work rate by the fast leg using Cohen's  $d$  (49, 50) using the distribution of rate constants obtained in the bootstrap analyses.


We compared step length asymmetry during early postadaptation after 45 and 15 min of adaptation using independent samples  $t$  tests. We also computed the effect size of the difference between model parameters for washout data after 45 min and after 15 min of adaptation using Cohen's  $d$  (49, 50) using the distribution of rate constants derived from bootstrap analyses.

## RESULTS

Fifteen individuals participated in our study (nine female, six male, age  $27 \pm 7$  yr, weight  $62 \pm 11$  kg, height  $168 \pm 9$  cm). One participant terminated the experiment after the split-belt adaptation condition, but we included their data for all other experimental conditions.

### After Prolonged Adaptation, People Overshoot Symmetry and Adopt Positive Step Length Asymmetries

We hypothesized that extending the duration of experience on a split-belt treadmill would allow participants to step further forward with the leg on the fast belt, generate positive step length asymmetries, and allow the treadmill to generate net positive work on the person. In agreement with our hypothesis, participants continued to lengthen the step length on the fast belt, with an average increase of 18 mm from 15 to 45 min of adaptation (Fig. 2, A–C, paired  $t$  test, one tail, 95% CI  $> 7.942$ ,  $P = 0.003$ ), whereas the step length on the slow belt did not change systematically from 15 to 45 min of adaptation (Fig. 2, D and E,  $P = 0.481$ ). After 45 min of adaptation, step length asymmetry was on average 0.018 more positive compared with 15 min (Fig. 2, F–H, paired  $t$  test, one tail, 95%



**Figure 2.** Participants continuously modified step lengths with the fast leg over 45 min of adaptation. **A:** fast (orange) and slow (blue) step lengths during adaptation. Solid lines represent sample means. Shaded areas are standard deviations. After 45 min of adaptation, the step length on the fast belt was longer than that on the slow belt ( $P = 0.003$ ). **B:** average step lengths on the fast belt at 15 and 45 min for all participants. **C:** 95% CI of the differences in fast step lengths at 45 min compared with 15 min. **D:** average step lengths on the slow belt at 15 and 45 min for all participants. **E:** 95% CI of the differences in slow step lengths at 45 min compared with 15 min. **F:** mean step length asymmetry during adaptation (black). Gray-shaded areas are standard deviations. **G:** average step length asymmetry at 15 and 45 min for all participants. **H:** 95% CI of the differences in step length asymmetry at 45 min compared with 15 min. Subscripts 15 and 45 refer to the specific timepoints used for analyses. Averages in **B**, **D**, and **G** were obtained for 100 strides before the identified timepoint. CI, confidence interval; SL, step lengths; SLA, step length asymmetry.


CI > 0.001,  $P = 0.042$ ). At 45 min, step length asymmetry was significantly more positive than baseline, with an average of 0.030 (Fig. 2F, one-sample  $t$  test, 95% CI > 0.004,  $P = 0.020$ ). Step length asymmetry data for individual participants are included in Supplemental Fig. S1 (all Supplemental material is available at <https://doi.org/10.17605/OSF.IO/WRSU8>).

Participants could take longer steps with the leg on the fast belt by either allowing the leg on the slow belt to trail further back or by placing the leg on the fast belt further forward at heel strike. Based on our hypothesis, placing the leg further forward on the fast belt would increase the fore-aft component of the ground reaction force, allowing the treadmill to perform more positive work on the body. In agreement with this hypothesis, participants continued to adjust fast leg leading placement from 15 to 45 min by 13 mm on average (Fig. 3, A, D, and E, paired  $t$  test, one tail, 95% CI > 6.464,  $P = 0.003$ ). Neither slow leg leading or trailing placement relative to the body (Fig. 3, C, F, G, J, and K) nor fast leg trailing placement relative to the body changed systematically (Fig. 3, H and I). We excluded one participant from this analysis due to loss of greater trochanter marker data.

Adaptation of step lengths occurred with a simultaneous adaptation of stance, swing, and double support times. However, each of these temporal variables adapted over different timescales. Participant's double support time plateaued quickly, with no changes in fast ( $P = 0.500$ ) or slow ( $P = 0.915$ ) double-support times between 15 and 45 min. In contrast, participants' stance time on the slow leg increased from 15 to 45 min by  $0.03 \pm 0.05$  s (paired  $t$  test, one tail, 95% CI > 0.001,  $P = 0.044$ ), whereas stance time on the fast leg remained unchanged from 15 to 45 min ( $P = 0.080$ ). Accordingly, swing time on the fast leg was  $0.02 \pm 0.04$  s longer at 45 min compared to 15 min (CI > 0.001,  $P = 0.049$ ), whereas swing time on the slow leg remained unchanged ( $P = 0.155$ ). This indicates that participants also continue to adjust step timing over the prolonged split-belt adaptation.

### Participants Quickly Accept Positive Work from the Treadmill and Gradually Reduce Positive Work by the Legs

We hypothesized that prolonged exposure to the split-belt treadmill would allow participants to learn to gain positive work from the treadmill to reduce positive work by the legs.




**Figure 3.** Modifications in foot placement relative to the body. We present data for 14 participants, as one participant lost a marker by the end of the trial. **A:** mean and standard deviation of leading (orange) and trailing (blue) limb placement relative to the body on the fast belt. The increase in step length by the fast leg was due to increased forward placement of the leading limb at heel strike. **B:** bird's-eye view of a participant as they take a step with the leg on the slow belt. The top and bottom dashed lines correspond to the location of the leading and trailing feet in **C** and **A**, respectively. The opposite would apply for a fast step. **C:** mean and standard deviation of leading (blue) and trailing (orange) limb placement relative to the body for steps on the slow belt. The difference in the values of the orange curves in **A** and **C** corresponds to the fast step length. The difference in the values of the blue curves in **C** and **A** correspond to the slow step length. **D:** fast limb placement relative to the body at heel strike at 15 and 45 min. **E:** 95% CI of the change in fast limb distance to the body at heel strike from 15 to 45 min. **F:** slow limb placement relative to the body at 15 and 45 min. **G:** 95% CI of the change in slow limb distance to the body at heel strike from 15 to 45 min. **H:** fast limb placement relative to the body at toe off at 15 and 45 min. **I:** 95% CI of the change in fast limb distance to the body at toe off from 15 to 45 min. **J:** slow limb placement to the body at toe off 15 and 45 min. **K:** 95% CI of the change in slow limb distance to the body at toe off from 15 to 45 min. Averages in **D**, **F**, **H**, and **J** were obtained for 100 strides. CI, confidence interval.

Consistent with our hypothesis, participants performed net negative work rate with the legs by the end of adaptation with an average of  $-0.06$  W/kg (one-sample  $t$  test, one tail, 95% CI  $< -0.0438$ ,  $P = 3.720 \times 10^{-6}$ ). Participants increased negative work rate with the fast leg within the first hundreds of strides (Fig. 4A), with no significant changes in negative work rate from 15 min of adaptation to 45 min of adaptation for the fast leg (Figs. 4A and 5A, paired  $t$  test,  $P = 0.094$ ). The negative work rate by the slow leg did not change systematically between 15 min and 45 min of adaptation (Fig. 4B and 5B,  $t$  test,  $P = 0.087$ ).

To take advantage of the assistance provided by the treadmill, it is not enough to only increase negative work rate by the legs. Participants must also learn to decrease


positive work rate, which could ultimately lead to a reduction in metabolic cost. Learning to reduce this positive work rate by the legs also begins quickly (Fig. 4C), with an average reduction from initial exposure to 15 min of adaptation of  $0.20$  W/kg (95% CI  $> 0.16$ ,  $P = 1.14 \times 10^{-6}$ ), or  $42 \pm 14\%$  for the fast leg. This learning process continues throughout the entire experiment, with an average reduction in positive work rate performed by the fast leg of  $11 \pm 9\%$ , which corresponds to  $0.025$  W/kg from 15 to 45 min (Figs. 4C and 5C, paired  $t$  test, 95% CI  $> 0.019$ ,  $P < 1.48 \times 10^{-4}$ ). Note that the reduction in positive work rate by the fast leg was not complete at the end of the experiment, as evidenced by the presence of a negative slope in the work rate time series. The positive work rate by the slow leg



**Figure 4.** Positive and negative work rate by the legs during adaptation. *A*: negative work rate generated by the fast leg. *B*: negative work rate generated by the slow leg. *C*: positive work rate by the fast leg. Participants continued to reduce positive work rate by the fast leg during the adaptation trial. *D*: positive work rate by the slow legs. Shaded areas are standard deviations.

stabilized almost immediately, with no changes between initial exposure and 15 min (Fig. 4D, *t* test,  $P = 0.215$ ) or between 15 and 45 min (Figs. 4D and 5D,  $P = 0.109$ ). Positive work rate data for the fast leg for individual participants are included in Supplemental Fig. S2.

We hypothesized that adoption of a more positive step length asymmetry would be associated with a reduction in positive work by the fast leg. In agreement with this hypothesis, we observed that changes in positive work rate by the fast leg ( $\Delta W_{45-15}^+$ ) were negatively correlated with changes in step



**Figure 5.** Changes in mechanical work rate by the fast and slow leg during adaptation. *A*: negative work rate by the fast leg at 15 and 45 min. *B*: 95% CI of the change in negative work rate from 15 to 45 min. *C*: negative work rate by the slow leg at 15 and 45 min. *D*: 95% CI of the change in negative work rate by the slow leg from 15 to 45 min. *E*: positive work rate by the fast leg at 15 and 45 min. *F*: 95% CI of the change in positive work rate by the fast leg from 15 to 45 min. *G*: positive work rate by the slow leg at 15 and 45 min. *H*: 95% CI of the change in slow limb positive work rate from 15 to 45 min.  $W^+$  refers to positive work.  $W^-$  refers to negative work. Subscripts 15 and 45 refer to the specific timepoints used for analyses. Averages in *A*, *C*, *E*, and *G* were obtained for 100 strides. CI, confidence interval.

length asymmetry during adaptation ( $\Delta SLA_{45-0}$ ;  $r = -0.64$ ,  $P = 0.010$ , Fig. 6E) but were not correlated with changes in fast leg leading limb placement ( $r = -0.51$ ,  $P = 0.060$ , Fig. 6F).


### Metabolic Cost Continues to Decrease as Participants Learn to Take Advantage of Assistance from the Treadmill

We hypothesized that continuous reductions in the positive work rate by the legs over 45 min of adaptation would lead to reductions in the metabolic cost of walking beyond the cost of walking with steps of equal lengths. Consistent with this prediction, participants reduced their metabolic cost by an additional 7% (IQR 8%), which corresponds to a reduction of 0.17 W/kg from 15 to 45 min of walking (Fig. 6, A–C, paired  $t$  test, 95% CI  $> 0.059$ ,  $P = 0.007$ ). The lower metabolic cost observed at 45 min indicates that people continue to refine their gait pattern to increase economy despite the demands of walking on a split-belt treadmill continuously for 45 min. Metabolic power data for individual participants are included in Supplemental Fig. S3.

In agreement with our hypothesis that walking with more positive step length asymmetry is less energetically costly, we found a negative correlation between the change in asymmetry ( $\Delta SLA_{45-0}$ ) and the change in metabolic cost over the course of adaptation ( $\Delta MetCost_{45-0}$ ,  $r = -0.66$ ,  $P = 0.026$ , Fig. 6G). The change in metabolic cost measured during adaptation ( $\Delta MetCost_{45-0}$ ) was also positively correlated with the change in positive work rate by the fast leg,  $\Delta W_{45-0}$  ( $r = 0.52$ ,  $P = 0.048$ , Fig. 6D). Thus, individuals reduced the positive work rate by the legs when obtaining assistance provided by the treadmill, and this ultimately led to reductions in metabolic cost.

### Adaptation of Step Length Asymmetry Can Be Characterized by a Sum of Two Exponential Models That Plateau at Positive Step Length Asymmetries

Changes in step length asymmetry during conventional studies of split-belt adaptation occur over two distinct timescales (23, 25, 44). Similarly, we found that a sum of two exponentials provided a good fit to step length



**Figure 6.** Metabolic power continuously decreased during adaptation. **A:** average metabolic power during adaptation trial. Shaded area represents the standard deviation across participants. **B:** average metabolic cost at 15 min and at 45 min of adaptation. Averages for metabolic data were obtained across 3-min periods. **C:** reduction in metabolic cost from 15 to 45 min of adaptation. **D:** correlation between the change in positive work by the fast leg from initial exposure to 45 min and change in metabolic power from initial exposure to 45 min. **E:** correlation between the change in step length asymmetry and positive work by the fast leg from initial exposure to 45 min. **F:** correlation between the change in fast leg foot placement relative to the body at heel strike and changes in work rate by the fast leg was not significant. **G:** correlation between the change in step length asymmetry and metabolic power from initial exposure to 45 min.  $W^+$  refers to positive work. Met refers to metabolic power. Subscripts 0, 15, and 45 refer to the specific timepoints used for analyses. Shaded areas in correlation plots are 95% CI. CI, confidence interval; SLA, step length asymmetry.

asymmetry data during prolonged adaptation (Eq. 4). To obtain the confidence intervals for the model parameters, we ran bootstrap analyses and obtained an  $R^2$  for the two exponential model of 0.356, bootstrap 95% CI [0.265, 0.447]. The model parameters were as follows:  $a_{fast} = -0.31$  (95% CI [-0.39, -0.24]),  $a_{slow} = -0.15$  (95% CI [-0.20, -0.11]),  $b_{fast} = 20$  strides (95% CI [9, 35]), and  $b_{slow} = 347$  strides (95% CI [195, 553]). Finally, the step length asymmetry plateau,  $c$ , was equal to 0.033 (95% CI [0.015, 0.053]), which supports our hypothesis that participants adapt toward positive asymmetries. Based on this model, it would take  $\sim 1,564$  strides (95% CI [1,089, 2,060]) to adapt 95% of the difference between the initial step length asymmetry during early adaptation and the final plateau. This is on average twice as long as traditional adaptation experiments. Note that the average duration of our study was  $\sim 2,300$  strides (range between 2,066 and 2,853 strides for all 15 participants).


### Adaptation of Positive Work Rate by the Fast Leg Can Be Characterized by a Single Exponential Model

We fit positive work rate for the fast leg using a single exponential model (Eq. 3). The exponential model had an  $R^2$  of 0.07, bootstrap 95% CI [0.03, 0.11]. The model parameters were as follows:  $a = 0.11$  (95% CI [0.07, 0.25]),  $b = 452$  strides (95% CI [10, 1,192]), and plateau,  $c$ , was equal to 0.24 W/kg (95% CI [0.21, 0.26]). The time constant for work rate by the fast leg was longer than the slow rate constant for step length asymmetry with a large effect size, as shown by a Cohen's  $d$  of 7.4, indicating that adaptation of work rate toward steady state occurs in a longer timescale than adaptation of step length asymmetry.

### The Duration of Washout Is Proportional to the Duration of Adaptation

We expected that when the belts returned to the same speed, individuals would have greater retention of the strategy adopted after 45 min of adaptation compared with 15 min, indicated by both a larger initial step length asymmetry and a longer time constant for washout when participants walked with belts tied at 1.0 m/s. We did not observe differences in step length asymmetry during early postadaptation after 45 min ( $0.34 \pm 0.15$ ) or 15 min ( $0.32 \pm 0.08$ ) of split-belt walking (Fig. 7, A and B, independent samples  $t$  test,  $P = 0.758$ ). However, the time course of washout varied with the duration of adaptation. A single exponential model (Eq. 3) best characterized the time course of changes in step length asymmetry during washout both after 15 and 45 min of adaptation.

We ran bootstrap analyses to obtain the confidence intervals of the washout model parameters both after 45 and 15 min of adaptation. We observed that the  $R^2$  values were 0.48 (95% CI [0.41, 0.56]) after 45 min of adaptation and 0.60 (95% CI [0.54, 0.66]) after 15 min of adaptation. The means and 95% confidence intervals of the model parameters were (Fig. 7, B–D) as follows:  $a_{45} = 0.22$  (95% CI [0.18, 0.26]),  $a_{15} = 0.27$  (95% CI [0.22, 0.31]),  $b_{45} = 63$  strides (95% CI [41, 82]),  $b_{15} = 25$  strides (95% CI [17, 33]),  $c_{45} = 0.05$  (95% CI [0.035, 0.066]), and  $c_{15} = 0.038$  (95% CI [0.034, 0.043]). Participants required almost three times longer to washout



**Figure 7.** Step length asymmetry (SLA) during washout trial. **A:** step length asymmetry time series. Data in purple were collected in  $n = 14$  participants who completed all walking trials of our study protocol. Data in green were collected from a previous study following a 15-min adaptation period in  $n = 12$  healthy participants (46). Shaded areas are standard deviations. **B–D:** distribution of the bootstrapped parameters for the single exponential model coefficients  $a$ ,  $b$ , and  $c$  as defined in Eq. 3.

the learned gait pattern after 45 min versus 15 min of adaptation (Fig. 7, A and C). The effect size for the difference in initial asymmetry ( $a_{45}$  vs.  $a_{15}$ ) was small, with a Cohen's  $d$  of 0.02. The difference in washout rates ( $b_{45}$  vs.  $b_{15}$ ) had a large effect size, with a Cohen's  $d$  of 13. Finally, the differences in the plateau ( $c_{45}$  vs.  $c_{15}$ ) had a small effect size, with a Cohen's  $d$  of 0.14.

## DISCUSSION

Learning to take advantage of external assistance from the environment during walking poses a complex problem for the neuromotor system. Here, we asked how people acquire and accept assistance during adaptation to walking on a split-belt treadmill. Acquiring work from the treadmill and then accepting this work requires coordination between limbs and across different parts of the gait cycle, all while maintaining a steady pace to avoid drifting back or walking off the treadmill. We found that during adaptation, participants adopt asymmetric step lengths to acquire positive work from the treadmill and reduce the positive work by the legs. This reduction in positive work by the legs was associated with continuous reductions in metabolic cost over the course of 45 min of walking. Our findings support the idea that energetic cost, which can be reduced when learning to

use external assistance, plays a critical role in shaping locomotor adaptation to a prolonged split-belt perturbation.

Previous studies have considered adaptation of locomotor patterns during split-belt walking to be a process driven by minimization of sensory prediction error, which is complete once participants achieve symmetric step lengths (11, 17, 19, 20, 23, 25, 38). This is largely because people transition from a gait with marked step length asymmetries to taking steps of nearly equal lengths after 10–15 min of adaptation. We can draw some insights from previous studies in arm reaching adaptation, which have found that minimization of error and effort may occur over different timescales, with error minimization being faster and taking preference over minimization of effort, which occurs over a longer timescale (26, 33, 51, 52). Here we find that adaptation of step length asymmetry is indeed faster than minimization of energetic cost. However, we find that individuals do not minimize step length asymmetry but adapt it toward a positive value consistent with taking advantage of assistance. We also show that adaptation of step length asymmetry, work, and reduction of metabolic cost continue beyond 15 min of adaptation. Thus, we conclude that minimizing step length asymmetry, which was traditionally thought of as the goal of adaptation, is perhaps better viewed as an initial point along the slower path toward a less energetically costly gait.

Although it may seem intuitive that external assistance leads to a reduction in metabolic cost, providing participants with a source of external positive work does not imply that they will know how to use it for assistance. For example, *participant 15* in our sample did not change their step lengths during adaptation (Supplemental Fig. S1), and they maintained the same rate of positive work with the fast leg through the entire 45 min of adaptation (Supplemental Fig. S2). Consequently, they did not reduce metabolic cost during adaptation (Supplemental Fig. S3). In participants who do learn how to take advantage of assistance, they learn to acquire positive work from the treadmill by first increasing forward limb placement of the fast leg, which increases negative work by the fast leg, and this occurs within hundreds of strides. Then, they accept the positive work by the treadmill by reducing the positive work by the leg on the fast belt, but this process requires thousands of strides and had not ended even after 45 min. The time during the gait cycle when participants generate positive work is also crucial for reducing energetic cost (41, 42). Participants adjust the time when they generate positive work from single limb support during early adaptation to the step-to-step transition during late adaptation (13). Generating positive work during the step-to-step transition at the same time when negative work is generated is less energetically costly, as it decreases the cost of redirecting the body velocity (42). This learning problem of coordinating multiple degrees of freedoms and coordinating spatiotemporal, kinematic, and kinetic processes to take advantage of external assistance appears to be complex enough that it requires the neuromotor system even longer than 45 min to solve.

How long is required for full adaptation? Based on our two exponential model for step length asymmetry, step length asymmetry is near the predicted plateau after 45 min of split-belt walking. However, adaptation of positive work rate did not achieve a stable plateau even after 45 min. Thus, adaptation of step length asymmetry does not mean that the

entire adaptation process is complete even after  $\sim 2,300$  strides. Although we cannot speculate on the amount of time required to minimize positive work rate, we know that it is markedly longer than is traditionally allotted. This is similar to what has been observed during adaptation of reaching (53): here, we find that the slow rate constant for step length asymmetry and the rate constant for positive work indicate that the time required for behavior to plateau is longer than what is traditionally allotted experimentally. This does not imply that adaptation cannot occur rapidly with other experimental manipulations. There is evidence that people adopt longer steps with the leg on the fast belt when walking on an inclined split-belt treadmill for 10 min (54) or when adapting to belt speeds close to running for 5 min (55). Given that these studies did not calculate mechanical work, and we do not know what the energetically optimal solution for the above studies would be, we cannot speculate if the rate of adaptation toward the energy optimal behavior occurred more rapidly than in the current study.

We found different timescales for adaptation of step length asymmetry and positive work rate by the fast leg empirically and using exponential models. One might wonder why step length asymmetry, work rate by the fast leg, and metabolic cost are not all synchronized in time and modeled by the same sum of two exponentials. This could result from a combination of measurement limitations and behavioral strategies. For example, metabolic cost often changes over a slower time-scale than changes in the underlying mechanical behavior largely as a result of the temporal dynamics of blood circulation and substrate utilization and lag in measurement of metabolic cost using expired gas analysis (48, 56, 57). Furthermore, one can modify step length asymmetry without necessarily reducing the work by the legs (as is the case for *participant 4*, Supplemental Figs. S1 and S2). One can also acquire work from the treadmill without reducing metabolic cost by not learning to reduce muscle activation or by generating excessive cocontraction. Therefore, learning to take advantage of assistance to ultimately reduce energetic cost is a complex problem, given the high dimensionality of the space of potential muscle activation patterns and the interplay between spinal-level pattern generating circuits, brainstem contributions, and supraspinal control (58). Together, these results support our general conclusion that adapting step length asymmetry and learning to take advantage of the assistance provided by the treadmill occur over multiple timescales.

One remaining question is whether individuals could further reduce the positive work performed by the legs. A previous study used computer modeling (59) to show that a dynamic walker (60, 61) can passively walk on a split-belt treadmill by harnessing energy from the treadmill, demonstrating that simple bipeds can walk with zero positive work. In addition, we previously established that participants could theoretically use a split-belt treadmill moving at a 3:1 ratio with an effectiveness of 67% (12), given that the trailing leg on the slow belt would need to perform positive work that is roughly one-third of the negative work performed by the leading leg on the fast belt. Here, we found that participants used the work by the treadmill with an effectiveness of 28%. The observation that the theoretically achievable effectiveness is  $\sim 40\%$  greater than what was observed during prolonged

exposure suggests that a gait that further reduces work rate and, potentially, metabolic cost could exist. Whether people can achieve this level of effectiveness given task and anatomical constraints has yet to be determined.

We found that the amount of experience did not affect the amount of retention of step length asymmetry, as indicated by the similar step length asymmetries measured during early postadaptation following 15 and 45 min of adaptation. That the magnitude of step length asymmetry at early postadaptation does not depend on prior exposure duration might be explained by task constraints: an initial asymmetry of  $\sim 0.30$  with belts tied might be the largest asymmetry possible to avoid falling, walking off, or drifting on the treadmill, but this remains to be determined. However, prolonged experience was associated with longer time constants for wash-out of the adapted pattern. Previous studies in reaching (62) and walking (63) have observed similar results for both the magnitude of the aftereffects and the duration of washout. After reaching in a force field, errors during error-clamp trials were similar after short versus long adaptation, yet errors persisted longer after adapting for more trials (62). These findings combined with our own might be explained by a two-state model of motor learning (64), where the time course of the decay of a motor memory depends on the number of adaptation trials, which here would correspond to the number of strides during adaptation. Thus, prolonged wash-out may reflect the state of the slower components of the adaptation process that are stabilized with practice.

Several findings in our study provide novel insights into the field of assistive devices in general and lower limb exoskeletons in particular. Similar to the split-belt treadmill, lower limb exoskeletons assist locomotion by performing mechanical work for the user, and the user has to learn to accept this assistance to reduce the work performed by muscles (3, 6–9, 65). As shown by two participants who did not reduce work by the fast leg, we demonstrate that just because a device performs work on the person, it does not imply that the person will learn how to reduce the work performed by the legs. This is similar to findings showing that people's ability to accept assistance from exoskeletons is modest compared with predictions of exoskeleton performance (66). Second, we show that participants continuously adapt their gait to reduce the work performed by the fast leg, and this process appears to continue beyond the 45 min of exposure that we provided. We previously found that visual feedback can be used to effectively guide individuals to take advantage of the assistance from the treadmill (12). Therefore, in the absence of explicit guidance, training periods for the use of assistive devices might require longer exposures than traditionally allotted to determine the set of parameters necessary to minimize energetic cost. Finally, our work demonstrates that split-belt treadmills, which are common in many biomechanics laboratories, can be used to study the principles underlying the processes by which people learn to use assistive devices.

## CONCLUSIONS

Prolonged adaptation to walking on a split-belt treadmill leads to continuous and gradual spatiotemporal, kinematic, and kinetic changes over different timescales, which result

in a reduction in positive work by the legs and reduced energetic cost compared with what is traditionally observed after 15 min of adaptation. These findings demonstrate that participants learn to take advantage of the assistance provided by the treadmill, but this process requires much longer than is traditionally allotted during studies of adaptation. Learning to take advantage of external assistance is a complex problem since participants must first determine how to acquire assistance, and then learn how to take advantage of this assistance to reduce energetic cost. Thus, providing participants with enough time is crucial for the neuromotor system to adjust coordination and adapt the many different processes required to be able to exploit external assistance and reduce energetic cost. Other factors, such as a desire to maintain balance (15), may be important objectives during early phases of adaptation. However, given the continuous reductions in work and energetic cost observed here, which continue even after adaptation of step lengths and step times has plateaued, our results are consistent with the theory that people seek to minimize energetic cost when faced with continuous perturbations during walking.

## ACKNOWLEDGMENTS

Preprint is available at <https://doi.org/10.1101/2020.04.04.025619>.

## GRANTS

This work was funded by the NIH National Center for Advancing Translational Science (KL2TR001854) to N. Sánchez, the NIH National Institute of Child Health and Human Development (R01-HD091184) to J. M. Finley, and a National Sciences and Engineering Research Council Discovery Grant to J. M. Donelan.

## DISCLOSURES

No conflicts of interest, financial or otherwise, are declared by the authors.

## AUTHOR CONTRIBUTIONS

N.S. and J.M.F. conceived and designed research; N.S. performed experiments; N.S. analyzed data; N.S., S.N.S., J.M.D., and J.M.F. interpreted results of experiments; N.S. prepared figures; N.S. drafted manuscript; N.S., S.N.S., J.M.D., and J.M.F. edited and revised manuscript; N.S., S.N.S., J.M.D., and J.M.F. approved final version of manuscript.

## ENDNOTE

At the request of the authors, readers are herein alerted to the fact that additional materials related to this manuscript may be found at <https://osf.io/wrsu8/>. These materials are not a part of this manuscript, and have not undergone peer review by the American Physiological Society (APS). APS and the journal editors take no responsibility for these materials, for the website address, or for any links to or from it.

## REFERENCES

- Hunter LC, Hendrix EC, Dean JC. The cost of walking downhill: is the preferred gait energetically optimal? *J Biomech* 43: 1910–1915, 2010. doi:10.1016/j.jbiomech.2010.03.030.

2. **Collins SH, Wiggan MB, Sawicki GS.** Reducing the energy cost of human walking using an unpowered exoskeleton. *Nature* 522: 212–215, 2015. doi:10.1038/nature14288.
3. **Ding Y, Kim M, Kuindersma S, Walsh CJ.** Human-in-the-loop optimization of hip assistance with a soft exosuit during walking. *Sci Robot* 3: eaar5438, 2018. doi:10.1126/scirobotics.aar5438.
4. **Kang I, Hsu H, Young A.** The effect of hip assistance levels on human energetic cost using robotic hip exoskeletons. *IEEE Robot Autom Lett* 4: 430–437, 2019. doi:10.1109/LRA.2019.2890896.
5. **Kim J, Lee G, Heimgartner R, Revi DA, Karavas N, Nathanson D, Galiana I, Eckert-Erdheim A, Murphy P, Perry D, Menard N, Choe DK, Malcolm P, Walsh CJ.** Reducing the metabolic rate of walking and running with a versatile, portable exosuit. *Science* 365: 668–672, 2019. doi:10.1126/science.aav7536.
6. **Malcolm P, Derave W, Galle S, De Clercq D.** A simple exoskeleton that assists plantarflexion can reduce the metabolic cost of human walking. *PLoS One* 8: e56137, 2013. doi:10.1371/journal.pone.0056137.
7. **Sawicki GS, Beck ON, Kang I, Young AJ.** The exoskeleton expansion: improving walking and running economy. *J Neuroeng Rehabil* 17: 25, 2020. doi:10.1186/s12984-020-00663-9.
8. **Sawicki GS, Ferris DP.** Mechanics and energetics of level walking with powered ankle exoskeletons. *J Exp Biol* 211: 1402–1413, 2008. doi:10.1242/jeb.009241.
9. **Zhang J, Fiers P, Witte KA, Jackson RW, Poggensee KL, Atkeson CG, Collins SH.** Human-in-the-loop optimization of exoskeleton assistance during walking. *Science* 356: 1280–1284, 2017. doi:10.1126/science.aal5054.
10. **Dietz V, Zijlstra W, Duysens J.** Human neuronal interlimb coordination during split-belt locomotion. *Exp Brain Res* 101: 513–520, 1994. doi:10.1007/BF00227344.
11. **Reisman DS, Block HJ, Bastian AJ.** Interlimb coordination during locomotion: what can be adapted and stored? *J Neurophysiol* 94: 2403–2415, 2005. doi:10.1152/jn.00089.2005.
12. **Sánchez N, Simha SN, Donelan JM, Finley JM.** Taking advantage of external mechanical work to reduce metabolic cost: the mechanics and energetics of split-belt treadmill walking. *J Physiol* 15: 1–44, 2019. doi:10.1113/JP277725.
13. **Selgrade BP, Thajchayapong M, Lee GE, Toney ME, Chang Y-H.** Changes in mechanical work during neural adaptation to asymmetric locomotion. *J Exp Biol* 220: 2993–2300, 2017. doi:10.1242/jeb.149450.
14. **Selgrade BP, Toney ME, Chang YH.** Two biomechanical strategies for locomotor adaptation to split-belt treadmill walking in subjects with and without transbilateral amputation. *J Biomech* 53: 136–143, 2017. doi:10.1016/j.jbiomech.2017.01.012.
15. **Buurke TJ, Lamoth CJ, Vervoort D, van der Woude LH, den Otter R.** Adaptive control of dynamic balance in human gait on a split-belt treadmill. *J Exp Biol* 221: jeb.174896, 2018. doi:10.1242/jeb.174896.
16. **Finley JM, Bastian AJ, Gottschall JS.** Learning to be economical: the energy cost of walking tracks motor adaptation. *J Physiol* 591: 1081–1095, 2013. doi:10.1113/jphysiol.2012.245506.
17. **Day KA, Leech KA, Roemmich RT, Bastian AJ.** Accelerating locomotor savings in learning: compressing four training days to one. *J Neurophysiol* 119: 2100–2113, 2018. doi:10.1152/jn.00903.2017.
18. **Hoogkamer W, Brujin SM, Potocanac Z, Van Calenbergh F, Swinnen SP, Duysens J.** Gait asymmetry during early split-belt walking is related to perception of belt speed difference. *J Neurophysiol* 114: 1705–1712, 2015. doi:10.1152/jn.00937.2014.
19. **Leech KA, Roemmich RT.** Independent voluntary correction and savings in locomotor learning. *J Exp Biol* 221: jeb.181826, 2018. doi:10.1242/jeb.181826.
20. **Long AW, Roemmich RT, Bastian AJ.** Blocking trial-by-trial error correction does not interfere with motor learning in human walking. *J Neurophysiol* 115: 2341–2348, 2016. doi:10.1152/jn.00941.2015.
21. **Malone LA, Bastian AJ, Torres-Oviedo G.** How does the motor system correct for errors in time and space during locomotor adaptation? *J Neurophysiol* 108: 672–683, 2012. doi:10.1152/jn.00391.2011.
22. **Mawase F, Haizler T, Bar-Haim S, Karniel A.** Kinetic adaptation during locomotion on a split-belt treadmill. *J Neurophysiol* 109: 2216–2227, 2013. doi:10.1152/jn.00938.2012.
23. **Mawase F, Bar-Haim S, Shmuelof L.** Formation of long-term locomotor memories is associated with functional connectivity changes in the cerebellar–thalamic–cortical network. *J Neurosci* 37: 349–361, 2017. doi:10.1523/JNEUROSCI.2733-16.2016.
24. **Roemmich RT, Long AW, Bastian AJ.** Seeing the errors you feel enhances locomotor performance but not learning. *Curr Biol* 26: 2707–2716, 2016. doi:10.1016/j.cub.2016.08.012.
25. **Torres-Oviedo G, Bastian AJ.** Seeing is believing: effects of visual contextual cues on learning and transfer of locomotor adaptation. *J Neurosci* 30: 17015–17022, 2010. doi:10.1523/JNEUROSCI.4205-10.2010.
26. **Izawa J, Shadmehr R.** Learning from sensory and reward prediction errors during motor adaptation. *PLoS Comput Biol* 7: e1002012–11, 2011. doi:10.1371/journal.pcbi.1002012.
27. **Shadmehr R, Smith MA, Krakauer JW.** Error correction, sensory prediction, and adaptation in motor control. *Annu Rev Neurosci* 33: 89–108, 2010. doi:10.1146/annurev-neuro-060909-153135.
28. **Emken JL, Benitez R, Sideris A, Bobrow JE, Reinkensmeyer DJ.** Motor adaptation as a greedy optimization of error and effort. *J Neurophysiol* 97: 3997–4006, 2007. doi:10.1152/jn.01095.2006.
29. **Zarrugh MY, Todd FN, Ralston HJ.** Optimization of energy expenditure during level walking. *Eur J Appl Physiol Occup Physiol* 33: 293–306, 1974. doi:10.1007/BF00430237.
30. **Sánchez N, Park S, Finley JM.** Evidence of energetic optimization during adaptation differs for metabolic, mechanical, and perceptual estimates of energetic cost. *Sci Rep* 7: 7682, 2017. doi:10.1038/s41598-017-08147-y.
31. **Charalambous CC, French MA, Morton SM, Reisman DS.** A single high-intensity exercise bout during early consolidation does not influence retention or relearning of sensorimotor locomotor long-term memories. *Exp Brain Res* 237: 2799–2810, 2019. doi:10.1007/s00221-019-05635-7.
32. **Leech KA, Day KA, Roemmich RT, Bastian AJ.** Movement and perception recalibrate differently across multiple days of locomotor learning. *J Neurophysiol* 120: 2130–2137, 2018. doi:10.1152/jn.00355.2018.
33. **Huang HJ, Kram R, Ahmed AA.** Reduction of metabolic cost during motor learning of arm reaching dynamics. *J Neurosci* 32: 2182–2190, 2012. doi:10.1523/JNEUROSCI.4003-11.2012.
34. **Huang HJ, Ahmed AA.** Reductions in muscle coactivation and metabolic cost during visuomotor adaptation. *J Neurophysiol* 112: 2264–2274, 2014. doi:10.1152/jn.00014.2014.
35. **Selinger JC, O'Connor SM, Wong JD, Donelan JM.** Humans can continuously optimize energetic cost during walking. *Curr Biol* 25: 1–5, 2015. doi:10.1016/j.cub.2014.10.064.
36. **Selinger JC, Wong JD, Simha SN, Donelan JM.** How humans initiate energy optimization and converge on their optimal gaits. *J Exp Biol* 222: jeb.198234, 2019. doi:10.1242/jeb.198234.
37. **Lay BS, Sparrow WA, Hughes KM, O'Dwyer NJ.** Practice effects on coordination and control, metabolic energy expenditure, and muscle activation. *Hum Mov Sci* 21: 807–830, 2002. doi:10.1016/S0167-9457(02)00166-5.
38. **Sparrow WA, Newell KM.** Energy expenditure and motor performance relationships in humans learning a motor task. *Psychophysiology* 31: 338–346, 1994. doi:10.1111/j.1469-8986.1994.tb02442.x.
39. **Choi JT, Vining EPG, Reisman DS, Bastian AJ.** Walking flexibility after hemispherectomy: split-belt treadmill adaptation and feedback control. *Brain* 132: 722–733, 2009. doi:10.1093/brain/awn333.
40. **Zeni JA, Richards JG, Higginson JS.** Two simple methods for determining gait events during treadmill and overground walking using kinematic data. *Gait Posture* 27: 710–714, 2008. doi:10.1016/j.gaitpost.2007.07.007.
41. **Donelan JM, Kram R, Kuo AD.** Simultaneous positive and negative external mechanical work in human walking. *J Biomech* 35: 117–124, 2002. doi:10.1016/S0021-9290(01)00169-5.
42. **Donelan JM, Kram R, Kuo AD.** Mechanical work for step-to-step transitions is a major determinant of the metabolic cost of human walking. *J Exp Biol* 205: 3717–3727, 2002. ]
43. **Brockway JM.** Derivation of formulae used to calculate energy expenditure in man. *Hum Nutr Clin Nutr* 41: 463–471, 1987. ]
44. **Darmohray DM, Jacobs JR, Marques HG, Carey MR.** Spatial and temporal locomotor learning in mouse cerebellum. *Neuron* 102: 217–231.e4, 2019. doi:10.1016/j.neuron.2019.01.038.
45. **Akaike H.** Likelihood of a model and information criteria. *J Econom* 16: 3–14, 1981. doi:10.1016/0304-4076(81)90071-3.

46. **Park S, Finley JM.** Characterizing dynamic balance during adaptive locomotor learning. *Proc Annu Int Conf IEEE Eng Med Biol Soc EMBS* 50–53, 2017. doi:10.1109/EMBC.2017.8036760.
47. **Park S, Finley JM.** Manual stabilization reveals a transient role for balance control during locomotor adaptation (Preprint). *bioRxiv* 647453, 2019. doi:10.1101/647453.
48. **Selinger JC, Donelan JM.** Estimating instantaneous energetic cost during non-steady-state gait. *J Appl Physiol* 117: 1406–1415, 2014. doi:10.1152/jappphysiol.00445.2014.
49. **Cohen J.** Statistical power analysis [Online]. *Curr Dir Psychol Sci* 1: 98–101, 1992. doi:10.1111/1467-8721.ep10768783.
50. **Lakens D.** Calculating and reporting effect sizes to facilitate cumulative science: a practical primer for t-tests and ANOVAs. *Front Psychol* 4: 863, 2013. doi:10.3389/fpsyg.2013.00863.
51. **Balasubramanian R, Howe RD, Matsuoka Y.** Task performance is prioritized over energy reduction. *IEEE Trans Biomed Eng* 56: 1310–1317, 2009. doi:10.1109/TBME.2008.2006683.
52. **Scheidt RA, Reinkensmeyer DJ, Conditt MA, Rymer WZ, Mussa-Ivaldi FA.** Persistence of motor adaptation during constrained, multi-joint, arm movements. *J Neurophysiol* 84: 853–862, 2000. doi:10.1152/jn.2000.84.2.853.
53. **Maeda RS, Cluff T, Gribble PL, Pruszynski JA.** Feedforward and feedback control share an internal model of the arm's dynamics. *J Neurosci* 38: 10505–10514, 2018. doi:10.1523/JNEUROSCI.1709-18.2018.
54. **Sombric CJ, Calvert JS, Torres-Oviedo G.** Large propulsion demands increase locomotor learning at the expense of step length symmetry. *Front Physiol* 10: 60, 2019. doi:10.3389/fphys.2019.00060.
55. **Yokoyama H, Sato K, Ogawa T, Yamamoto S-I, Nakazawa K, Kawashima N.** Characteristics of the gait adaptation process due to split-belt treadmill walking under a wide range of right-left speed ratios in humans. *PLoS One* 13: e0194875, 2018. doi:10.1371/journal.pone.0194875.
56. **Turner DL.** Cardiovascular and respiratory control mechanisms during exercise: an integrated view. *J Exp Biol* 160: 309–340, 1991.
57. **Whipp BJ, Ward SA.** Physiological determinants of pulmonary gas exchange kinetics during exercise. *Med Sci Sports Exerc* 22: 62–71, 1990.
58. **Dietz V.** Human neuronal control of automatic functional movements: interaction between central programs and afferent input. *Physiol Rev* 72: 33–69, 1992. doi:10.1152/physrev.1992.72.1.33.
59. **Simha SN, Chui VL, Butterfield JK, Donelan JM, Collins SH.** Passive machines on a split-belt treadmill. *Dynamic Walking*. Canmore, Alberta, June 3–6, 2019.
60. **Kuo AD.** A simple model of bipedal walking predicts the preferred speed-step length relationship. *J Biomech Eng* 123: 264–269, 2001. doi:10.1115/1.1372322.
61. **McGeer T.** Passive dynamic walking. *Int J Rob Res* 9: 62–82, 1990. doi:10.1177/027836499000900206.
62. **Huang VS, Shadmehr R.** Persistence of motor memories reflects statistics of the learning event. *J Neurophysiol* 102: 931–940, 2009. doi:10.1152/jn.00237.2009.
63. **Roemmich RT, Bastian AJ.** Two ways to save a newly learned motor pattern. *J Neurophysiol* 113: 3519–3530, 2015. doi:10.1152/jn.00965.2014.
64. **Smith MA, Ghazizadeh A, Shadmehr R.** Interacting adaptive processes with different timescales underlie short-term motor learning. *PLoS Biol* 4: e179, 2006. doi:10.1371/journal.pbio.0040179.
65. **Galle S, Malcolm P, Collins SH, De Clercq D.** Reducing the metabolic cost of walking with an ankle exoskeleton: interaction between actuation timing and power. *J Neuroeng Rehabil* 14: 1–16, 2017. doi:10.1186/s12984-016-0214-x.
66. **Handford ML, Srinivasan M.** Robotic lower limb prosthesis design through simultaneous computer optimizations of human and prosthesis costs. *Sci Rep* 6: 19983, 2016. doi:10.1038/srep19983.