

Classical Mechanics

Lecture 7

UNIT 10: WORK AND ENERGY

Today's Concepts:
Work & Kinetic Energy



"Knowing is not enough; we must apply. Willing is not enough; we must do."

Bruce Lee

SCIENCE SPOOKTACULAR



featuring

WICKED SCIENCE ACTIVITIES!

**SPOOKTACULAR SHOWS OF
PHANTOM PHYSICS & CRYPTIC CHEMISTRY!**



Free Family Fun at Simon Fraser University

**Halloween Saturday
Saturday October 25, 2014**

Demos and activities 12:00-3:30
Academic Quadrangle, South Concourse 3000 level

Spooktacular shows 1:00 and 3:00
Shrum Science Centre, B9200 Lecture Theatre
Priority tickets available free (see website below)

Information: <http://www.sfu.ca/physics/outreach/science-spooktacular.html>

Volunteer: <http://www.sfu.ca/~exafs/spookvolunteer.html>



Costumes welcomed!

Karate



Will not do Session 3 of Unit 10.

It is a Karate thing.

You can do it as a “project” in Unit 14 instead of the assigned project on the pendulum.



Stuff you asked about:

WHAT IS GOING ON WITH ALL THE INTEGRALS?????? HELP :(

Why is it necessary to use integrals in the equation for work?

everything. everything is difficult

Everything is fine except the work done by Gravity far from earth

The slides had a lot of information that was confusing and it flew by. I didn't have time to understand what was going on.

The dot product is confusing


Pretty much everything. This prelecture sort of went over my head. I'll definitely have to watch it again.

The concept of positive and negative work is very confusing. please explain

Uh oh. Hotdog.

I hope you have enough funny quotes to fit into your slides.

The Dot Product



Dot Product

$$\vec{A} \cdot \vec{B} \equiv AB \cos \theta$$

$$\vec{A} \cdot \vec{B} = AB \quad (\text{Parallel Vectors})$$

$$\vec{A} \cdot \vec{B} = 0 \quad (\text{Perpendicular Vectors})$$

$$\vec{A} \cdot \vec{B} = -AB \quad (\text{Anti-Parallel Vectors})$$

Dot Product using Unit Vectors

$$\hat{i} \cdot \hat{i} = 1$$

$$\hat{i} \cdot \hat{j} = 0$$

$$\hat{j} \cdot \hat{j} = 1$$

$$\hat{j} \cdot \hat{k} = 0$$

$$\hat{k} \cdot \hat{k} = 1$$

$$\hat{k} \cdot \hat{j} = 0$$

etc.

$$\vec{A} = A_x \hat{i} + A_y \hat{j} + A_z \hat{k}$$

$$\vec{B} = B_x \hat{i} + B_y \hat{j} + B_z \hat{k}$$

Dot Product in Rectangular Coordinates

$$\begin{aligned}\vec{A} \cdot \vec{B} &= (A_x \hat{i} + A_y \hat{j} + A_z \hat{k}) \cdot (B_x \hat{i} + B_y \hat{j} + B_z \hat{k}) \\&= (A_x B_x \hat{i} \cdot \hat{i} + A_x B_y \hat{i} \cdot \hat{j} + \cdots + A_z B_y \hat{k} \cdot \hat{j} + A_z B_z \hat{k} \cdot \hat{k}) \\&= (A_x B_x \times 1 + \cancel{A_x B_y \times 0} + \cdots + \cancel{A_z B_y \times 0} + A_z B_z \times 1) \\ \vec{A} \cdot \vec{B} &= (A_x B_x + A_y B_y + A_z B_z)\end{aligned}$$

Finding the angle between vectors

If we know the xyz components of two vectors we can find the angle between them:

$$\begin{aligned}\vec{A} \cdot \vec{B} &= (A_x B_x + A_y B_y + A_z B_z) \\ &= |\vec{A}| |\vec{B}| \cos \theta\end{aligned}$$

therefore

$$\theta = \arccos \left(\frac{\vec{A} \cdot \vec{B}}{|\vec{A}| |\vec{B}|} \right)$$

Example

$$\vec{A} = 3\hat{i} + 4\hat{j}$$

$$|\vec{A}| = |\vec{B}| = 5$$

$$\vec{B} = 4\hat{i} + 3\hat{j}$$

$$\vec{A} \cdot \vec{B} = 3 \cdot 4 + 4 \cdot 3 = 24$$

$$\theta = \arccos \left(\frac{\vec{A} \cdot \vec{B}}{|\vec{A}| |\vec{B}|} \right)$$

$$\theta = \arccos \left(\frac{24}{25} \right) = \arccos 0.96$$
$$= 0.28 \text{ rad} = 16^\circ$$

Work-Kinetic Energy Theorem


The work done by the net force F as it acts on an object that moves between positions r_1 and r_2 is equal to the change in the object's kinetic energy:

$$W = \Delta K$$

$$W = \int_{\vec{r}_1}^{\vec{r}_2} \vec{F} \cdot d\vec{l} \qquad K = \frac{1}{2}mv^2$$

Work-Kinetic Energy Theorem: 1-D Example

If the force is constant and the directions aren't changing then this is very simple to evaluate:



$$W = \int_{\vec{r}_1}^{\vec{r}_2} \vec{F} \cdot d\vec{l} = \vec{F} \cdot \vec{d}$$

In this case $= Fd$ since $\cos(0)=1$


$$\left(\int_{\vec{r}_1}^{\vec{r}_2} d\vec{l} = \vec{d} \right)$$

This is probably what you remember from High School.
(The notation may be confusing though.)

Clicker Question



A lighter car and a heavier van, each initially at rest, are pushed with the same constant force F . After both vehicles travel a distance d , which of the following statements is true? (Ignore friction)




- A) They will have the same velocity
- B) They will have the same kinetic energy
- C) They will have the same momentum

$$\int_{\vec{r}_1}^{\vec{r}_2} \vec{F} \cdot d\vec{l} = \Delta K$$


Derivation – not so important

Concept – very important




A force pushing over some distance
will change the kinetic energy.

Work done by gravity near the Earth's surface




Work done by gravity near the Earth's surface

$$W_{TOT} = W_1 + W_2 + \dots + W_N$$
$$= m\vec{g} \cdot d\vec{l}_1 + m\vec{g} \cdot d\vec{l}_2 + \dots + m\vec{g} \cdot d\vec{l}_N$$



Work done by gravity near the Earth's surface

$$\begin{aligned}W_{TOT} &= W_1 + W_2 + \dots + W_N \\&= m\vec{g} \cdot d\vec{l}_1 + m\vec{g} \cdot d\vec{l}_1 + \dots + m\vec{g} \cdot d\vec{l}_N \\&= -mgdy_1 - mgdy_2 \dots - mgdy_N \\&= -mg\Delta y\end{aligned}$$



$$W_g = -m g \Delta y$$

Work-Kinetic Energy Theorem


If there are several forces acting then W is the work done by the net (total) force:

$$W_{NET} = \Delta K$$

$$= W_1 + W_2 + \dots$$


You can just add up the work done by each force

$$W_{NET} = W_{TOT}$$




CheckPoint

Falling Sliding and Swinging: Question 1 (N = 54)



Three objects having the same mass begin at the same height, and all move down the same vertical distance H . One falls straight down, one slides down a frictionless inclined plane, and one swings on the end of a string.



In which case does the object have the biggest net work done on it by all forces during its motion?

Free Fall

Frictionless incline

String

A) Free Fall

B) Incline

C) String


D) All the same

Clicker Question



Three objects having the same mass begin at the same height, and all move down the same vertical distance H . One falls straight down, one slides down a frictionless inclined plane, and one swings on the end of a string.

What is the relationship between their speeds when they reach the bottom?




A) $v_f > v_i > v_p$

B) $v_f > v_p > v_i$

C) $v_f = v_p = v_i$

CheckPoint / Clicker Question



A) $v_f > v_i > v_p$

B) $v_f > v_p > v_i$

C) $v_f = v_p = v_i$

Only gravity will do work: $W_g = \Delta K$

$$\left\{ \begin{array}{l} W_g = mgH \\ \Delta K = \frac{1}{2} mv_f^2 \end{array} \right\} \quad v = \sqrt{2gH}$$



$$v_f = v_i = v_p = \sqrt{2gH}$$


CheckPoint

A car drives up a hill with constant speed. Which statement best describes the total work W_{TOT} done on the car by all forces as it moves up the hill?

A) $W_{TOT} = 0$

B) $W_{TOT} > 0$

C) $W_{TOT} < 0$



Less than 40% got this right...

CheckPoint





A car drives up a hill with constant speed.
Which statement best describes the
total work W_{TOT} done on the car by all
forces as it moves up the hill?

A) $W_{TOT} = 0$

B) $W_{TOT} > 0$

C) $W_{TOT} < 0$



A) The change in kinetic energy is zero because the velocity is constant.
By the work-kinetic energy theorem, we know that work must also be zero.

B) The force of friction between the car's wheels and the ramp exert a force up the
ramp over a particular distance. $W=Fd$, so the work is positive.


C) gravity is downward and car moves up the hill.

Clicker Question




A car drives up a hill with constant speed. How does the kinetic energy of the car change as it moves up the hill?

- A) It increases
- B) It stays the same
- C) It decreases



CheckPoint

A box sits on the horizontal bed of a moving truck. Static friction between the box and the truck keeps the box from sliding around as the truck drives.




The work done on the box by the static frictional force as the truck moves a distance D is:

- A) Positive B) Negative C) Zero


About 60% got this right...

From Before


A box sits on the horizontal bed of a moving truck. Static friction between the box and the truck keeps the box from sliding around as the truck drives.




If the truck moves with constant accelerating to the left as shown, which of the following diagrams best describes the static frictional force acting on the box:



CheckPoint



Work on Box in Accelerating Truck: Question (N = 53)



The work done on the box by the static frictional force as the truck moves a distance D is:

A) Positive B) Negative C) Zero



A) because the direction of the static frictional force and the direction the box travels are all same.

B) Since the friction force always has opposite direction from the movement direction, the work done is negative


C) Friction keeps the box from sliding, thus $D=0$, and work done is 0.

Work done by a Spring

$$W_{1 \rightarrow 2} = -\frac{1}{2}k(x_2^2 - x_1^2)$$



I am confused about the positive work and negative work and also the positive and negative forces for the spring problems.



Use the formula to get the magnitude of the work

Use a picture to get the sign (look at directions)


$$W_{1 \rightarrow 2} = -\frac{1}{2}k(x_2^2 - x_1^2)$$

In this example the spring does –ve work since F and Δx are in opposite direction.
The axes don't matter.

Clicker Question



A box attached at rest to a spring at its equilibrium length. You now push the box with your hand so that the spring is compressed a distance D , and you hold the box at rest in this new location.



During this motion, the spring does:

A) Positive Work


B) Negative Work

C) Zero work

Clicker Question



A box attached at rest to a spring at its equilibrium length. You now push the box with your hand so that the spring is compressed a distance D , and you hold the box at rest in this new location.



During this motion, your hand does:

A) Positive Work


B) Negative Work

C) Zero work

Clicker Question




A box attached at rest to a spring at its equilibrium length. You now push the box with your hand so that the spring is compressed a distance D , and you hold the box at rest in this new location.



During this motion, the total work done on the box is:

- A) Positive B) Negative C) Zero



$$W = \int_{\vec{r}_1}^{\vec{r}_2} \vec{F}(\vec{r}) \cdot d\vec{r} = - \int_{r_1}^{r_2} \frac{GM_e m}{r^2} dr = \frac{GM_e m}{r} \Big|_{r_1}^{r_2} = GM_e m \left(\frac{1}{r_2} - \frac{1}{r_1} \right)$$

Clicker Question



In Case 1 we send an object from the surface of the earth to a height above the earth surface equal to one earth radius.

In Case 2 we start the same object a height of one earth radius above the surface of the earth and we send it infinitely far away.

In which case is the magnitude of the work done by the Earth's gravity on the object biggest?

A) Case 1 B) Case 2 C) They are the same

$$W = GM_em \left(\frac{1}{r_2} - \frac{1}{r_1} \right)$$



Case 1



Case 2




Clicker Question Solution



Case 1: $W = GM_em \left(\frac{1}{2R_E} - \frac{1}{R_E} \right) = -\frac{GM_em}{2R_E}$

Case 2: $W = GM_em \left(\frac{1}{\infty} - \frac{1}{2R_E} \right) = -\frac{GM_em}{2R_E}$

Same!



SCIENCE SPOOKTACULAR



featuring

WICKED SCIENCE ACTIVITIES!

**SPOOKTACULAR SHOWS OF
PHANTOM PHYSICS & CRYPTIC CHEMISTRY!**



Free Family Fun at Simon Fraser University

**Halloween Saturday
Saturday October 25, 2014**

Demos and activities 12:00-3:30
Academic Quadrangle, South Concourse 3000 level

Spooktacular shows 1:00 and 3:00
Shrum Science Centre, B9200 Lecture Theatre
Priority tickets available free (see website below)

Information: <http://www.sfu.ca/physics/outreach/science-spooktacular.html>

Volunteer: <http://www.sfu.ca/~exafs/spookvolunteer.html>



Costumes welcomed!