



# Electricity & Magnetism Lecture 17

## Today's Concept:



Faraday's Law

$$emf = \oint \vec{E} \cdot d\vec{\ell} = -\frac{d\Phi_B}{dt}$$



## Stuff you said

- There are so many types of RHR it is difficult to remember them all and which to apply
- what will happen if a magnet is moving towards a loop what if the magnet stays in a loop, I read from the book says the induced current will produce its own magnetic field thus increasing external flux don't understand
- Can you please explain why the magnet falling down in a metal tube is slower?
- I like the blue Ncdonalds fries with a halo on top. Grade A OC right there.

Faraday's Law: 
$$emf = \oint \vec{E} \cdot d\vec{\ell} = -\frac{d\Phi_B}{dt}$$
 where  $\Phi_B = \int \vec{B} \cdot d\vec{A}$ 

$$\Phi_B \equiv \int \vec{B} \cdot d\vec{A}$$

Looks scary but it's not — its amazing and beautiful!



A changing magnetic flux produces an electric field.




Electricity and magnetism are deeply connected.

Faraday's Law: 
$$emf = \oint \vec{E} \cdot d\vec{\ell} = -\frac{d\Phi_B}{dt}$$
 where  $\Phi_B \equiv \int \vec{B} \cdot d\vec{A}$ 

$$\Phi_B = \int \vec{B} \cdot d\vec{A}$$

#### In Practical Words:

- 1) When the flux  $\Phi_B$  through a loop changes, an *emf* is induced in the loop.
- 2) The *emf* will make a current flow if it can (like a battery).




Faraday's Law: 
$$emf = \oint \vec{E} \cdot d\vec{\ell} = -\frac{d\Phi_B}{dt}$$
 where  $\Phi_B = \int \vec{B} \cdot d\vec{A}$ 

$$\Phi_B = \int \vec{B} \cdot d\vec{A}$$

#### In Practical Words:

- 1) When the flux  $\Phi_B$  through a loop changes, an *emf* is induced in the loop.
- 2) The *emf* will make a current flow if it can (like a battery).
- 3) The current that flows induces a new magnetic field.




Faraday's Law: 
$$emf = \oint \vec{E} \cdot d\vec{\ell} = -\frac{d\Phi_B}{dt}$$

where

$$\Phi_B \equiv \int \vec{B} \cdot d\vec{A}$$

#### In Practical Words:

- 1) When the flux  $\Phi_R$  through a loop changes, an *emf* is induced in the loop.
- 2) The *emf* will make a current flow if it can (like a battery).
- 3) The current that flows induces a new magnetic field.
- 4) The new magnetic field opposes the change in the original magnetic field that created it. (Lenz' Law)



Faraday's Law: 
$$emf = \oint \vec{E} \cdot d\vec{\ell} = -\frac{d\Phi_B}{dt}$$


where 
$$\Phi_B \equiv \int \vec{B} \cdot d\vec{A}$$

## **Executive Summary:**



- $emf \rightarrow current \rightarrow field$  a) induced only when flux is changing
  - b) opposes the change


A copper loop is placed in a uniform magnetic field. You are looking from the right






Suppose the loop is moving to the right, The current induced in the loop is

- A) zero
- B) clockwise
- C) counterclockwise




A copper loop is placed in a uniform magnetic field. You are looking from the right




The induced B tries to boost up the decreasing external field.

Now suppose the loop is stationary and that the magnetic field is *decreasing*. The current induced in the loop is

- A) zero
- B) clockwise
- C) counterclockwise




Suppose the loop is spun around a vertical axis, and that it makes on complete revolution every second



The current induced in the loop


- A) is zero
- B) changes once per second
- C) changes twice per second








A horizontal copper ring is dropped from rest directly above the north pole of a permanent magnet.





Will the acceleration |a| of the falling ring in the presence of the magnet be any different than it would have been under the influence of just gravity?

- A) |a| > g
- B) |a| = g
- C) |a| < g

A rectangular loop (height = a, length = b, resistance = R, mass = m) coasts with a constant velocity  $v_0$  in +x direction as shown. At t = 0, the loop enters a region of constant magnetic field B directed in the -z direction.

x

What is the direction and the magnitude of the force on the loop when half of it is in the field?

#### **Conceptual Analysis**

Once loop enters B field region, flux will be changing in time Faraday's Law then says emf will be induced

#### Strategic Analysis


Find the emf

Find the current in the loop

Find the force on the current

A rectangular loop (height = a, length = b, resistance = R, mass = m) coasts with a constant velocity  $v_0$  in +x direction as shown. At t = 0, the loop enters a region of constant magnetic field B directed in the -z direction.

What is the magnitude of the emf induced in the loop just after it enters the field?




$$emf = -\frac{d\Phi_B}{dt}$$

A) 
$$\mathcal{E} = Babv_0^2$$
 B)  $\mathcal{E} = \frac{1}{2}Bav_0$  C)  $\mathcal{E} = \frac{1}{2}Bbv_0$  D)  $\mathcal{E} = Bav_0$  E)  $\mathcal{E} = Bbv_0$ 

A rectangular loop (height = a, length = b, resistance = R, mass = m) coasts with a constant velocity  $v_0$  in +x direction as shown. At t = 0, the loop enters a region of constant magnetic field B directed in the -z direction.

What is the direction of the current induced in the loop just after it enters the field?




$$emf = -\frac{d\Phi_B}{dt}$$

A rectangular loop (height = a, length = b, resistance = R, mass = m) coasts with a constant velocity  $v_0$  in +xdirection as shown. At t = 0, the loop enters a region of constant magnetic field B directed in the -z direction.

What is the direction of the net force on the loop just after it enters the field?


B) 
$$-y$$
 C)  $+x$ 



$$emf = -\frac{d\Phi_B}{dt}$$

A rectangular loop (height = a, length = b, resistance = R, mass = m) coasts with a constant velocity  $v_0$  in +x direction as shown. At t = 0, the loop enters a region of constant magnetic field B directed in the -z direction.

What is the magnitude of the net force on the loop just after it enters the field?



$$\vec{F} = I\vec{L} \times \vec{B}$$
  $\mathcal{E} = Bav_0$   $emf = -\frac{d\Phi_B}{dt}$ 

A) 
$$F = 4aBv_{o}R$$


B) 
$$F = a^2 B v_o R$$

A) 
$$F = 4aBv_o R$$
 B)  $F = a^2 Bv_o R$  C)  $F = a^2 B^2 v_o^2 / R$ 

$$D) F = a^2 B^2 v_o / R$$

## Follow Up

A rectangular loop (sides = a,b, resistance = R, mass = m) coasts with a constant velocity  $v_0$  in +x direction as shown. At t = 0, the loop enters a region of constant magnetic field B directed in the -z direction.



## What is the velocity of the loop when half of it is in the field?

t = dt:  $\mathcal{E} = Bav_0$ 

Which of these plots best represents the velocity as a function of time as the loop moves from entering the field to halfway through?

