

Mathematical Preliminaries

Ling324

Reading: *Meaning and Grammar*, pg. 529-540

Outline

- Set theory
- Ordered pairs and cartesian products
- Relations
- Functions

What is a Set?

- A SET is a collection of objects. It can be finite or infinite.

$$A = \{a, b, c\}$$

$$N = \{1, 2, 3, \dots\}$$

- An object is an ELEMENT of a set A if that object is a member of the collection A .

Notation: “ \in ” reads as “is an element of”, “is a member of”, or “belongs to”.

$$a \in A$$

$$2 \in N$$

A set can have another set as a member.

Let $B = \{a, b, c, \{d, e\}\}$, then $\{d, e\} \in B$

- A set with only one member is called a SINGLETON.
- A set with no members is called the EMPTY SET or NULL SET.

Notation: \emptyset or $\{ \}$

Specification of Sets

- List notation

A set consists of the objects named, not of the names themselves.

$$B = \{\text{The Amazon River, George Washington, 3}\}$$
$$C = \{\text{The Amazon River, 'George Washington', 3}\}$$

A set is unordered.

Writing the name of a member more than once does not change its membership status. For a given object, either it is a member of a given set or it is not.

$$\{a, b, c, e, e, e\}$$
$$\{a, b, c, e\}$$

Specification of Sets (cont.)

- Predicate notation

A better way to describe an infinite set is to indicate a property the members of the set share.

$\{x \mid x \text{ is an even number greater than } 3\}$ is read as
“the set of all x such that x is an even number greater than 3.”

‘ x ’ is a variable.

- Recursive rules

A rule for generating elements ‘recursively’ from a finite basis.

a) $4 \in E$

b) if $x \in E$, then $x + 2 \in E$

c) Nothing else belongs to E .

QUESTION: Give a list notation for the above recursive rules.

Set-theoretic Identity and Cardinality

- Two sets are IDENTICAL if and only if they have exactly the same members.

$$\{1, 2, 3, 4\} = \{x \mid x \text{ is a positive integer less than } 5\}$$

$$\begin{aligned} &\{x \mid x \text{ is a member of the Japanese male olympic gymnastics team in 2004}\} \\ &= \{x \mid x \text{ is a member of the team that won the gold medal in the 2004} \\ &\text{olympics for the male all-around gymnastics competition}\} \end{aligned}$$

- The number of members in a set A is called the CARDINALITY of A .

Notation: $|A|$

Let $A = \{1, 3, 5, a, b\}$.

$|A| =$

Subsets

- A set A is a SUBSET of a set B if all the elements of A are also in B .

Notation: $A \subseteq B$

- PROPER SUBSET

Notation: $A \subset B$

- $A \not\subseteq B$ means that A is not a subset of B .

- A set A is a subset of itself. $A \subseteq A$.

- If $A \subseteq B$, and $B \subseteq C$, then $A \subseteq C$.

Subsets (cont.)

QUESTION: Fill in the blank with either \subseteq or $\not\subseteq$.

- a) $\{a, b, c\}$ $\{s, b, a, e, g, i, c\}$
- b) $\{a, b, j\}$ $\{s, b, a, e, g, i, c\}$
- c) \emptyset $\{a\}$
- d) $\{a, \{a\}\}$ $\{a, b, \{a\}\}$
- e) $\{\{a\}\}$ $\{a\}$
- f) $\{a\}$ $\{\{a\}\}$
- g) $\{\emptyset\}$ $\{a\}$
- h) A A

QUESTION: Are the following statements true or false?

Let $A = \{b, \{c\}\}$.

- a) $c \in A$
- b) $\{c\} \in A$
- c) $\{b\} \subseteq A$
- d) $\{c\} \subseteq A$
- e) $\{\{c\}\} \subseteq A$
- f) $\{b\} \notin A$
- g) $\{b, \{c\}\} \subset A$
- h) $\{\{b, \{c\}\}\} \subseteq A$

Power Sets

- The POWER SET of A , $\wp(A)$, is the set whose members are all the subsets of A .

The set A itself and the null set are always members of $\wp(A)$.

Let $A = \{a, b\}$.

$$\wp(A) = \{\{a\}, \{b\}, \{a, b\}, \emptyset\}$$

- $|\wp(A)| = 2^n$
- From the definition of power set, it follows that $A \subseteq B$ iff $A \in \wp(B)$.

QUESTION: Let $B = \{a, b, c\}$. What is $\wp(B)$?

Set-theoretic Operations

- $A \cap B$: The INTERSECTION of two sets A and B is the set containing all and only the objects that are elements of both A and B .

$$A \cap B \cap C = \cap\{A, B, C\}$$

- $A \cup B$: The UNION of two sets A and B is the set containing all and only the objects that are elements of A , of B , or of both A and B .

$$A \cup B \cup C = \cup\{A, B, C\}$$

- $A - B$: The DIFFERENCE between two sets A and B subtracts from A all objects which are in B .

- A' : The COMPLEMENT of a set A is the set of all the individuals in the universe of discourse except for the elements of A (i.e., $U - A$).

Set-theoretic Operations (cont.)

QUESTION: Let $K = \{a, b\}$, $L = \{c, d\}$, and $M = \{b, d\}$.

a) $K \cup L = \{a, b, c, d\}$

b) $K \cup M =$

c) $(K \cup L) \cup M =$

d) $L \cup \emptyset =$

e) $K \cap L = \emptyset$

f) $L \cap M =$

g) $K \cap K =$

h) $K \cap \emptyset =$

i) $K \cap (L \cap M) =$

j) $K \cap (L \cup M) =$

k) $K - M = \{a\}$

m) $L - M =$

n) $M - L =$

o) $K - \emptyset =$

p) $\emptyset - K =$

Set-theoretic Equalities

- Some fundamental set-theoretic equalities

- Commutative Laws

$$X \cup Y = Y \cup X$$

$$X \cap Y = Y \cap X$$

- Associative Laws

$$(X \cup Y) \cup Z = X \cup (Y \cup Z)$$

$$(X \cap Y) \cap Z = X \cap (Y \cap Z)$$

- Distributive Laws

$$X \cup (Y \cap Z) = (X \cup Y) \cap (X \cup Z)$$

$$X \cap (Y \cup Z) = (X \cap Y) \cup (X \cap Z)$$

- Identity Laws

$$X \cup \emptyset = X$$

$$X \cup U = U$$

$$X \cap \emptyset = \emptyset$$

$$X \cap U = X$$

- Complement Laws

$$X \cup X' = U$$

$$(X')' = X$$

$$X \cap X' = \emptyset$$

$$X - Y = X \cap Y'$$

- DeMorgan's Laws

$$(X \cup Y)' = X' \cap Y'$$

$$(X \cap Y)' = X' \cup Y'$$

Set-theoretic Equalities (cont.)

- Set-theoretic equalities can be used to simplify a complex set-theoretic expression, or to prove the truth of other statements about sets.

Simplify the expression $(A \cup B) \cup (B \cap C)'$.

- | | | |
|----|--------------------------------|-------------|
| 1. | $(A \cup B) \cup (B \cap C)'$ | |
| 2. | $(A \cup B) \cup (B' \cup C')$ | DeMorgan |
| 3. | $A \cup (B \cup (B' \cup C'))$ | Associative |
| 4. | $A \cup ((B \cup B') \cup C')$ | Associative |
| 5. | $A \cup (U \cup C')$ | Complement |
| 6. | $A \cup (C' \cup U)$ | Commutative |
| 7. | $A \cup U$ | Identity |
| 8. | U | Identity |

QUESTION: Show that $(A \cap B) \cap (A \cap C)' = A \cap (B - C)$.

Ordered Pairs and Cartesian Products

- A SEQUENCE of objects is a list of these objects in some order. (cf., Recall that a set is unordered.)

$\langle a, b, c \rangle, \langle 7, 21, 57 \rangle, \langle 1, 2, 3, \dots \rangle$

- Finite sequences are called TUPLES. A sequence with k elements is a K-TUPLE.

A 2-tuple is also called an (ordered) PAIR.

$\langle a, b \rangle$

- If A and B are two sets, the CARTESIAN PRODUCT of A and B , written as $A \times B$, is the set containing all pairs wherein the first element is a member of A and the second element is a member of B .
- Although each member of a Cartesian product is an ordered pair, the Cartesian product itself is an unordered set of them.

Ordered Pairs and Cartesian Products (cont.)

QUESTION: Let $K = \{a, b, c\}$ and $L = \{1, 2\}$.

$$K \times L = \{ \langle a, 1 \rangle, \langle a, 2 \rangle, \langle b, 1 \rangle, \langle b, 2 \rangle, \langle c, 1 \rangle, \langle c, 2 \rangle \}$$

$$L \times K =$$

$$L \times L =$$

QUESTION: Let $A \times B = \{ \langle a, 1 \rangle, \langle a, 2 \rangle, \langle a, 3 \rangle \}$. What is $\wp(A \times B)$?

Relations

- A RELATION is a set of ordered n-tuples.

Binary relation: e.g., mother of, kiss, subset

Ternary relation: e.g., give

Unary relation: a set of individuals. e.g., being a ling324 student, being a Canadian

- A relation from A to B is a subset of the Cartesian product $A \times B$.

Rab , or aRb : The relation R holds between objects a and b .

$R \subseteq A \times B$: A relation between objects from two sets A and B . A relation *from* A *to* B .

$R \subseteq A \times A$: A relation holding of objects from a single set A is called a relation *in* A .

Relations (cont.)

- $\text{Domain}(R) = \{a \mid \text{there is some } b \text{ such that } \langle a, b \rangle \in R\}$
 $\text{Range}(R) = \{b \mid \text{there is some } a \text{ such that } \langle a, b \rangle \in R\}$

Let $A = \{a, b\}$ and $B = \{c, d, e\}$. $R = \{\langle a, d \rangle, \langle a, e \rangle, \langle b, e \rangle\}$.
 $\text{Domain}(R) = \{a, b\}$; $\text{Range}(R) = \{d, e\}$

Note: A relation may relate one object in its domain to more than one object in its range.

- The COMPLEMENT of a relation $R \subseteq A \times B$, written R' , contains all ordered pairs of the Cartesian product which are not members of the relation R .

The INVERSE of a relation, written as R^{-1} , has as its members all the ordered pairs in R , with their first and second elements reversed.

$$(R')' = R; (R^{-1})^{-1} = R$$

If $R \subseteq A \times B$, then $R^{-1} \subseteq B \times A$, but $R' \subseteq A \times B$.

QUESTION: Let $A = \{1, 2, 3\}$ and $R \subseteq A \times A$ be
 $\{\langle 3, 2 \rangle, \langle 3, 1 \rangle, \langle 2, 1 \rangle\}$, which is 'greater than' relation in A .
What is R' ? What is R^{-1} ?

Types of Relations

- Reflexive: for all a in the domain, $\langle a, a \rangle \in R$. e.g., being the same age as
Nonreflexive: e.g., like
Irreflexive: for all a in the domain, $\langle a, a \rangle \notin R$. e.g., proper subset
- Symmetric: whenever $\langle a, b \rangle \in R$, $\langle b, a \rangle \in R$. e.g., being five miles from
Nonsymmetric: e.g., being the sister of
Asymmetric: it is never the case that $\langle a, b \rangle \in R$ and $\langle b, a \rangle \in R$. e.g., being the mother of
- Transitive: whenever $\langle a, b \rangle \in R$ and $\langle b, c \rangle \in R$, $\langle a, c \rangle \in R$. e.g., being older than
Nontransitive: e.g., like
Intransitive: whenever $\langle a, b \rangle \in R$ and $\langle b, c \rangle \in R$, it is not the case that $\langle a, c \rangle \in R$. e.g., being the mother of
- Equivalence: reflexive, transitive and symmetric. e.g., being the same age as
An equivalence relation PARTITIONS a set A into EQUIVALENCE CLASSES, which are DISJOINT and whose union is identical with A .

Functions

- A relation R from A to B is a FUNCTION from A to B if and only if:
 - a) Each element in the domain is paired with just one element in the range.
 - b) The domain of R is equal to A (except for PARTIAL FUNCTIONS).

QUESTION: Let $A = \{a, b, c\}$ and $B = \{1, 2, 3, 4\}$. Which of the following relations are functions?

- a) $P = \{ \langle a, 1 \rangle, \langle b, 2 \rangle, \langle c, 3 \rangle \}$
 - b) $Q = \{ \langle a, 1 \rangle, \langle b, 2 \rangle \}$
 - c) $R = \{ \langle a, 2 \rangle, \langle a, 3 \rangle, \langle b, 4 \rangle \}$
 - d) $S = \{ \langle a, 3 \rangle, \langle b, 2 \rangle, \langle c, 2 \rangle \}$
- A function that is a subset of $A \times B$ is a function *from* A to B .
A function in $A \times A$ is a function *in* A .
 - $F : A \rightarrow B$ is read as “ F is function from A to B ”.
 - $F(a) = b$ is read as “ F maps a to b ”.
 - Given $F(a) = b$, a is an ARGUMENT and b is the VALUE.

Functions (cont.)

- Functions from A to B are generally said to be INTO B .
Functions from A to B such that the range of the function equals B are called ONTO B .
- A function $F : A \rightarrow B$ is called a ONE-TO-ONE function just in case no member of B is assigned to more than one member of A . Otherwise, we will call them MANY-TO-ONE function.
- A function which is both one-to-one and onto is called a ONE-TO-ONE CORRESPONDENCE.
If a function F is a one-to-one correspondence, F^{-1} is also a function.
- A function with k arguments is called a K-ARY FUNCTION, and k is called the ARITY of the function.

Unary function takes one argument. $F(a)$.

Binary function takes two arguments. $F(a, b)$.

- Infix notation: e.g., $a + b$.
Prefix notation: e.g., $+(a, b)$.

Functions (cont.)

- A PREDICATE or PROPERTY is a function whose range is $\{\text{True}, \text{False}\}$.

$\text{even_number}(2) = \text{True}$, $\text{even_number}(3) = \text{False}$.

take_ling324 , male , freshman .

- The specification of a function from a domain D to $\{\text{True}, \text{False}\}$ defines a unique subset of domain D , and the specification of a set defines a unique function.

For instance, let the domain D be a set of numbers. If we collect all the elements in D that are mapped to True by function *even_number*, we end up with a set of even numbers, which is a subset of D .

Also, if a number is an element of the set of even numbers, then it is mapped to True by function *even_number*, and if it is not an element of the set of even numbers, then it is mapped to False by function *even_number*.

- We call the unique function that is associated with set A , the CHARACTERISTIC FUNCTION of A .