

SIMON FRASER UNIVERSITY
Department of Economics

Econ 835
Quantitative Methods

Prof. Kasa
Fall 2001

MIDTERM EXAM

Answer the following questions True, False, or Uncertain. Briefly explain your answers. No credit without explanation. (5 points each).

1. Adding too many variables to a regression model leads to biased estimates.
2. The sum of the Least Squares residuals is always zero.
3. When estimating the model, $y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \beta_3 x_3 + u$, we know that if the individual t -statistics for β_2 and β_3 are insignificant then the F -statistic for the joint hypothesis $\beta_2 = \beta_3 = 0$ must also be insignificant.
4. The ‘usual’ t -statistics only have a t -distribution if the model’s error term is normally distributed.
5. If the explanatory variables in a multiple regression are not correlated with each other, the multiple regression slope coefficients are the same as the slope coefficient estimates we would get by running a set of individual bivariate regressions.
6. Although multicollinearity does not cause biased coefficient estimates, it does cause biased estimates of their variances.
7. Heteroskedasticity causes the Ordinary Least Squares estimator to be biased.

The following questions are short answer. Briefly explain your answer. Clarity will be rewarded.

8. (15 points). Suppose you want to study the effectiveness of a certain drug in reducing cholesterol. You have a sample of data on individuals measuring the change in their cholesterol levels over a given time period. For each individual you also have data on whether they took the drug or not and on whether they are male or female.
 - (a) Describe how you would formulate a model to test whether the drug is effective. Write down the exact model you would estimate. State explicitly what your null and alternative hypotheses would be, and what test statistic you would compute.
 - (b) Now suppose you want to test the hypothesis that the drug is more effective for women than for men. What model would you estimate? What test statistic would you compute? (Note: Since the dependent variable is the *change* in cholesterol, you don’t need to worry about the possibility that average cholesterol levels differ between men and women).

9. (15 points). State the Gauss-Markov Theorem. On what assumptions is it based? Why is it important?
10. (10 points). In a study with a large number of observations, the income elasticity of demand is estimated to be 0.70 with a standard error of 0.12. Is the elasticity significantly less than 1.0? Assume that you are only willing to tolerate a 5% chance of incorrectly rejecting the null hypothesis.
11. (10 points). Briefly describe White's test for heteroskedasticity. How do you compute it? What is its (asymptotic) distribution? Explain the intuition behind it.
12. (10 points). Your friend has correctly pointed out that correlation between a model's error term and the explanatory variables causes Least Squares to produce biased estimates. She then proposes the following strategy to test for the presence of such correlation. First, estimate the original equation, and then regress the residuals on the X variables in order to see if there is a significant relationship. Briefly comment on this strategy. Do you see any potential problems with this approach?
13. (5 points). Suppose you are interested in studying the influence of macroeconomic conditions on industry profitability. You have a cross-sectional sample of 16 industries, containing data on each industry's profits during a given year and the growth rate of GDP during that year. One idea is to simply regress profits on GDP growth. Do you see any problems with this approach?