STAT 380
Poisson Processes

Richard Lockhart

Simon Fraser University

Spring 2021

Poisson Processes

- Particles arriving over time at a particle detector.
- Several ways to describe most common model.
- Approach 1:
  - numbers of particles arriving in an interval has Poisson distribution,
  - mean proportional to length of interval,
  - numbers in several non-overlapping intervals independent.

Notation, formal assumptions

- For $s < t$, denote number of arrivals in $(s, t]$ by $N(s, t)$. Model is
  1. $N(s, t)$ has a Poisson$(\lambda(t - s))$ distribution.
  2. For $0 \le s_1 < t_1 \le s_2 < t_2 \cdots \le s_k < t_k$ the variables $N(s_i, t_i); i = 1, \ldots, k$ are independent.
  3. Jargon: *independent Poisson increments*.

Approach 2

- Let $0 < S_1 < S_2 < \cdots$ be the times at which the particles arrive.
- Let $T_i = S_i - S_{i-1}$ with $S_0 = 0$ by convention.
- Then $T_1, T_2, \ldots$ are independent Exponential random variables with mean $1/\lambda$.
- Note $P(T_i > x) = e^{-\lambda x}$ is called **survival function** of T_i.
- Approaches 1 and 2 are equivalent.
- Both are deductions of a model based on **local** behaviour of process.

Approach 3

Assume:

1. given all the points in $[0, t]$ the probability of 1 point in the interval $(t, t+h]$ is of the form
$$\lambda h + o(h)$$
2. given all the points in $[0, t]$ the probability of 2 or more points in interval $(t, t+h]$ is of the form
$$o(h)$$

- All 3 approaches are equivalent.
- I show: 3 implies 1, 1 implies 2 and 2 implies 3.
- First explain o, O.

Landau's big and little 'O'

- **Notation**: given functions f and g we write

$$f(h) = g(h) + o(h)$$

provided

$$\lim_{h \to 0} \frac{f(h) - g(h)}{h} = 0$$

- Aside: if there is a constant M such that

$$\limsup_{h \to 0} \left| \frac{f(h) - g(h)}{h} \right| \leq M$$

we say

$$f(h) = g(h) + O(h)$$

- Another form is $f(h) = g(h) + O(h)$ means there is $\delta > 0$ and M s.t. for all $|h| < \delta$

$$|f(h) - g(h)| \leq Mh$$

- Idea: $o(h)$ is tiny compared to h while $O(h)$ is (very) roughly the same size as h.

Model 3 implies 1

- Fix t, define $f_t(s)$ to be conditional probability of 0 points in $(t, t+s]$ given value of process on $[0, t]$.
- Derive differential equation for f.
- Given process on $[0, t]$ and 0 points in $(t, t+s]$ probability of no points in $(t, t+s+h]$ is

$$f_{t+s}(h) = 1 - \lambda h + o(h)$$

- Given the process on $[0, t]$ the probability of no points in $(t, t+s]$ is $f_t(s)$.
- Using $P(AB|C) = P(A|BC)P(B|C)$ gives

$$\begin{aligned} f_t(s+h) &= f_t(s) f_{t+s}(h) \\ &= f_t(s)(1 - \lambda h + o(h)) \end{aligned}$$

3 implies 1 continued

- Now rearrange, divide by h to get

$$\frac{f_t(s+h) - f_t(s)}{h} = -\lambda f_t(s) + \frac{o(h)}{h}$$

- Let $h \to 0$ and find

$$\frac{\partial f_t(s)}{\partial s} = -\lambda f_t(s)$$

- Differential equation has solution

$$f_t(s) = f_t(0)\exp(-\lambda s) = \exp(-\lambda s)\,.$$

- Notice: survival function of exponential rv.

General case

- Notation: $N(t) = N(0, t)$.
- $N(t)$ is a non-decreasing function of t.
- Let
$$P_k(t) = P(N(t) = k)$$
- Evaluate $P_k(t + h)$ by conditioning on $N(s); 0 \leq s < t$ and $N(t) = j$.
- Given $N(t) = j$ probability that $N(t + h) = k$ is conditional probability of $k - j$ points in $(t, t + h]$.
- So, for $j \leq k - 2$:
$$P(N(t + h) = k | N(t) = j, N(s), 0 \leq s < t) = o(h).$$

General case, continued

- For $j = k - 1$ we have

$$P(N(t+h) = k | N(t) = k-1, N(s), 0 \le s < t) = \lambda h + o(h)$$

- For $j = k$ we have

$$P(N(t+h) = k | N(t) = k, N(s), 0 \le s < t) = 1 - \lambda h + o(h)$$

- N is increasing so only consider $j \le k$.

$$\begin{aligned} P_k(t+h) &= \sum_{j=0}^{k} P(N(t+h) = k | N(t) = j) P_j(t) \\ &= P_k(t)(1 - \lambda h) + \lambda h P_{k-1}(t) + o(h) \end{aligned}$$

- Rearrange, divide by h and let $h \to 0$ t get

$$P_k'(t) = -\lambda P_k(t) + \lambda P_{k-1}(t)$$

General case, continued III

- For $k = 0$ the term P_{k-1} is dropped and

$$P_0'(t) = -\lambda P_0(t)$$

- Using $P_0(0) = 1$ we get

$$P_0(t) = e^{-\lambda t}$$

- Put this into the equation for $k = 1$ to get

$$P_1'(t) = -\lambda P_1(t) + \lambda e^{-\lambda t}$$

- Multiply by $e^{\lambda t}$ to see

$$\left(e^{\lambda t} P_1(t)\right)' = \lambda$$

- With $P_1(0) = 0$ we get

$$P_1(t) = \lambda t e^{-\lambda t}$$

General case, continued IV

- For general k we have $P_k(0) = 0$ and

$$\left(e^{\lambda t} P_k(t)\right)' = \lambda e^{\lambda t} P_{k-1}(t)$$

- Check by induction that

$$e^{\lambda t} P_k(t) = (\lambda t)^k / k!$$

- Hence: $N(t)$ has Poisson(λt) distribution.
- Similar ideas permit proof of

$$P\left(N(s,t) = k | N(u); 0 \le u \le s\right) = \frac{\{\lambda(t-s)\}^k e^{-\lambda(t-s)}}{k!}$$

- By induction can prove N has independent Poisson increments.

Exponential Interarrival Times

- Suppose N is a Poisson Process.
- Define $T_1, T_2, \ldots$ to be the times between 0 and the first point, the first point and the second and so on.
- Fact: $T_1, T_2, \ldots$ are iid exponential rvs with mean $1/\lambda$.
- Already did T_1 rigorously.
- The event $T_1 > t$ is exactly the event $N(t) = 0$.
- So

$$P(T_1 > t) = P(N(t) = 0) = \exp(-\lambda t)$$

which is the survival function of an exponential rv.

Exponential Interarrival Times deduced

- I do case of T_1, T_2.
- Let t_1, t_2 be two positive numbers and $s_1 = t_1$, $s_2 = t_1 + t_2$.
- Consider event

$$\{t_1 < T_1 \le t_1 + \delta_1\} \cap \{t_2 < T_2 \le t_2 + \delta_2\}.$$

- This is almost the same as the intersection of the four events:

$$\begin{aligned} N(0, t_1] &= 0 \\ N(t_1, t_1 + \delta_1] &= 1 \\ N(t_1 + \delta_1, t_1 + \delta_1 + t_2] &= 0 \\ N(s_2 + \delta_1, s_2 + \delta_1 + \delta_2] &= 1 \end{aligned}$$

which has probability

$$e^{-\lambda t_1} \times \lambda \delta_1 e^{-\lambda \delta_1} \times e^{-\lambda t_2} \times \lambda \delta_2 e^{-\lambda \delta_2}$$

Less Rigor

- Divide by $\delta_1\delta_2$.
- Let δ_1 and δ_2 go to 0 to get joint density of T_1, T_2:

$$\lambda^2 e^{-\lambda t_1} e^{-\lambda t_2}$$

- This is joint density of two independent exponential variates.

More Rigor

- Find joint density of $S_1, \ldots, S_k$.
- Use **change of variables** to find joint density of $T_1, \ldots, T_k$.

More Rigor, II

- First step: Compute

$$P(0 < S_1 \le s_1 < S_2 \le s_2 \cdots < S_k \le s_k)$$

- This is just the event of exactly 1 point in each interval $(s_{i-1}, s_i]$ for $i = 1, \ldots, k-1$ ($s_0 = 0$) and at least one point in $(s_{k-1}, s_k]$.
- This event has probability

$$\prod_1^{k-1} \left\{ \lambda(s_i - s_{i-1}) e^{-\lambda(s_i - s_{i-1})} \right\} \left(1 - e^{-\lambda(s_k - s_{k-1})} \right)$$

- Second step: write this in terms of joint cdf of $S_1, \ldots, S_k$.

More Rigor, III, $k = 2$

$$P(0 < S_1 \leq s_1 < S_2 \leq s_2) = F_{S_1,S_2}(s_1, s_2) - F_{S_1,S_2}(s_1, s_1)$$

- Notice tacit assumption $s_1 < s_2$.
- Differentiate twice, that is, take

$$\frac{\partial^2}{\partial s_1 \partial s_2}$$

to get

$$f_{S_1,S_2}(s_1, s_2) = \frac{\partial^2}{\partial s_1 \partial s_2} \lambda s_1 e^{-\lambda s_1} \left(1 - e^{-\lambda(s_2 - s_1)}\right)$$

- Simplify to

$$\lambda^2 e^{-\lambda s_2}$$

- Recall tacit assumption to get

$$f_{S_1,S_2}(s_1, s_2) = \lambda^2 e^{-\lambda s_2} 1(0 < s_1 < s_2)$$

More Rigor, IV, change of variables

- Now compute the joint cdf of T_1, T_2 by

$$F_{T_1,T_2}(t_1,t_2) = P(S_1 < t_1, S_2 - S_1 < t_2)$$

- This is

$$\begin{aligned} P(S_1 < t_1, S_2 - S_1 < t_2) &= \int_0^{t_1} \int_{s_1}^{s_1+t_2} \lambda^2 e^{-\lambda s_2} ds_2 ds_1 \\ &= \lambda \int_0^{t_1} \left(e^{-\lambda s_1} - e^{-\lambda(s_1+t_2)} \right) ds_1 \\ &= 1 - e^{-\lambda t_1} - e^{-\lambda t_2} + e^{-\lambda(t_1+t_2)} \end{aligned}$$

- Differentiate twice to get

$$f_{T_1,T_2}(t_1,t_2) = \lambda e^{-\lambda t_1} \lambda e^{-\lambda t_2}$$

- This is joint density of 2 independent exponential rvs.

Summary so far

Have shown:

- Instantaneous rates model implies independent Poisson increments model
- Independent Poisson increments model implies independent exponential interarrivals.
- Next: show independent exponential interarrivals implies the instantaneous rates model.

Exponential interarrivals implies rates

- Suppose $T_1, \ldots$ iid exponential rvs with means $1/\lambda$.
- Define N_t by $N_t = k$ if and only if

$$T_1 + \cdots + T_k \le t \le T_1 + \cdots + T_{k+1}$$

- Let A be the event $N(s) = n(s); 0 < s \le t$.
- We are to show

$$P(N(t, t+h] = 1 | N(t) = k, A) = \lambda h + o(h)$$

and

$$P(N(t, t+h] \ge 2 | N(t) = k, A) = o(h)$$

- If $n(s)$ is a possible trajectory consistent with $N(t) = k$ then n has jumps at points

$$s_1 \equiv t_1, s_2 \equiv t_1 + t_2, \ldots, s_k \equiv t_1 + \cdots + t_k < t$$

and at no other points in $(0, t]$.

Continued

- So given $N(s) = n(s); 0 < s \leq t$ with $n(t) = k$ we are essentially being given
$$T_1 = t_1, \ldots, T_k = t_k, T_{k+1} > t - s_k$$
and asked the conditional probability in the first case of the event B given by
$$B = \{t - s_k < T_{k+1} \leq t - s_k + h < T_{k+2} + T_{k+1}\} .$$
- Conditioning on $T_1, \ldots, T_k$ irrelevant (independence).
$$\begin{aligned} P(N(t, t+h] = 1 | N(t) = k, A)/h \\ &= P(B | T_{k+1} > t - s_k)/h \\ &= \frac{P(B)}{he^{-\lambda(t-s_k)}} \end{aligned}$$

Continued

- The numerator may be evaluated by integration:

$$P(B) = \int_{t-s_k}^{t-s_k+h} \int_{t-s_k+h-s_1}^{\infty} \lambda^2 e^{-\lambda(s_1+s_2)} ds_2 ds_1$$

- Let $h \to 0$ to get the limit

$$P(N(t, t+h] = 1 | N(t) = k, A)/h \to \lambda$$

as required.

- The computation of

$$\lim_{h \to 0} P(N(t, t+h] \geq 2 | N(t) = k, A)/h$$

is similar.

Properties of exponential rvs

- **Convolution**: If X and Y independent rvs with densities f and g respectively and $Z = X + Y$ then

$$P(Z \le z) = \int_{-\infty}^{\infty} \int_{-\infty}^{z-x} f(x)g(y)dydx$$

- Differentiating wrt z we get

$$f_Z(z) = \int_{-\infty}^{\infty} f(x)g(z-x)dx$$

- This integral is called the **convolution** of densities f and g.
- If $T_1, \ldots, T_n$ iid Exponential(λ) then $S_n = T_1 + \cdots + T_n$ has a Gamma(n, λ) distribution.
- Density of S_n is

$$f_{S_n}(s) = \lambda(\lambda s)^{n-1} e^{-\lambda s}/(n-1)!$$

for $s > 0$.

Proof

$$P(S_n > s) = P(N(0,s] < n) = \sum_{j=0}^{n-1} (\lambda s)^j e^{-\lambda s}/j!$$

- Then

$$\begin{aligned} f_{S_n}(s) &= \frac{d}{ds} P(S_n \le s) \\ &= \frac{d}{ds} \left\{1 - P(S_n > s)\right\} \\ &= -\lambda \sum_{j=1} n-1 \left\{ j(\lambda s)^{j-1} - (\lambda s)^j \right\} \frac{e^{-\lambda s}}{j!} + \lambda e^{-\lambda s} \\ &= \lambda e^{-\lambda s} \sum_{j=1}^{n-1} \left\{ \frac{(\lambda s)^j}{j!} - \frac{(\lambda s)^{j-1}}{(j-1)!} \right\} + \lambda e^{-\lambda s} \end{aligned}$$

- This telescopes to

$$f_{S_n}(s) = \lambda (\lambda s)^{n-1} e^{-\lambda s}/(n-1)!$$

Extreme Values

- Suppose $X_1, \ldots, X_n$ are independent exponential rvs with means $1/\lambda_1, \ldots, 1/\lambda_n$
- Then $Y = \min\{X_1, \ldots, X_n\}$ has exponential distribution with mean

$$\frac{1}{\lambda_1 + \cdots + \lambda_n}$$

- Proof:

$$\begin{aligned} P(Y > y) &= P(\forall k X_k > y) \\ &= \prod e^{-\lambda_k y} \\ &= e^{-\sum \lambda_k y} \end{aligned}$$

Memoryless Property

- Suppose X has exponential distribution.
- Conditional distribution of $X - x$ given $X \geq x$ is also exponential.
- Proof:

$$P(X - x > y | X \geq x) = \frac{P(X > x + y, X \geq x)}{P(X > x)}$$

$$= \frac{PX > x + y)}{P(X \geq x)}$$

$$= \frac{e^{-\lambda(x+y)}}{e^{-\lambda x}}$$

$$= e^{-\lambda y}$$

Hazard Rates

- Assume rv $T > 0$ with density f and cdf F
- Hazard rate, or instantaneous failure rate, is

$$r(t) = \lim_{\delta \to 0} \frac{P(t < T \leq t + \delta | T \geq t)}{\delta}$$

- This is just

$$r(t) = \frac{f(t)}{1 - F(t)}$$

- For an exponential random variable with mean $1/\lambda$ this is

$$h(t) = \frac{\lambda e^{-\lambda t}}{e^{-\lambda t}} = \lambda$$

- The exponential distribution has constant failure rate.

The Weibull distribution

- **Weibull** random variables have density, for $t > 0$,

$$f(t|\lambda, \alpha) = \lambda(\lambda t)^{\alpha-1} e^{-(\lambda t)^{\alpha}}.$$

- The corresponding survival function is, for $t > 0$,

$$1 - F(t) = e^{-(\lambda t)^{\alpha}}$$

- The hazard rate is

$$r(t) = \lambda(\lambda t)^{\alpha-1}$$

- Hazard rate is increasing for $\alpha > 1$, decreasing for $\alpha < 1$.
- For $\alpha = 1$ this is the exponential distribution.

Weibull distribution, continued

- Since
$$r(t) = \frac{dF(t)/dt}{1 - F(t)} = -\frac{d\log(1 - F(t))}{dt}$$
we can integrate to find
$$1 - F(t) = \exp\{-\int_0^t r(s)ds\}.$$
- So r determines F and f.

Properties of Poisson Processes

1) If N_1 and N_2 are independent Poisson processes with rates λ_1 and λ_2, respectively, then $N = N_1 + N_2$ is a Poisson process with rate $\lambda_1 + \lambda_2$.
2) Suppose N is a Poisson process with rate λ. Suppose each point is marked with a label, say one of $L_1, \ldots, L_r$, independently of all other occurrences.

- Suppose p_i is the probability that a given point receives label L_i.
- Let N_i count the points with label i (so that $N = N_1 + \cdots + N_r$).
- Then $N_1, \ldots, N_r$ are independent Poisson processes with rates $p_i\lambda$.

Properties of Poisson Processes

3) Suppose $U_1, U_2, \ldots$ independent rvs, each uniformly distributed on $[0, T]$.

- Suppose M is a Poisson(λT) random variable independent of the U's.
- Let

$$N(t) = \sum_1^M 1(U_i \le t)$$

- Then N is a Poisson process on $[0, T]$ with rate λ.

Properties of Poisson Processes

4) Suppose N is a Poisson process with rate λ.

- Let $S_1 < S_2 < \cdots$ be the times at which points arrive
- Given $N(T) = n$ $S_1, \ldots, S_n$ have the same distribution as the order statistics of a sample of size n from the uniform distribution on $[0, T]$.

5) Given $S_{n+1} = T$, $S_1, \ldots, S_n$ have the same distribution as the order statistics of a sample of size n from the uniform distribution on $[0, T]$.

Indications of some proofs

1) $N_1, \ldots, N_r$ independent Poisson processes rates λ_i, $N = \sum N_i$.

- Let A_h be the event of 2 or more points in N in the time interval $(t, t+h]$
- Let B_h be the event of exactly one point in N in the time interval $(t, t+h]$.
- Let A_{ih} and B_{ih} be the corresponding events for N_i.
- Let H_t denote the history of the processes up to time t; we condition on H_t.
- We are given:

$$P(A_{ih}|H_t) = o(h)$$

and

$$P(B_{ih}|H_t) = \lambda_i h + o(h)\,.$$

Indications of some proofs, II

- Note that

$$A_h \subset \bigcup_{i=1}^{r} A_{ih} \cup \bigcup_{i \neq j} (B_{ih} \cap B_{jh})$$

- Since

$$\begin{aligned} P(B_{ih} \cap B_{jh} | H_t) &= P(B_{ih} | H_t) P(B_{jh} | H_t) \\ &= (\lambda_i h + o(h))(\lambda_j h + o(h)) \\ &= O(h^2) \\ &= o(h) \end{aligned}$$

we have checked one of the two infinitesimal conditions for a Poisson process.

Indications of some proofs, III

- Next let C_h be the event of no points in N in the time interval $(t, t+h]$.
- Let C_{ih} be the same for N_i.
- Then

$$\begin{aligned} P(C_h|H_t) &= P(\cap C_{ih}|H_t) \\ &= \prod P(C_{ih}|H_t) \\ &= \prod (1 - \lambda_i h + o(h)) \\ &= 1 - (\sum \lambda_i)h + o(h) \end{aligned}$$

shows

$$\begin{aligned} P(B_h|H_t) &= 1 - P(C_h|H_t) - P(A_h|H_t) \\ &= (\sum \lambda_i)h + o(h) \end{aligned}$$

- Hence N is a Poisson process with rate $\sum \lambda_i$.

Indications of some proofs, IV

2) The infinitesimal approach used for 1 can do part of this.

- See text for rest.
- Events defined as in **1)**:
- Define event B_{ih}: there is one point in N_i in $(t, t+h]$
- Define event B_h:there is exactly one point in any of the r processes,
- B_{ih} is union of B_h and the subset of A_h where there are two or more points in N in $(t, t+h]$ but exactly one is labelled i.
- Since $P(A_h|H_t) = o(h)$

$$\begin{aligned} P(B_{ih}|H_t) &= p_i P(B_h|H_t) + o(h) \\ &= p_i(\lambda h + o(h)) + o(h) \\ &= p_i \lambda h + o(h) \end{aligned}$$

Indications of some proofs, V

- Similarly, A_{ih} is a subset of A_h so

$$P(A_{ih}|H_t) = o(h)$$

- This shows each N_i is Poisson with rate λp_i.
- To get independence requires more work; see the text for the algebraic method which is easier.

Indications of some proofs, VI

3) Fix $s < t$.

- Let $N(s,t)$ be the number of points in $(s,t]$.
- Given $N = n$ the conditional distribution of $N(s,t)$ is Binomial(n,p) with $p = (t-s)/T$.

Indications of some proofs, VII

- So
$$
\begin{aligned}
P(N(s,t)=k) &= \sum_{n=k}^{\infty} P(N(s,t)=k, N=n) \\
&= \sum_{n=k}^{\infty} P(N(s,t)=k|N=n)P(N=n) \\
&= \sum_{n=k}^{\infty} \frac{n!}{k!(n-k)!} p^k (1-p)^{n-k} \frac{(\lambda T)^n}{n!} e^{-\lambda T} \\
&= \frac{e^{-\lambda T}}{k!} (\lambda T p)^k \sum_{n=k}^{\infty} \frac{(1-p)^{n-k}(\lambda T)^{n-k}}{(n-k)!} \\
&= \frac{e^{-\lambda T}}{k!} (\lambda T p)^k \sum_{m=0}^{\infty} (1-p)^m (\lambda T)^m / m! \\
&= \frac{e^{-\lambda T}}{k!} (\lambda T p)^k e^{\lambda T(1-p)} \\
&= \frac{e^{-\lambda(t-s)}(\lambda(t-s))^k}{k!}
\end{aligned}
$$

Indications of some proofs, VIII

4) Fix s_i, h_i for $i = 1, \ldots, n$ such that

$$0 < s_1 < s_1 + h_1 < s_2 < \cdots < s_n < s_n + h_n < T$$

- Given $N(T) = n$ we compute the probability of the event

$$A = \bigcap_{i=1}^{n} \{s_i < S_i < s_i + h_i\}$$

- Intersection of A, $N(T) = n$ is ($s_0 = h_0 = 0$):

$$B \equiv \bigcap_{i=1}^{n} \{N(s_{i-1} + h_{i-1}, s_i] = 0, N(s_i, s_i + h_i] = 1\}$$

$$\cap \{N(s_n + h_n, T] = 0\}$$

whose probability is

$$\left(\prod \lambda h_i\right) e^{-\lambda T}$$

Indications of some proofs, IX

- So

$$
\begin{aligned}
P(A|N(t)=n) &= \frac{P(A, N(T)=n)}{P(N(T)=n)} \\
&= \frac{\lambda^n e^{-\lambda T} \prod h_i}{(\lambda T)^n e^{-\lambda T}/n!} \\
&= \frac{n! \prod h_i}{T^n}
\end{aligned}
$$

- Divide by $\prod h_i$ and let all h_i go to 0 to get joint density of $S_1, \ldots, S_n$ is

$$\frac{n!}{T^n} 1(0 < s_1 < \cdots < s_n < T);$$

density of order statistics from Uniform$[0, T]$ sample of size n.

Indications of some proofs, X

5) Replace the event $S_{n+1} = T$ with $T < S_{n+1} < T + h$.

- With A as before we want

$$P(A|T < S_{n+1} < T + h) = \frac{P(B, N(T, T+h] \geq 1)}{P(T < S_{n+1} < T + h)}$$

- Note that B is independent of $\{N(T, T+h] \geq 1\}$ and that we have already found the limit

$$\frac{P(B)}{\prod h_i} \to \lambda^n e^{-\lambda T}$$

- We are left to compute the limit of

$$\frac{P(N(T, T+h] \geq 1)}{P(T < S_{n+1} < T + h)}$$

- The denominator is

$$\sum_{k=0}^{n} P(N(0, T] = k, N(T, T+h] = n+1-k) + o(h)$$

$$= P(N(0, T] = n)\lambda h + o(h)$$

Indications of some proofs, XI

- Thus

$$\frac{P(N(T, T+h] \geq 1)}{P(T < S_{n+1} < T+h)} = \frac{\lambda h + o(h)}{\frac{(\lambda T)^n}{n!} e^{-\lambda T} \lambda h + o(h)}$$
$$\to \frac{n!}{(\lambda T)^n e^{-\lambda T}}$$

- This gives the conditional density of $S_1, \ldots, S_n$ given $S_{n+1} = T$ as in **4)**.

Inhomogeneous Poisson Processes

- The idea of hazard rate can be used to extend the notion of Poisson Process.
- Suppose $\lambda(t) \geq 0$ is a function of t.
- Suppose N is a counting process such that

$$P(N(t+h) = k+1 | N(t) = k, H_t) = \lambda(t)h + o(h)$$

and

$$P(N(t+h) \geq k+2 | N(t) = k, H_t) = o(h)$$

Inhomogeneous Poisson Processes

- Then N has independent increments and $N(t+s) - N(t)$ has a Poisson distribution with mean

$$\int_t^{t+s} \lambda(u)du$$

- If we put

$$\Lambda(t) = \int_0^t \lambda(u)du$$

then mean of $N(t+s) - N(t)$ is $\Lambda(t+s) - \Lambda(t)$.

- Jargon: λ is the **intensity** or **instantaneous intensity** and Λ the **cumulative intensity**.
- Can use the model with Λ any non-decreasing right continuous function, possibly without a derivative. This allows ties.

Compound Poisson Processes

- Imagine insurance claims arise at times of a Poisson process, $N(t)$, (more likely for an inhomogeneous process).
- Let Y_i be the value of the ith claim associated with the point whose time is S_i.
- Assume that the Y's are independent of each other and of N.
- Let
$$\mu = \mathrm{E}(Y_i) \text{ and } \sigma^2 = \mathrm{var}(Y_i)$$
- Let
$$X(t) = \sum_{i=1}^{N(t)} Y_i$$
be the total claim up to time t.
- We call X a compound Poisson Process.

Useful properties

$$
\begin{aligned}
\mathrm{E}\left\{X(t)|N(t)\right\} &= N(t)\mu \\
\mathrm{Var}\left\{X(t)|N(t)\right\} &= N(t)\sigma^2 \\
\mathrm{E}\left\{X(t)\right\} &= \mu\mathrm{E}\left\{N(t)\right\} \\
&= \mu\lambda t \\
\mathrm{Var}\left\{X(t)\right\} &= \mathrm{Var}\left[\mathrm{E}\left\{X(t)|N(t)\right\}\right] \\
&\qquad + \mathrm{E}\left[\mathrm{Var}\left\{X(t)|N(t)\right\}\right] \\
&= \lambda t\mu^2 + \lambda t\sigma^2
\end{aligned}
$$

(Look at all familiar? See homework.)