

STAT 450: Statistical Theory

The Multivariate Normal Distribution

Def'n: $Z \in R^1 \sim N(0,1)$ iff

$$f_Z(z) = \frac{1}{\sqrt{2\pi}} e^{-z^2/2}$$

Def'n: $Z \in R^p \sim MVN(0, I)$ if and only if $Z = (Z_1, \dots, Z_p)^t$ with the Z_i independent and each $Z_i \sim N(0,1)$.

In this case according to our theorem

$$\begin{aligned} f_Z(z_1, \dots, z_p) &= \prod \frac{1}{\sqrt{2\pi}} e^{-z_i^2/2} \\ &= (2\pi)^{-p/2} \exp\{-z^t z/2\}; \end{aligned}$$

superscript t denotes matrix transpose.

Def'n: $X \in R^p$ has a multivariate normal distribution if it has the same distribution as $AZ + \mu$ for some $\mu \in R^p$, some $p \times p$ matrix of constants A and $Z \sim MVN(0, I)$.

If the matrix A is singular then X will not have a density.

If A is invertible then we can derive the multivariate normal density by the change of variables formula:

$$X = AZ + \mu \Leftrightarrow Z = A^{-1}(X - \mu)$$

$$\frac{\partial X}{\partial Z} = A \quad \frac{\partial Z}{\partial X} = A^{-1}.$$

So

$$\begin{aligned} f_X(x) &= f_Z(A^{-1}(x - \mu)) |\det(A^{-1})| \\ &= \frac{\exp\{-(x - \mu)^t (A^{-1})^t A^{-1} (x - \mu)/2\}}{(2\pi)^{p/2} |\det A|}. \end{aligned}$$

Now define $\Sigma = AA^t$ and notice that

$$\Sigma^{-1} = (A^t)^{-1} A^{-1} = (A^{-1})^t A^{-1}$$

and

$$\det \Sigma = \det A \det A^t = (\det A)^2.$$

Thus f_X is

$$\frac{\exp\{-(x - \mu)^t \Sigma^{-1} (x - \mu)/2\}}{(2\pi)^{p/2} (\det \Sigma)^{1/2}};$$

the $MVN(\mu, \Sigma)$ density.

Note density is the same for all A such that $AA^t = \Sigma$. This justifies the notation $MVN(\mu, \Sigma)$.

For which vectors μ and matrices Σ is this a density? Any μ but if $x \in R^p$ then

$$\begin{aligned} x^t \Sigma x &= x^t A A^t x \\ &= (A^t x)^t (A^t x) \\ &= \sum_1^p y_i^2 \geq 0 \end{aligned}$$

where $y = A^t x$.

Inequality strict except for $y = 0$ which is equivalent to $x = 0$. Thus Σ is a positive definite symmetric matrix.

Conversely, if Σ is a positive definite symmetric matrix then there is a square invertible matrix A such that $AA^t = \Sigma$ so that there is a $MVN(\mu, \Sigma)$ distribution. (A can be found via the Cholesky decomposition, e.g.)

More generally X has MVN distribution if it has the same distribution as $AZ + \mu$ (no restriction that A be non-singular). When A is singular X will not have a density: $\exists a$ such that $P(a^t X = a^t \mu) = 1$; X is confined to a hyperplane. Still true that the distribution of X depends only on the matrix $\Sigma = AA^t$: if $AA^t = BB^t$ then $AZ + \mu$ and $BZ + \mu$ have the same distribution.

Properties of the MVN distribution

1: All margins are multivariate normal: if

$$X = \begin{bmatrix} X_1 \\ X_2 \end{bmatrix}$$

$$\mu = \begin{bmatrix} \mu_1 \\ \mu_2 \end{bmatrix}$$

and

$$\Sigma = \begin{bmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{bmatrix}$$

then $X \sim MVN(\mu, \Sigma) \Rightarrow X_1 \sim MVN(\mu_1, \Sigma_{11})$.

2: All conditionals are normal: the conditional distribution of X_1 given $X_2 = x_2$ is

$$MVN(\mu_1 + \Sigma_{12}\Sigma_{22}^{-1}(x_2 - \mu_2), \Sigma_{11} - \Sigma_{12}\Sigma_{22}^{-1}\Sigma_{21}).$$

3: $MX + \nu \sim MVN(M\mu + \nu, M\Sigma M^t)$: affine transformation of MVN is normal.