

**STAT 801: Mathematical Statistics**

**Normal samples: Distribution Theory**

**Theorem:** Suppose  $X_1, \dots, X_n$  are independent  $N(\mu, \sigma^2)$  random variables. Then

1.  $\bar{X}$  (sample mean) and  $s^2$  (sample variance) independent.
2.  $n^{1/2}(\bar{X} - \mu)/\sigma \sim N(0, 1)$ .
3.  $(n - 1)s^2/\sigma^2 \sim \chi_{n-1}^2$ .
4.  $n^{1/2}(\bar{X} - \mu)/s \sim t_{n-1}$ .

**Proof:** Let  $Z_i = (X_i - \mu)/\sigma$ .

Then  $Z_1, \dots, Z_p$  are independent  $N(0, 1)$ .

So  $Z = (Z_1, \dots, Z_p)^t$  is multivariate standard normal.

Note that  $\bar{X} = \sigma\bar{Z} + \mu$  and  $s^2 = \sum(X_i - \bar{X})^2/(n - 1) = \sigma^2 \sum(Z_i - \bar{Z})^2/(n - 1)$  Thus

$$\frac{n^{1/2}(\bar{X} - \mu)}{\sigma} = n^{1/2}\bar{Z}$$

$$\frac{(n - 1)s^2}{\sigma^2} = \sum(Z_i - \bar{Z})^2$$

and

$$T = \frac{n^{1/2}(\bar{X} - \mu)}{s} = \frac{n^{1/2}\bar{Z}}{s_Z}$$

where  $(n - 1)s_Z^2 = \sum(Z_i - \bar{Z})^2$ .

So: reduced to  $\mu = 0$  and  $\sigma = 1$ .

**Step 1:** Define

$$Y = (\sqrt{n}\bar{Z}, Z_1 - \bar{Z}, \dots, Z_{n-1} - \bar{Z})^t.$$

(So  $Y$  has same dimension as  $Z$ .) Now

$$Y = \begin{bmatrix} \frac{1}{\sqrt{n}} & \frac{1}{\sqrt{n}} & \cdots & \frac{1}{\sqrt{n}} \\ 1 - \frac{1}{n} & -\frac{1}{n} & \cdots & -\frac{1}{n} \\ -\frac{1}{n} & 1 - \frac{1}{n} & \cdots & -\frac{1}{n} \\ \vdots & \vdots & \vdots & \vdots \end{bmatrix} \begin{bmatrix} Z_1 \\ Z_2 \\ \vdots \\ Z_n \end{bmatrix}$$

or letting  $M$  denote the matrix

$$Y = MZ.$$

It follows that  $Y \sim MVN(0, MM^t)$  so we need to compute  $MM^t$ :

$$MM^t = \left[ \begin{array}{c|cccc} 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 - \frac{1}{n} & -\frac{1}{n} & \cdots & -\frac{1}{n} \\ \vdots & -\frac{1}{n} & \ddots & \cdots & -\frac{1}{n} \\ 0 & \vdots & \cdots & & 1 - \frac{1}{n} \end{array} \right]$$

$$= \left[ \begin{array}{c|c} 1 & 0 \\ \hline 0 & Q \end{array} \right].$$

Solve for  $Z$  from  $Y$ :  $Z_i = n^{-1/2}Y_1 + Y_{i+1}$  for  $1 \leq i \leq n - 1$ . Use the identity

$$\sum_{i=1}^n (Z_i - \bar{Z}) = 0$$

to get  $Z_n = -\sum_{i=2}^n Y_i + n^{-1/2}Y_1$ . So  $M$  invertible:

$$\Sigma^{-1} \equiv (MM^t)^{-1} = \left[ \begin{array}{c|c} 1 & 0 \\ \hline 0 & Q^{-1} \end{array} \right]$$

Use change of variables to find  $f_Y$ . Let  $\mathbf{y}_2$  denote vector whose entries are  $y_2, \dots, y_n$ . Note that

$$\mathbf{y}^t \Sigma^{-1} \mathbf{y} = y_1^2 + \mathbf{y}_2^t Q^{-1} \mathbf{y}_2$$

Then

$$\begin{aligned} f_Y(\mathbf{y}) &= (2\pi)^{-n/2} \exp[-\mathbf{y}^t \Sigma^{-1} \mathbf{y} / 2] / |\det M| \\ &= \frac{1}{\sqrt{2\pi}} e^{-y_1^2/2} \times \\ &\quad \frac{(2\pi)^{-(n-1)/2} \exp[-\mathbf{y}_2^t Q^{-1} \mathbf{y}_2 / 2]}{|\det M|} \end{aligned}$$

Note:  $f_Y(\mathbf{y})$  is function of  $y_1$  times a function of  $y_2, \dots, y_n$ .

Thus  $\sqrt{n}\bar{Z}$  is independent of  $Z_1 - \bar{Z}, \dots, Z_{n-1} - \bar{Z}$ .

Since  $s_Z^2$  is a function of  $Z_1 - \bar{Z}, \dots, Z_{n-1} - \bar{Z}$  we see that  $\sqrt{n}\bar{Z}$  and  $s_Z^2$  are independent.

Also, density of  $Y_1$  is a multiple of the function of  $y_1$  in the factorization above. But factor is standard normal density so  $\sqrt{n}\bar{Z} \sim N(0, 1)$ .

First 2 parts done. Third part is a homework exercise.

Derivation of the  $\chi^2$  density:

Suppose  $Z_1, \dots, Z_n$  are independent  $N(0, 1)$ . Define  $\chi_n^2$  distribution to be that of  $U = Z_1^2 + \dots + Z_n^2$ . Define angles  $\theta_1, \dots, \theta_{n-1}$  by

$$\begin{aligned} Z_1 &= U^{1/2} \cos \theta_1 \\ Z_2 &= U^{1/2} \sin \theta_1 \cos \theta_2 \\ &\vdots \\ Z_{n-1} &= U^{1/2} \sin \theta_1 \cdots \sin \theta_{n-2} \cos \theta_{n-1} \\ Z_n &= U^{1/2} \sin \theta_1 \cdots \sin \theta_{n-1} \end{aligned}$$

(Spherical co-ordinates in  $n$  dimensions. The  $\theta$  values run from 0 to  $\pi$  except last  $\theta$  from 0 to  $2\pi$ .) Derivative formulas:

$$\frac{\partial Z_i}{\partial U} = \frac{1}{2U} Z_i$$

and

$$\frac{\partial Z_i}{\partial \theta_j} = \begin{cases} 0 & j > i \\ -Z_i \tan \theta_i & j = i \\ Z_i \cot \theta_j & j < i \end{cases}$$

Fix  $n = 3$  to clarify the formulas.

Use shorthand  $R = \sqrt{U}$ .

Matrix of partial derivatives is

$$\begin{bmatrix} \frac{\cos \theta_1}{2R} & -R \sin \theta_1 & 0 \\ \frac{\sin \theta_1 \cos \theta_2}{2R} & R \cos \theta_1 \cos \theta_2 & -R \sin \theta_1 \sin \theta_2 \\ \frac{\sin \theta_1 \sin \theta_2}{2R} & R \cos \theta_1 \sin \theta_2 & R \sin \theta_1 \cos \theta_2 \end{bmatrix}$$

Find determinant by adding  $2U^{1/2} \cos \theta_j / \sin \theta_j$  times col 1 to col  $j + 1$  (no change in determinant).

Resulting matrix is lower triangular; diagonal entries

$$U^{-1/2} \cos \theta_1 / 2,$$

$$U^{1/2} \cos \theta_2 / \cos \theta_1$$

and

$$U^{1/2} \sin \theta_1 / \cos \theta_2.$$

We multiply these together to get

$$U^{1/2} \sin(\theta_1) / 2$$

(non-negative for all  $U$  and  $\theta_1$ ).

General  $n$ : every term in the first column contains a factor  $U^{-1/2}/2$  while every other entry has a factor  $U^{1/2}$ .

FACT: Multiplying a column in a matrix by  $c$  multiplies the determinant by  $c$ .

SO: Jacobian of the transformation is  $u^{(n-1)/2} u^{-1/2} / 2$  times some function, say  $h$ , which depends only on the angles.

Thus the joint density of  $U, \theta_1, \dots, \theta_{n-1}$  is

$$(2\pi)^{-n/2} \exp(-u/2) u^{(n-2)/2} h(\theta_1, \dots, \theta_{n-1}) / 2$$

To compute the density of  $U$  we must do an  $n - 1$  dimensional multiple integral  $d\theta_{n-1} \cdots d\theta_1$ .

Answer has the form

$$c u^{(n-2)/2} \exp(-u/2)$$

for some  $c$ .

Evaluate  $c$  by making

$$\int f_U(u) du = c \int u^{(n-2)/2} \exp(-u/2) du = 1$$

Substitute  $y = u/2, du = 2dy$  to see that

$$c 2^{n/2} \int y^{(n-2)/2} e^{-y} dy = c 2^{n/2} \Gamma(n/2) = 1$$

CONCLUSION: the  $\chi_n^2$  density is

$$\frac{1}{2\Gamma(n/2)} \left(\frac{u}{2}\right)^{(n-2)/2} e^{-u/2} 1(u > 0).$$

Fourth part: consequence of first 3 parts and definition of  $t_\nu$  distribution.

**Definition:**  $T \sim t_\nu$  if  $T$  has same distribution as

$$Z / \sqrt{U/\nu}$$

where  $Z \sim N(0, 1), U \sim \chi_\nu^2$  and  $Z$  and  $U$  are independent.

Derive density of  $T$  in this definition:

$$\begin{aligned} P(T \leq t) &= P(Z \leq t\sqrt{U/\nu}) \\ &= \int_0^\infty \int_{-\infty}^{t\sqrt{u/\nu}} f_Z(z) f_U(u) dz du \end{aligned}$$

Differentiate wrt  $t$  by differentiating inner integral:

$$\frac{\partial}{\partial t} \int_{at}^{bt} f(x) dx = b f(bt) - a f(at)$$

by fundamental thm of calculus. Hence

$$\frac{d}{dt}P(T \leq t) = \int_0^\infty f_U(u) \left(\frac{u}{\nu}\right)^{1/2} \frac{\exp[-t^2 u/(2\nu)]}{\sqrt{2\pi}} du$$

Plug in

$$f_U(u) = \frac{1}{2\Gamma(\nu/2)} (u/2)^{(\nu-2)/2} e^{-u/2}$$

to get

$$f_T(t) = \frac{\int_0^\infty (u/2)^{(\nu-1)/2} \exp[-u(1+t^2/\nu)/2] du}{2\sqrt{\pi\nu}\Gamma(\nu/2)}.$$

Substitute  $y = u(1+t^2/\nu)/2$ , to get

$$\begin{aligned} dy &= (1+t^2/\nu)du/2 \\ (u/2)^{(\nu-1)/2} &= [y/(1+t^2/\nu)]^{(\nu-1)/2} \end{aligned}$$

leading to

$$f_T(t) = \frac{(1+t^2/\nu)^{-(\nu+1)/2}}{\sqrt{\pi\nu}\Gamma(\nu/2)} \int_0^\infty y^{(\nu-1)/2} e^{-y} dy$$

or

$$f_T(t) = \frac{\Gamma((\nu+1)/2)}{\sqrt{\pi\nu}\Gamma(\nu/2)} \frac{1}{(1+t^2/\nu)^{(\nu+1)/2}},$$