STAT 804: Notes on Lecture 3

Definition: If $\{\epsilon_t\}$ is a white noise series and μ and $b_0, \ldots, b_p$ are constants then

$$X_t = \mu + b_0\epsilon_t + b_1\epsilon_{t-1} + \cdots + b_p\epsilon_{t-p}$$

is a moving average of order p; we write $MA(p)$.
Question: From observations on X can we estimate the b's and $\sigma^2 = \mathrm{Var}(\epsilon_t)$ accurately? NO.
Definition: A **model** for data X is a family $\{P_\theta; \theta \in \Theta\}$ of possible distributions for X.
Definition: A model is **identifiable** if $\theta_1 \neq \theta_2$ implies that $P_{\theta_1} \neq P_{\theta_2}$; that is different θ's give different distributions for the data.

When a model is unidentifiable there are different values of θ which make exactly the same predictions about the data so the data do not permit you to distinguish between these θ values.
Example: Suppose ϵ is an iid $N(0, \sigma^2)$ series and that $X_t = b_0\epsilon_t + b_1\epsilon_{t-1}$. Then the series X has mean 0 and covariance

$$C_X(h) = \begin{cases} (b_0^2 + b_1^2)\sigma^2 & h = 0 \\ b_0 b_1 \sigma^2 & h = 1 \\ 0 & \text{otherwise} \end{cases}$$

Now a normal distribution is specified by its mean and its variance so two normal time series with mean 0 and the same covariance function have the same distribution. You can see that if you multiply the ϵ's by a and divide both b_0 and b_1 by a then the covariance function of X is unchanged. Thus we cannot hope to estimate all three parameters, b_0, b_1 and σ. We choose to set the parameter b_0 to be 1. Now are the parameters b_1 and σ identifiable? We try to solve the equations

$$C(0) = (1 + b^2)\sigma^2$$

and

$$C(1) = b\sigma^2$$

to see if the solution is unique. Divide the two equations to see

$$\frac{C(1)}{C(0)} = \frac{b}{1 + b^2}$$

or

$$b^2 - \frac{C(0)}{C(1)}b + 1 = 0$$

which has the solutions

$$\frac{\frac{C(0)}{C(1)} \pm \sqrt{\left(\frac{C(0)}{C(1)}\right)^2 - 4}}{2}$$

You should notice two things:

1. If

$$\left|\frac{C(0)}{C(1)}\right| < 2$$

there are no solutions. Since $C(0) = \sqrt{\text{Var}(X_t)\text{Var}(X_{t+1})}$ we can see that $C(1)/C(0)$ is the correlation between X_t and X_{t+1}. We have proved that for an $MA(1)$ process this correlation cannot be more than $1/2$ in absolute value.

2. If

$$\left|\frac{C(0)}{C(1)}\right| > 2$$

there are two solutions.

The two solutions multiply together to give the constant term 1 in the quadratic equation. If the two roots are distinct it follows that one of them is larger than 1 and the other smaller in absolute value. Let b and b^* denote the two roots. Let $\alpha = C(1)/b$ and $\alpha^* = C(1)/b^*$. Let ϵ_t be iid $N(0, \alpha)$ and ϵ_t^* be iid $N(0, \alpha^*)$. Then

$$X_t \equiv \epsilon_t + b\epsilon_{t-1}$$

and

$$X_t^* \equiv \epsilon_t + b^*\epsilon_{t-1}^*$$

have identical means and covariance functions. Observing X_t you cannot distinguish the first of these models from the second. We will fit $MA(1)$ models by **requiring** our estimated b to have $|\hat{b}| \leq 1$.

Reason: We can manipulate the model equation for X just as we did for and autoregressive process last time:

$$\begin{aligned}\epsilon_t &= X_t - b\epsilon_{t-1} \\ &= X_t - b(X_{t-1} - b\epsilon_{t-2}) - b\epsilon_{t-1} \\ &\vdots \\ &= \sum_0^\infty (-b)^j X_{t-j}\end{aligned}$$

This manipulation makes sense if $|b| < 1$. If so then we can rearrange the equation to get

$$X_t = \epsilon_t - \sum_1^\infty (-b)^j X_{t-j}$$

which is an autoregressive process.

If, on the other hand, $|b| > 1$ then we can write

$$X_t = b(\frac{1}{b}\epsilon_t - \epsilon_{t-1})$$

Let $\epsilon_t^* = \epsilon_t/b$; ϵ^* is also white noise. We find

$$\begin{aligned}\epsilon_{t-1}^* &= X_t - \frac{1}{b}\epsilon_t^* \\ &= X_t - \frac{1}{b}(X_{t+1} - \frac{1}{b}\epsilon_{t+1}^*) \\ &\vdots \\ &= \sum_0^\infty (-\frac{1}{b})^j X_{t+j}\end{aligned}$$

which means

$$X_t = \epsilon_{t-1}^* - \sum_1^\infty (-\frac{1}{b})^j X_{t+j}$$

This represents the current value as depending on the future which seems physically far less natural than the other choice.

Definition: An $MA(p)$ process is invertible if it can be written in the form

$$X_t = \sum_1^\infty a_j X_{t-j} + \epsilon_t$$

Definition: A process X is an autoregression of order p (written $AR(p)$) if

$$X_t = \sum_1^p a_j X_{t-j} + \epsilon_t$$

(so an invertible MA is an infinite order autoregression).
Definition: The backshift operator transforms a time series into another time series by shifting it back one time unit; if X is a time series then BX is the time series with

$$(BX)_t = X_{t-1} .$$

The identity operator I satisfies $IX = X$. We use B^j for $j = 1, 2, \ldots$ to denote B composed with itself j times so that

$$(B^j X)_t = X_{t-j}$$

For $j = 0$ this gives $B^0 = I$.

Now we use B to develop a formal method for studying the existence of a given $AR(p)$ and the invertibility of a given $MA(p)$. An $AR(1)$ process satisfies

$$(I - a_1 B)X = \epsilon$$

If you think of $I - a_1 B$ as some sort of infinite dimensional matrix then you get the formal identity

$$X = (I - a_1 B)^{-1} \epsilon$$

So how will we define this inverse of an infinite matrix? We use the idea of a geometric series expansion.

If b is a real number then

$$(1 - ab)^{-1} = \frac{1}{1 - ab} = \sum_{j=0}^{\infty} (ab)^j$$

so we hope that $(I - a_1 B)^{-1}$ can be defined by

$$(I - a_1 B)^{-1} = \sum_{j=0}^{\infty} a_1^j B^j$$

This would mean

$$X = \sum_{j=0}^{\infty} a_1^j B^j \epsilon$$

or looking at the formula for a particular t and remembering the meaning of B^j we get

$$X_t = \sum_{j=0}^{\infty} a_1^j \epsilon_{t-j}$$

This is the formula I had in lecture 2.

Now consider a general $AR(p)$ process:

$$(I - \sum_1^p a_j B^j) X = \epsilon$$

We will factor the operator applied to x. Let

$$\phi(x) = 1 - \sum_1^p a_j x^j$$

Then ϕ is a polynomial of degree p. It thus has (a theorem of C. F. Gauss) p roots $1/b_1, \ldots, 1/b_p$. (None of the roots is 0 because the constant term in ϕ is 1.) This means we can factor ϕ as

$$\phi(x) = \prod_1^p (1 - b_j x)$$

Now back to the definition of X:

$$\prod_1^p (I - b_j B) X = \epsilon$$

can be solved by inverting each term in the product (in any order — the terms in the product commute) to get

$$X = \prod_1^p (I - b_j B)^{-1} \epsilon$$

The inverse of $I - b_1 B$ will exist if the sum

$$\sum_{k=0}^{\infty} b_j^k B^k$$

converges; this requires $|b_j| < 1$. Thus a stationary $AR(p)$ solution of the equations exists if every root of the characteristic polynomial ϕ is larger than 1 in absolute value (actually the roots can be complex and I mean larger than 1 in modulus).

Summary

- An $MA(q)$ process $X_t = \epsilon_t - \sum_{j=1}^{q} b_j \epsilon_{t-j}$ is invertible if and only if all roots of the characteristic polynomial $\psi(x) = 1 - \sum_{j=1}^{q} b_j x^j$ lie outside the unit circle in the complex plain.

- For a given covariance function of an $MA(q)$ process there is only one set of coefficients $b_1, \ldots, b_q$ for which the process is invertible.

- An $AR(p)$ process $X_t - \sum_{j=1}^{q} a_j X_{t-j} = \epsilon_t$ is asymptotically stationary if and only if all roots of the characteristic polynomial $\phi(x) = 1 - \sum_{1}^{p} a_j x^j$ lie outside the unit circle in the complex plain.

  (Asymptotically stationary means this: if you make $X_{-1}, X_{-2}, \ldots, X_{-p}$ anything at all and use the equation defining the $AR(p)$ to define all the rest of the X values then as $t \to \infty$ the process gets closer to being stationary. The assertion of asymptotic stationarity is equivalent here to the existence of an exactly stationary solution of the equations.)

Definition: A process X is an $ARMA(p, q)$ (mixed autoregressive of order p and moving average of order q) if it satisfies

$$\phi(B)X = \psi(B)\epsilon$$

where ϵ is white noise and

$$\phi(B) = I - \sum_{1}^{p} a_j B^j$$

and

$$\psi(B) = I - \sum_{1}^{p} b_j B^j$$

The ideas we used above can be stretched to show that the process X is identifiable and causal (can be written as an infinite order autoregression on the past) if the roots of $\psi(x)$ lie outside the unit circle. A stationary solution, which can be written as an infinite order causal (no future ϵs in the average) moving average, exists if all the roots of $\phi(x)$ lie outside the unit circle.
Other Stationary Processes:

1. Periodic processes. Suppose Z_1 and Z_2 are independent $N(0, \sigma^2)$ random variables and that ω is a constant. Then

$$X_t = Z_1 \cos(\omega t) + Z_2 \sin(\omega t)$$

has mean 0 and

$$\begin{aligned} \text{Cov}(X_t, X_{t+h}) &= \sigma^2 \left[\cos(\omega t)\cos(\omega(t+h)) + \sin(\omega t)\sin(\omega(t+h))\right] \\ &= \sigma^2 \cos(\omega h) \end{aligned}$$

Since X is Gaussian we find that X is second order and strictly stationary. In fact (see your homework) You can write

$$X_t = R \sin(\omega t + \Phi)$$

where R and Φ are suitable random variables so that the trajectory of X is just a sine wave.

2. Poisson shot noise processes:

A Poisson process is a process $N(A)$ indexed by subsets A of the real line with the property that each $N(A)$ has a Poisson distribution with parameter $\lambda \text{length}(A)$ and if $A_1, \ldots A_p$ are any non-overlapping subsets of R then $N(A_1), \ldots, N(A_p)$ are independent. We often use $N(t)$ for $N([0,t])$.

To define a shot noise process we let $X(t) = 1$ at those t where there is a jump in N and 0 elsewhere. The process X is stationary. If we have some function g defined on $[0, \infty)$ and decreasing sufficiently quickly to 0 (like say $g(x) = e^{-x}$) then the process

$$Y(t) = \sum g(t - \tau) 1(X(\tau) = 1) 1(\tau \leq t)$$

is stationary. It has a jump every time t passes a jump in the Poisson process and otherwise follows the trajectory of the sum of several copies of g (shifted around in time). We commonly write

$$Y(t) = \int_0^\infty g(t - \tau) dN(\tau)$$