

# STAT 830

## Confidence Sets

Richard Lockhart

Simon Fraser University

STAT 890 = 830 — Fall 2011



# Purposes of These Notes

- Discuss exact and approximate confidence intervals.
- Discuss role of pivots in finding confidence intervals.



# Confidence Intervals

- **Def'n:** A level  $\beta$  confidence set for a parameter  $\phi(\theta)$  is a random subset  $C$ , of the set of possible values of  $\phi$  such that for each  $\theta$

$$P_{\theta}(\phi(\theta) \in C) \geq \beta$$

- Confidence sets are very closely connected with hypothesis tests:
- First from confidence sets to hypothesis tests.
- Suppose  $C$  is a level  $\beta = 1 - \alpha$  confidence set for  $\phi$ .
- To test  $\phi = \phi_0$ : reject if  $\phi \notin C$ .
- This test has level  $\alpha$ .



## From tests to confidence sets

- Conversely, suppose that for each  $\phi_0$  we have available a level  $\alpha$  test of  $\phi = \phi_0$  whose rejection region is say  $R_{\phi_0}$ .
- Define  $C = \{\phi_0 : \phi = \phi_0 \text{ is not rejected}\}$ ; get level  $1 - \alpha$  confidence set for  $\phi$ .
- **Example:** Usual  $t$  test gives rise in this way to the usual  $t$  confidence intervals

$$\bar{X} \pm t_{n-1, \alpha/2} \frac{s}{\sqrt{n}}.$$



## Confidence sets from Pivots

- **Def'n:** A **pivot** (pivotal quantity) is a function  $g(\theta, X)$  whose distribution is the same for all  $\theta$ .
- Note  $\theta$  in pivot is same  $\theta$  as being used to calculate distribution of  $g(\theta, X)$ .
- Using pivots to generate confidence sets:
- Pick a set  $A$  in space of possible values for  $g$ .
- Let  $\beta = P_{\theta}(g(\theta, X) \in A)$ ; since  $g$  is pivotal  $\beta$  is the same for all  $\theta$ .
- Given data  $X$  solve the relation

$$g(\theta, X) \in A$$

to get

$$\theta \in C(X, A).$$



## Example: Normal variance interval

- Note  $(n-1)s^2/\sigma^2 \sim \chi_{n-1}^2$  is pivot in  $N(\mu, \sigma^2)$  model.
- Given  $\beta = 1 - \alpha$  consider the two points

$$\chi_{n-1, 1-\alpha/2}^2 \text{ and } \chi_{n-1, \alpha/2}^2.$$

- Then

$$P(\chi_{n-1, 1-\alpha/2}^2 \leq (n-1)s^2/\sigma^2 \leq \chi_{n-1, \alpha/2}^2) = \beta$$

for all  $\mu, \sigma$ .

- Solve this relation:

$$P\left(\frac{(n-1)^{1/2}s}{\chi_{n-1, \alpha/2}} \leq \sigma \leq \frac{(n-1)^{1/2}s}{\chi_{n-1, 1-\alpha/2}}\right) = \beta$$

so interval

$$\left[ \frac{(n-1)^{1/2}s}{\chi_{n-1, \alpha/2}}, \frac{(n-1)^{1/2}s}{\chi_{n-1, 1-\alpha/2}} \right]$$

is a level  $1 - \alpha$  confidence interval.



## Other intervals

- In the same model we also have

$$P(\chi_{n-1,1-\alpha}^2 \leq (n-1)s^2/\sigma^2) = \beta$$

which can be solved to get

$$P(\sigma \leq \frac{(n-1)^{1/2}s}{\chi_{n-1,1-\alpha}}) = \beta$$

- This gives a level  $1 - \alpha$  interval

$$(0, (n-1)^{1/2}s/\chi_{n-1,1-\alpha}).$$

- Right hand end of interval usually called *confidence upper bound*.
- In general the interval from

$$(n-1)^{1/2}s/\chi_{n-1,\alpha_1} \text{ to } (n-1)^{1/2}s/\chi_{n-1,1-\alpha_2}$$

has level  $\beta = 1 - \alpha_1 - \alpha_2$ .

- For fixed  $\beta$  can minimize length of interval numerically — rarely used
- See homework for an example.

