


A Few Surprises Yet in Steady 2D Topographic Wave Flows

- ▷ nonlinearity & rotational influences on wave generation
- ▷ a rotating version of Long's theory




Epifanio

- ▷ Dave Muraki (Simon Fraser University)
- ▷ Craig Epifanio (Texas A&M)
- ▷ Chris Snyder (NCAR Boulder)


Linear Theory: Tiny Rossby Number

Quasigeostrophic Flow Over A Ridge

- ▷ small height gaussian ridge ($\mathcal{A} = NH/U = 0.25$)
- ▷ predominantly balanced QG flow ($\mathcal{R} = U/fL = 0.25$)
- ▷ very weak wave anomalies near leeward surface (Pierrehumbert, 1984)




Regimes for 2D Steady Topographic Flows



Linear Theory: Small Rossby Number

Appearance of Waves


- ▷ steady uniform flow, constant stratification
- ▷ intermediate case: QG summit flow with short waves ($\mathcal{R} = 0.50$)
- ▷ development of downstream (dispersive) wavetrain



Linear Theory: Intermediate Rossby Number

Fully Developed Wave Field

- ▷ strong waves with similar scale to QG summit flow ($\mathcal{R} = 1.0$)
- ▷ significant wave radiation aloft



- ▷ as \mathcal{R} ↗, waves grow in amplitude (exponentially) & wavelength (linearly)

Linear Theory: A Singular Numerical Problem

Fourier Integral Solution (Queney, 1947)

$$b(x, z) = -\frac{N^2}{\pi} \text{Real} \left\{ \int_0^\infty \hat{h}(k) e^{ikx} e^{m(k)z} dk \right\}$$



Buoyancy Anomaly

▷ linear waves with **rotation**, **stratification** & topography $h(x)$

$$A^2 b_{xx} + \mathcal{R}^{-2} b_{zz} + b_{xxz} = 0 \quad ; \quad b(x, 0) = -h(x)$$

▷ 2D linear dispersion relation gives a **singular exponent** at $k = \mathcal{R}^{-1}$

$$m(k) = \begin{cases} -\frac{Ak}{\sqrt{\mathcal{R}^{-2} - k^2}} & \text{for } 0 \leq k < \mathcal{R}^{-1} \quad (\text{vertical decay}) \\ i \frac{Ak}{\sqrt{\mathcal{R}^{-2} - k^2}} & \text{for } \mathcal{R}^{-1} < k < \infty \quad (\text{outgoing waves}) \end{cases}$$



djm

▷ rotating wave case prone to severe numerical Fourier errors

Three Questions

a: Is There an Analog to Long's Theory that includes Coriolis Rotation?

- ▷ Long's theory (1953) for buoyancy anomaly
- ▷ steady, **nonlinear** & **non-rotating** flows are obtained exactly via linear Helmholtz solutions
 - no obvious extension to include **rotation**

b: What is the Nature of Pierrehumbert's Finite \mathcal{R} Singularity?

- ▷ semi-geostrophic approximation: Pierrehumbert (1985)
- ▷ SG solutions have singular breakdown at finite Rossby number
 - a true **finite amplitude** flow transition, or merely a manifestation of SG approximation?

c: How can Waves be Generated at Small Rossby Number?



- ▷ Pierrehumbert/Wyman (1985) & Trüb/Davies (1995)
- ▷ wave generation by finite amplitude ridges at small \mathcal{R}
- ▷ relaxation of time-dependent flow computations
 - how does **nonlinearity** circumvent quasigeostrophic balance?

a: Long's Theory for Non-Rotating Topographic Waves

An Exact Nonlinear Theory for Buoyancy

- ▷ steady, **non-rotating** & hydrostatic/nonhydrostatic (Long, 1953)
- ▷ 2D helmholtz equation: stratified ($\mathcal{A} = NH/U$) & nonhydrostatic (δ^2)

$$\mathcal{A}^2 b + b_{zz} + \delta^2 \mathcal{A}^2 b_{xx} = 0 \quad ; \quad b(x, h(x)) = 0$$
- ▷ downstream waves derive from radiation boundary conditions
 - except hydrostatic waves ($\delta^2=0$) are nondispersive



- ▷ nonlinear fluid system reduces to a single equation for buoyancy

Isentropic Coordinates

2D Primitive Equations

- ▷ nondimensional: steady, **rotating** & nonhydrostatic
- ▷ potential temperature θ as vertical coordinate ($\theta_z = 1/z\theta$)
$$\begin{aligned} \mathcal{D}u & - \mathcal{R}^{-1}v & = & -\mathcal{A}^2 M_x & - \delta^2 z_x \mathcal{D}w \\ \mathcal{D}v & + \mathcal{R}^{-1}u & = & \mathcal{R}^{-1} & \\ \delta^2 z_\theta \mathcal{D}w & + \mathcal{A}^2 z & = & -\mathcal{A}^2 M_\theta & \\ \mathcal{D}z & - w & = & 0 & \end{aligned}$$
- ▷ Montgomery potential: $M = \phi - z\theta$
- ▷ steady 2D advection: $\mathcal{D} = u \partial_x$; incident wind $u^\infty = 1$
- ▷ 2D divergence: $z_\theta u_x - z_x u_\theta + w_\theta = 0$

Steady Streamline Property

- ▷ divergence + thermodynamic $\rightarrow \{u z_\theta\}_x = 0$
 - \rightarrow squeezing isentropes (streamlines) accelerates flow
- ▷ velocity relations: $u = 1/z_\theta$; $w = z_x/z_\theta$
- ▷ across-ridge flow: $v_x = \mathcal{R}^{-1} (z_\theta - 1)$
- ▷ eliminating M through vorticity . . . then a miracle happens . . .

A Master Equation for Buoyancy

Vertical Displacement Equation

- ▷ includes both **f-plane** and **non-hydrostatic** effects
$$A^2 z_{xx} + \mathcal{R}^{-2} z_{\theta\theta} - \eta_{xx} = 0 \quad ; \quad \eta = \frac{1}{2} (u^2 + \delta^2 w^2)_\theta - \delta^2 w_x$$
- ▷ surface condition: $z(x, 0) = h(x)$ & radiation BCs
- ▷ nonlinearity in horizontal vorticity η
- ▷ equivalent to Long's equation without rotation ($\mathcal{R}^{-2} \rightarrow 0$)

Hydrostatic Buoyancy Equation ($\delta^2 = 0$)

- ▷ constant stratification: $z = \theta - b(x, \theta)$

$$A^2 b_{xx} + \mathcal{R}^{-2} b_{\theta\theta} + \left\{ u^3 b_{\theta\theta} \right\}_{xx} = 0 \quad ; \quad u = \frac{1}{1 - b_\theta}$$

- ▷ surface condition: $b(x, 0) = -h(x)$ & radiation BCs
- ▷ linearizing ($u \sim 1$) recovers queney

b: Nonlinear Flows

Isentropic Coordinate Singularities

- ▷ breakdowns in coordinate inversion of $z = \theta - b(x, \theta)$

$$\theta_z = \frac{1}{z_\theta} = u = \frac{1}{1 - b_\theta} \rightarrow \begin{cases} \infty & \text{isentropes collapsing, } u \rightarrow \infty \\ 0 & \text{isentropes overturning, } u \rightarrow 0 \end{cases}$$

Semi-geostrophic Approximation

- ▷ small \mathcal{R} extension of quasigeostrophy: Robinson (1960), Pierrehumbert (1985)
- ▷ SG truncation of *hydrostatic master equation*
$$\mathcal{A}^2 b_{xx} + \mathcal{R}^{-2} b_{\theta\theta} = 0 \quad ; \quad b(x, 0) = -h(x)$$
- ▷ isentropes collapse must occur above $h(x)$ -dependent critical value of $\mathcal{R}\mathcal{A}$


Enhanced Wave Generation & Singularity Suppression?

- ▷ approach to collapse invalidates SG approximation, as nonlinearity u^3 must become large
- ▷ seems nonlinearity suppresses collapse singularity through enhanced wave generation

c: Nonlinear Waves at Tiny Rossby Number

Nonlinear Wave Generation

- ▷ moderate height gaussian ridge ($\mathcal{A} = NH/U = 1.00$)
- ▷ tiny Rossby number flow ($\mathcal{R} = U/fL = 0.25$)
- ▷ time-transient computation to steady state



- ▷ how are these waves generated?

Epifanio


Direct Steady Solve

Solution at $\mathcal{R} = 0.25$; $\mathcal{A} = 1.00$


▷ iterate on nonlinearity in *hydrostatic master equation*: $u^{old} \rightarrow b^{new} \rightarrow u^{new}$

$$\mathcal{A}^2 b_{xx}^n + \mathcal{R}^{-2} b_{\theta\theta}^n + \left\{ (u^o)^3 b_{\theta\theta}^n \right\}_{xx} = 0 \quad ; \quad u^n = \frac{1}{1 - b_{\theta}^n}$$


linear solution: $b^0(x, \theta)$



waves after convergent iterations: $b(x, \theta)$



Streamline Comparison



Epifanio & djm


Possible Nonlinear Mechanisms

- ▷ nonlinear modification of local Rossby number
 - enhanced topographic wave generation at ridge summit
 - modification of wave propagation (rays) in interior
- ▷ nonlinear wave generation in interior?

$$A^2 b_{xx}^n + \mathcal{R}^{-2} b_{\theta\theta}^n + b_{xx\theta\theta}^n = - \left(((u^o)^3 - 1) b_{\theta\theta}^o \right)_{xx}$$

Generation/Enhancement/Refraction









Nonlinear Corrections



▷ 1st correction:
→ *new waves*

▷ remaining corrections:
→ refraction by u^{QG}

Other Fields





djm

Nonlinear Waves at Small & Moderate Rossby Number ---

Nonlinear Wave Enhancement

- ▷ moderate height gaussian ridge ($\mathcal{A} = 1.00$)
- ▷ Rossby number flows ($\mathcal{R} = 0.50, 1.00$)
- ▷ time-transient computation to steady state



Epifanio


- ▷ wave amplitudes approach overturning as \mathcal{R} ↗

Summary


Master Equation for Buoyancy

- ▷ single equation for 2D topographic wave flow spanning non-hydrostatic to QG regimes
- ▷ quantitative tool for understanding nonlinear wave processes
- ▷ key issue: stability & accuracy of numerical solves

one iteration at $\mathcal{R} = 0.50$



time-dependent computation



Epifanio & djm