

73. Show that the function  $z = xe^y + ye^x$  is a solution of the equation

$$\frac{\partial^2 z}{\partial x^2} + \frac{\partial^2 z}{\partial y^2} = x \frac{\partial^2 z}{\partial x \partial y^2} + y \frac{\partial^2 z}{\partial x^2 \partial y}$$

74. Show that the Cobb-Douglas production function  $P = bL^\alpha K^\beta$  satisfies the equation

$$L \frac{\partial P}{\partial L} + K \frac{\partial P}{\partial K} = (\alpha + \beta)P$$

75. Show that the Cobb-Douglas production function satisfies  $P(L, K_0) = C_1(K_0)L^\alpha$  by solving the differential equation

$$\frac{dP}{dL} = \alpha \frac{P}{L}$$

(See Equation 5.)

76. The temperature at a point  $(x, y)$  on a flat metal plate is given by  $T(x, y) = 60/(1 + x^2 + y^2)$ , where  $T$  is measured in  $^\circ\text{C}$  and  $x, y$  in meters. Find the rate of change of temperature with respect to distance at the point  $(2, 1)$  in (a) the  $x$ -direction and (b) the  $y$ -direction.

77. The total resistance  $R$  produced by three conductors with resistances  $R_1, R_2, R_3$  connected in a parallel electrical circuit is given by the formula

$$\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3}$$

Find  $\partial R / \partial R_1$ .

78. The gas law for a fixed mass  $m$  of an ideal gas at absolute temperature  $T$ , pressure  $P$ , and volume  $V$  is  $PV = mRT$ , where  $R$  is the gas constant. Show that

$$\frac{\partial P}{\partial V} \frac{\partial V}{\partial T} \frac{\partial T}{\partial P} = -1$$

79. For the ideal gas of Exercise 78, show that

$$T \frac{\partial P}{\partial T} \frac{\partial V}{\partial T} = mR$$

80. The wind-chill index is modeled by the function

$$W = 13.12 + 0.6215T - 11.37v^{0.16} + 0.3965Tv^{0.16}$$

where  $T$  is the temperature ( $^\circ\text{C}$ ) and  $v$  is the wind speed (km/h). When  $T = -15^\circ\text{C}$  and  $v = 30$  km/h, by how much would you expect the apparent temperature to drop if the actual temperature decreases by  $1^\circ\text{C}$ ? What if the wind speed increases by 1 km/h?

81. The kinetic energy of a body with mass  $m$  and velocity  $v$  is  $K = \frac{1}{2}mv^2$ . Show that

$$\frac{\partial K}{\partial m} \frac{\partial K}{\partial v^2} = K$$

82. If  $a, b, c$  are the sides of a triangle and  $A, B, C$  are the opposite angles, find  $\partial A / \partial a, \partial A / \partial b, \partial A / \partial c$  by implicit differentiation of the Law of Cosines.

83. You are told that there is a function  $f$  whose partial derivatives are  $f_x(x, y) = x + 4y$  and  $f_y(x, y) = 3x - y$ . Should you believe it?



84. The paraboloid  $z = 6 - x - x^2 - 2y^2$  intersects the plane  $x = 1$  in a parabola. Find parametric equations for the tangent line to this parabola at the point  $(1, 2, -4)$ . Use a computer to graph the paraboloid, the parabola, and the tangent line on the same screen.

85. The ellipsoid  $4x^2 + 2y^2 + z^2 = 16$  intersects the plane  $y = 2$  in an ellipse. Find parametric equations for the tangent line to this ellipse at the point  $(1, 2, 2)$ .

86. In a study of frost penetration it was found that the temperature  $T$  at time  $t$  (measured in days) at a depth  $x$  (measured in feet) can be modeled by the function

$$T(x, t) = T_0 + T_1 e^{-\lambda x} \sin(\omega t - \lambda x)$$

where  $\omega = 2\pi/365$  and  $\lambda$  is a positive constant.

- (a) Find  $\partial T / \partial x$ . What is its physical significance?  
 (b) Find  $\partial T / \partial t$ . What is its physical significance?  
 (c) Show that  $T$  satisfies the heat equation  $T_t = k T_{xx}$  for a certain constant  $k$ .



- (d) If  $\lambda = 0.2, T_0 = 0$ , and  $T_1 = 10$ , use a computer to graph  $T(x, t)$ .  
 (e) What is the physical significance of the term  $-\lambda x$  in the expression  $\sin(\omega t - \lambda x)$ ?

87. Use Clairaut's Theorem to show that if the third-order partial derivatives of  $f$  are continuous, then

$$f_{xyy} = f_{yyx} = f_{yxy}$$

88. (a) How many  $n$ th-order partial derivatives does a function of two variables have?  
 (b) If these partial derivatives are all continuous, how many of them can be distinct?  
 (c) Answer the question in part (a) for a function of three variables.

89. If  $f(x, y) = x(x^2 + y^2)^{-3/2} e^{\sin(x^2 y)}$ , find  $f_x(1, 0)$ . [Hint: Instead of finding  $f_x(x, y)$  first, note that it's easier to use Equation 1 or Equation 2.]

90. If  $f(x, y) = \sqrt[3]{x^3 + y^3}$ , find  $f_x(0, 0)$ .

91. Let

$$f(x, y) = \begin{cases} \frac{x^3 y - xy^3}{x^2 + y^2} & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0) \end{cases}$$



- (a) Use a computer to graph  $f$ .  
 (b) Find  $f_x(x, y)$  and  $f_y(x, y)$  when  $(x, y) \neq (0, 0)$ .  
 (c) Find  $f_x(0, 0)$  and  $f_y(0, 0)$  using Equations 2 and 3.  
 (d) Show that  $f_{xy}(0, 0) = -1$  and  $f_{yx}(0, 0) = 1$ .  
 (e) Does the result of part (d) contradict Clairaut's Theorem? Use graphs of  $f_{xy}$  and  $f_{yx}$  to illustrate your answer.

