

## ORIGINAL ARTICLE

**Turner syndrome and the evolution of human sexual dimorphism**

Bernard Crespi

Department of Biosciences, Simon Fraser University, Burnaby, BC, Canada

**Keywords**

genomic imprinting, sexual dimorphism, Turner syndrome, X chromosome.

**Correspondence**

Bernard Crespi, Killam Research Professor, Department of Biosciences, Simon Fraser University, Burnaby, BC V5A 1S6, Canada. Tel.: 778 782 3533; fax: 778 782 3496; e-mail: crespi@sfu.ca

Received: 19 November 2007

Accepted: 17 January 2008

First published online: 22 February 2008

doi:10.1111/j.1752-4571.2008.00017.x

**Abstract**

Turner syndrome is caused by loss of all or part of an X chromosome in females. A series of recent studies has characterized phenotypic differences between Turner females retaining the intact maternally inherited versus paternally inherited X chromosome, which have been interpreted as evidence for effects of X-linked imprinted genes. In this study I demonstrate that the differences between Turner females with a maternal X and a paternal X broadly parallel the differences between males and normal females for a large suite of traits, including lipid profile and visceral fat, response to growth hormone, sensorineural hearing loss, congenital heart and kidney malformations, neuro-anatomy (sizes of the cerebellum, hippocampus, caudate nuclei and superior temporal gyrus), and aspects of cognition. This pattern indicates that diverse aspects of human sex differences are mediated in part by X-linked genes, via genomic imprinting of such genes, higher rates of mosaicism in Turner females with an intact X chromosome of paternal origin, karyotypic differences between Turner females with a maternal versus paternal X chromosome, or some combination of these phenomena. Determining the relative contributions of genomic imprinting, karyotype and mosaicism to variation in Turner syndrome phenotypes has important implications for both clinical treatment of individuals with this syndrome, and hypotheses for the evolution and development of human sexual dimorphism.

**Introduction**

Genomic syndromes are alterations to a suite of contiguous genes, such as deletions, duplications, or aneuploidies, that result in characteristic sets of phenotypic changes, some of which may require medical interventions (e.g. Feinstein and Singh 2007). Such syndromes provide unique insights into human evolution because they represent naturally occurring genomic variation that can be linked with specific phenotypic consequences for human growth, development and cognition. For example, Haig and Wharton (2003), Oliver et al. (2007) and Crespi and Badcock (2008) show how the phenotypes of Prader-Willi and Angelman syndromes, which are due to diametrical alterations of a region of chromosome 15 bearing a cluster of imprinted genes, provide insight into the evolution of human childhood and mother-offspring interactions.

Similarly, Williams syndrome, caused by deletions of a region of chromosome 7, involves an unusual cognitive profile of spared or enhanced expressive-language skills, but greatly impaired visual-spatial abilities, which has been interpreted as providing insights into the genetic and neurological architecture of human language (Tassabehji 2003; Meyer-Lindenberg et al. 2006; Brock 2007). Duplications of the Williams-syndrome region, by contrast, involve high rates of autism, with expressive language abilities selectively impaired (Berg et al. 2007).

Several genomic syndromes involve gains or loss of entire chromosomes. Loss of part or all of an X chromosome causes Turner syndrome in females, whereas gains of one or more X chromosomes result in Klinefelter syndrome in males (Simpson et al. 2003; Bondy 2006). These syndromes are of particular interest in human evolution

because the X chromosome evolves relatively rapidly and bears a concentration of genes related to reproduction and cognition (Vallender and Lahn 2004; Nielsen et al. 2005). Phenotypes related to reproduction and cognition are indeed notably altered in Turner and Klinefelter syndromes, with both syndromes involving dysregulation of gonadal development and alterations to neurocognitive profiles of verbal versus visual-spatial skills (Money 1993; Simpson et al. 2003; Bondy 2006; Kesler 2007). These findings indicate that studies of Turner and Klinefelter syndromes that integrate approaches from evolutionary biology and medical genetics should provide useful insights into both the developmental-genomic aetiologies of these conditions, and how X-linked genes have been involved in the evolution of modern humans.

In this paper I focus on the causes of phenotypic variation among individuals with Turner syndrome, and between Turner syndrome females, normal females and normal males. Turner syndrome is characterized phenotypically by short stature, gonadal dysgenesis, a range of anatomical stigmata, and a neurocognitive profile of spared or enhanced verbal abilities but impaired visual-spatial and social skills (Sybert and McCauley 2004; Bondy 2006; Kesler 2007). The syndrome is caused by partial or complete loss of one of the two X chromosomes in most or all cells, due to a range of cytogenetic alterations, with most cases associated with either: (i) the absence of one entire X (45,X), resulting in monosomy, (ii) deletion of part of the short, Xp, arm of the X chromosome (46,XdelXp), or (iii) formation of an Xq iso-chromosome (46,XiXq, with two identical arms of Xq and an Xp deletion) (see Bondy 2006 for more detail on karyotypic variation).

Turner females may also be mosaics of 45,X with 46,XX cells, or mosaics of 45,X cells with cells bearing 46,XdelXp, 46,XiXq, 46,XY, or other karyotypes. Estimates of the frequency of mosaicism range from 67% to 90% (Held et al. 1992; Fernández-García et al. 2000), but the presence and degree of mosaicism has been difficult to establish because multiple tissues must be studied and PCR-based methods must be used for accurate quantification, but most studies have relied on karyotype data from single tissues (Fernández-García et al. 2000). Chromosomal mosaicism of the forms 45,X with 46,XX, or 45,X with 46,XdelXp or 46,XiXq, notably mitigates the severity of Turner syndrome phenotypes (e.g. Murphy et al. 1997; El-Mansouri et al. 2007). Turner phenotypes are also mediated in part by preferential inactivation of structurally abnormal X chromosomes, or in some cases by failed or partial X-inactivation (Migeon et al. 1996; Wolff et al. 2000; Leppig and Disteche 2001).

Determining the nature and causes of karyotype–phenotype correlations in Turner syndrome is important both for clinical treatment of this condition, and for understanding the roles of sex-linked genes in human evolution and development. The primary genetic consequences of Turner syndrome aneuploidies, deletions and mosaicism, that may contribute to phenotypic variation between Turner syndrome females and 46,XX females, and among females with this syndrome, are twofold: (i) full or partial haploinsufficiency of noninactivated X-linked genes in pseudoautosomal region 1 (at the terminus of the Xp arm) or elsewhere on this chromosome; and (ii) the presence of a full or fragmentary Y chromosomal in some or all cells (Bondy 2006; Lynn and Davies 2007).

A third source of potential variation in Turner syndrome phenotypes is epigenetic. Given that the intact chromosome in Turner syndrome is inherited from either the father or the mother, imprinting (silencing by parent of origin) of genes may also influence gene expression on the X (Skuse et al. 1997; Skuse 1999, 2000, 2005, 2006; Davies et al. 2006), as it does for many autosomal imprinted genes. A series of studies has tested for phenotypic differences between Turner females with the intact X inherited either maternally or paternally (Hamelin et al. 2006; Bondy et al. 2007; Sagi et al. 2007). Some of these studies have employed small sample sizes, but more-recent and larger studies have demonstrated statistically significant differences for diverse traits, with important implications for genetic diagnosis and clinical treatment.

Skuse (1999) has suggested that X-linked imprinting may serve as a mechanism for the evolution of sexual dimorphism in humans, given that gene dosages of X-linked imprinted genes are expected to differ between the sexes, and a basis in population-genetic theory has been provided for this hypothesis by Iwasa and Pomiankowski (1999) and Mills and Moore (2006). In accordance with these ideas, Skuse et al. (1997) has shown that Turner-syndrome individuals with the maternally inherited X intact (the only X present in normal XY males) differ from paternal-X females in exhibiting a set of relatively male-typical cognitive traits including higher liability to autism.

Do other traits exhibit a similar pattern, of normal sex differences mirroring differences between Turner females with an intact maternal X (Xmat) versus an intact paternal X (Xpat)? If so, can these differences be ascribed to effects of X-linked imprinting, or to correlates of the parental origin of the X chromosome such as mosaicism or karyotype, given that dosages of noninactivated X-linked genes may also mediate human sexual dimorphism? Sufficient data are available to evaluate these patterns for seven categories of phenotype.

## Phenotypic differences between Xmat and Xpat females with Turner syndrome

In comparing the results from multiple studies that assess the same phenotypic trait in Turner syndrome females with an intact maternal versus paternal X chromosome (Table 1), it is important to recognize that the different studies have used different clinical populations; for example, Tsezou et al. (1999) and Bondy et al. (2007) included 45X/46,XX mosaics, whereas Hamelin et al. (2006) and Sagi et al. (2007) excluded them, and Sagi et al. (2007) included only females with 45,X or isodicentric karyotypes. This source of among-study variation, in conjunction with variation between and within studies in sample sizes and methods of quantifying phenotypes (e.g. Sagi et al. 2007), means that it is difficult to interpret failures of replication in terms of the presence or absence of biological effects. For each trait, data are compiled on patterns of concordance between differences between Xmat versus Xpat Turner females, differences between 45,X and other Turner females, and differences between males and females in normal populations (Table 1).

### Response to growth hormone

Females with Turner syndrome exhibit reduced adult stature that can be prevented in part via treatment with growth hormone (Sybert and McCauley 2004; Bondy 2006). Tsezou et al. (1999) found no significant difference between Xmat and Xpat Turner females in growth-hormone-stimulated height gain over 2 years, but Hamelin et al. (2006) reported significantly greater gain in height among Xmat than Xpat Turner females, over 5–6 years between ages 10 and 20, with parental origin explaining 36–53% of the response to growth hormone. Sagi et al. (2007) found a mean height gain per year in response to growth hormone treatment that was 28% greater in Xmat than Xpat females, but this difference was not statistically significant. Males are substantially and significantly more responsive to growth hormone treatment than females (Burman et al. 1997; Thangavel and Shapiro 2007).

### Lipid profiles and visceral fat

Turner syndrome females exhibit an atherogenic lipid profile (a distribution of serum fatty acid levels associated with high risk of atherosclerosis) and high levels of visceral fat compared to normal 46,XX females (Van et al. 2006a). Van et al. (2006b) reported significantly higher levels of visceral fat, and higher levels of total cholesterol, LDL cholesterol, and triglycerides, in Xmat than Xpat Turner females aged 27–31 years on average, and they note that this difference directly parallels the

difference between normal males and females. By contrast, in a population of Turner females with a mean age of 15 years, Sagi et al. (2007) found lower total and low-density lipoprotein levels in the Xmat than Xpat group.

Among normal middle-aged populations, males exhibit higher levels of visceral fat, and higher LDL levels, than do females (Freedman et al. 2004; Van et al. 2006a,b). However, such sex differences in LDL levels are absent or much less pronounced in children and adolescents (Freedman et al. 2000; Jolliffe and Janssen 2006).

Compared to 45,X/46,XX mosaics, levels of LDL, triglycerides, and body fat were higher in 45,X females by 10%, 26% and 18% respectively in the study of El-Mansouri et al. (2007), but these differences were not statistically significant. By contrast, levels of total cholesterol were significantly higher in 45,X than 45,X/46,XX mosaics, by 15% ( $P < 0.01$  in El-Mansouri et al. 2007). This population exhibited a mean age of 31, comparable with that of Van et al. (2006b).

### Sensorineural hearing loss

Turner syndrome females exhibit high rates of early-onset hearing loss, due to otitis media (middle-ear infections), auricular anomalies, and other causes (King et al. 2007), with symptoms notably more severe in cases with monosomy 45,X than in cases with mosaicism or structural X-chromosome defects (Barrenäs et al. 1999, 2000; Morimoto et al. 2006). Otitis media and aging-related hearing loss are also more common and severe in males than 46,XX females (see Barrenäs et al. 2000; Henry 2004), but early-onset hearing loss is very rare in such populations. In mouse models of hearing loss, females lose hearing earlier than males in a strain with early-onset hearing loss comparable in timing to that in Turner syndrome females, but males lose hearing earlier in strains with the late-onset, age-related loss that corresponds to the usual situation in humans (Henry 2004). These findings suggest that early-onset and late-onset hearing loss involve different mechanisms, that are mediated differently by sex.

Hamelin et al. (2006) found significantly less early-onset sensorineural hearing loss among Xmat (34% of patients) than Xpat (67%) females with Turner syndrome. This difference corresponds to the sex difference between normal male and female humans to the extent that mechanisms of early hearing loss are similar between females with Turner syndrome and mouse strains with early-onset hearing loss. El-Mansouri et al. (2007) reported a 51% incidence of impaired hearing in 45,X Turner females, compared to 26% in 45,X/46,XX mosaic females ( $P = 0.07$ ).

**Table 1.** The differences in phenotype between Turner syndrome females with a maternal versus paternal X chromosome are broadly consistent with the differences between normal males and females, and the differences between monosomic 45,X Turner females versus Turner females with other karyotypes. These parallel patterns may be caused by X-linked imprinting mediating the development of sexual dimorphism, by lower levels of mosaicism and higher rates of 45,X monosomy in Turner females with a maternal X chromosome, or by both processes (See text for details).

Trait	Xmat/Xpat difference	Sex difference	45,X/other difference	Comments
Response to growth hormone	Greater response in Xmat females (1) or no difference (2,3)	Males show larger response than females (4,5)	No data	Height in Turner syndrome affected by X-linked and autosomal genes (6)
Lipid profile and visceral fat	Higher cholesterol, LDL, and visceral fat in Xmat females, in middle age (7); lower total and LDL LDL cholesterol in Xmat females in adolescence (3)	Males have higher LDL and visceral fat than females across middle age; sexes similar in adolescence (7–11)	45,X females have higher cholesterol than 45,X/46,XX females (12)	In (7), females were age 27–31 on average; in (3), they were age 15 on average
Sensorineural hearing loss	Xmat females show lower levels of early-onset hearing loss (1)	Males show more overall hearing loss is rare (13,14); in mouse models, males have less early-onset hearing loss (14)	45,X females may have more hearing loss than 45,X/46,XX females ( $P = 0.07$ ) (12)	
Congenital heart defects	Xmat females exhibit more cardiac anomalies (15), or no difference (2,3,6)	Higher rates of aortic cardiac anomalies at birth in males (16,17)	Higher rates of aortic cardiac anomalies in 45,X than mosaic females (18,19)	Turner syndrome cardiac defects are found differentially in males (16–21).
Congenital kidney defects	Xmat females have higher rate of renal anomalies (3), or no difference (6)	Higher rates of renal anomalies at birth in males (16) or no difference (17)	No difference in rates of 'urinary track malformations' between 45,X and 45,X/46,XX females (12)	
Neuroanatomy	Larger cerebellum in Xmat than 46,XX females (22); larger superior temporal gyrus in Xmat than Xpat females (23); larger hippocampus and smaller caudate nuclei in Xmat than Xpat females (24); or no differences (25,26)	Males have larger cerebellum, larger left anterior superior temporal gyrus, and larger amygdala-hippocampus, but smaller caudate nuclei (27–29)	45X/46,XX females exhibit intermediacy between 45,X and 46,XX females for some neuroanatomical and neurological-function traits (30–32)	
Psychological traits	Xmat females show impaired social cognition, lower verbal skills, more attention, thought and aggression problems, higher rate of autism (33–35); Xmat females have better visual-spatial memory (35); twofold higher rate of ADHD in Xmat females but difference not significant (36)	Males exhibit poorer social and verbal skills than females, higher rates of autism and ADHD, and better visual-spatial skills (37–39)	Larger difference between high verbal and low performance skills in 45,X than 45,X/46,XX females (40)	

(1) Hamelin et al. 2006 (2) Tsezou et al. 1999 (3) Sagi et al. 2007 (4) Burman et al. 1997 (5) Thangavel and Shapiro 2007; (6) Bondy et al. 2007 (7) Van et al. 2006b (8) Van et al. 2006a (9) Freedman et al. 2000 (10) Freedman et al. 2000 (11) Jolliffe and Janssen 2006 (12) El-Mansouri et al. 2007 (13) Barrenäs et al. 2000 (14) Henry 2004 (15) Chu et al. 1994 (16) Lary and Paulozzi 2001 (17) Shaw et al. 2003 (18) Götzsche et al. 1994 (19) Prandstraller et al. 1999 (20) Geodakian and Sherman 1970 (21) Geodakian and Sherman 1971 (22) Brown et al. 2002 (23) Kesler et al. 2003 (24) Cutter et al. 2006 (25) Good et al. 2003 (26) Kesler et al. 2004 (27) Good et al. 2001 (28) Chen et al. 2007 (29) Wilke et al. 2007 (30) Murphy et al. 1997 (31) Murphy et al. 1993 (32) Murphy et al. 1994 (33) Skuse et al. 1997 (34) Skuse 1999 (35) Bishop et al. 2000 (36) Russell et al. 2006 (37) Geary 1998 (38) Baron-Cohen 2003 (39) Hermens et al. 2005 (40) Temple and Carney 1993.

### Congenital heart defects

Chu et al. (1994) reported a significantly higher incidence of cardiac anomalies in Turner females with an intact Xmat (34, 38% of 90) than an intact Xpat (4, 11% of 34, Fisher's exact test,  $P = 0.0026$ ), based on pooling of published data from four studies that individually yielded Fisher's exact values of 0.005 (Ross et al. 1991), 0.088 (Lorda-Sanchez et al. 1992), 0.15 (Chu et al. 1994) and 0.70 (Mathur et al. 1991). By contrast, three other studies have reported similar rates of cardiac anomalies between groups (Tsezou et al. 1999; Bondy et al. 2007; Sagi et al. 2007). Overall, using Fisher's combining test of probabilities, the difference between these two groups was not significant ( $\chi^2 = 20.3$ , 14 d.f.,  $P = 0.13$ ). Differences between studies may be due to the sensitivity of diagnostic methods (Bondy et al. 2007), and variation in the karyotypes present or the degrees of mosaicism, given substantially higher rates of cardiac anomalies in monosomic 45,X females than mosaic females (Götzsche et al. 1994; Prandstraller et al. 1999; El-Mansouri et al. 2007).

The cardiac defects most common in Turner syndrome include anomalies of the aorta, especially aortic coarctation and stenosis. These heart defects exhibit 20–50% higher rates in males (Geodakian and Sherman 1971; Lary and Paulozzi 2001; Shaw et al. 2003), and they have been considered as the most well-defined 'male' heart defects (Geodakian and Sherman 1970, 1971).

### Congenital kidney defects

Sagi et al. (2007) found a higher incidence of renal anomalies in Xmat Turner females (12/60) than in Xpat females (0/20;  $P = 0.03$ , Fisher's exact test). By contrast, a recent analysis with large samples sizes found no such difference (33/133 vs 12/50 respectively,  $P > 0.50$ ) (Bondy et al. 2007), as did earlier studies with small samples (reviewed in Sagi et al. 2007), including Chu et al. (1994). Fisher's combining test of the data from the seven studies to date showed a lack of overall significance ( $P > 0.50$ ). Congenital renal anomalies show a lack of sex bias (ratio 1:1) in one epidemiological study (Shaw et al. 2003), but a significant male bias (1.74:1) in another study (Lary and Paulozzi 2001).

### Neuroanatomy

For each of the three studies showing X-chromosome parent of origin effects on neuroanatomy in Turner syndrome (Brown et al. 2002; Kesler et al. 2003; Cutter et al. 2006), the observed parental-origin differences parallel the differences between males and 46,XX females (Good et al.

2001; Chen et al. 2007; Wilke et al. 2007), given the information available (Table 1).

### Psychological traits

Skuse et al. (1997) and Skuse (1999) reported that Xmat females exhibited higher levels of verbal, social, emotional and behavioural problems than Xpat females (Table 1); by contrast, Bishop et al. (2000) describe evidence that Xmat Turner females exhibit better visual-spatial memory, but worse verbal memory, than Xpat females, with females also better than males at this verbal-memory task. Russell et al. (2006) found that seven (35%) of 20 Xmat Turner females, and one (14%) of seven Xpat females were diagnosed with ADHD, but this difference was not statistically significant (Fisher's exact,  $P = 0.30$ ), nor was the difference significant between 45,X females (8, 30% of 27) and mosaic females (4, 17% of 23) (Fisher's exact test,  $P = 0.25$ ). Similarly, Sagi et al. (2007) reported that five of seven Xpat females, but only four of eleven Xmat females, had academic skills or degrees (Fisher's exact test,  $P = 0.17$ ).

The verbal versus visual-spatial differences between Turner syndrome females and 46,XX females contrast with the differences between males and 46,XX females, given that on average, males exhibit relatively better visual-spatial skills compared to verbal skills than do females (Geary 1998; Baron-Cohen 2003). However, as described by Skuse et al. (1997) and Skuse (1999), some of the neurocognitive differences between Xmat and Xpat Turner females, such as lower verbal, attentional, emotional and social skills in the Xmat genotype, notably parallel the differences between males and 46,XX females.

The effects of karyotype and mosaicism on cognitive functions in Turner syndrome have yet to be investigated in detail, but Temple and Carney (1993) reported that the differences between verbal IQ scores and performance IQ scores were larger in monosomic 45,X than in mosaic 45,X/46,XX Turner females, and Murphy et al. (1993, 1994, 1997) describes evidence from neuroimaging and cognitive studies for X-chromosome dosage effects on verbal versus visual-spatial/performance skills. Genetic evidence for such effects has been provided by Vawter et al. (2007), who found strong correlations of gene-expression levels with verbal skills, for several X-linked genes, in individuals with Klinefelter syndrome (usually XXY in males).

### Pleiotropic effects of growth

Of the traits in Table 1 that show evidence of differences between Xmat and Xpat Turner females, one trait, response to growth hormone, is a direct correlate of

growth parameters, and three additional traits, sensorineural hearing loss, lipid profile and body composition, and neuroanatomy, are also known to be growth-related. Thus, in females with Turner syndrome females, the extent of hearing loss is positively associated with reduced height (and lower IGF1 levels) (Barrenäs et al. 2000; Morimoto et al. 2006), and growth hormone treatment is associated with both reduced truncal (visceral) obesity (Gravholt et al. 2005) and increased levels of grey matter in the temporal, parietal and occipital lobes of the brain (Cutter et al. 2006). Taller females with Turner syndrome also bear a reduced number of anatomical stigmata (El-Mansouri et al. 2007), but there is no apparent effect of GH treatment or height on cognitive function in Turner syndrome (Ross 2005; Messina et al. 2007). Barrenäs et al. (2000) describe evidence that growth-related phenotypes in Turner syndrome (and other aneuploidies) are mediated by effects of aneuploidy on rates of cell turnover, which in Turner syndrome differentially modulate growth of SHOX-regulated mesodermal tissues with the shortest cell cycle time and highest cell cycle rate. This hypothesis of aneuploidy effects on cell cycle times is also supported by evidence for altered temporal control of cell replication in Turner syndrome (Reish et al. 2002), and changes in the proportions of 45,X vs 46,XX cell lines over time *in vivo* (Nielsen 1976; Held et al. 1992; Devi et al. 1998).

Despite patterns in covariation of clinical phenotype with height in Turner syndrome, Turner females clearly do not differ in height by parental origin of the X chromosome (Mathur et al. 1991; Bondy et al. 2007; Kochi et al. 2007; Sagi et al. 2007). Instead, the height of Turner syndrome females shows a strong, positive, highly significant correlation with their mother's height (regardless of parental origin of the X), but the association with father's height is smaller or nonexistent (Salerno and Job 1987; Chu et al. 1994; Tsezou et al. 1999; Hamelin et al. 2006; Bondy et al. 2007; Kochi et al. 2007). The simplest explanation for this remarkable, well-replicated finding is that haploinsufficiency of some X-linked, noninactivated gene or genes (such as SHOX) results in altered transactivation of one or more autosomal imprinted genes that regulate growth. The imprinted gene DLK1 represents a notable functional and positional candidate for such effects due to its roles in regulating growth, adiposity, and bone development (Abdallah et al. 2004, 2007; Ansell et al. 2007), its location at 14q32.2 where apparent imprinting effects on human height have been described (Mukhopadhyay and Weeks 2003), and the phenotypic effects of reduced or absent DLK1 expression, which include low birth weight, short stature, high palate, micrognathia (small teeth), small hands, hypotonia (weak muscle tone), scoliosis, recurrent otitis media, high cholesterol and

obesity (Kotzot 2004; Temple et al. 2007), all of which are relatively common in Turner syndrome. Comparable interactions between X-linked genes and autosomal imprinted genes affecting growth have been described in mice (Vrana et al. 2000; Loschiavo et al. 2007), and Pan et al. (2007) describe sex-specific X-chromosome effects on height and triglyceride levels that are consistent with an important role for sex linkage in phenotypic variation for these traits. Taken together, these findings suggest that growth-related phenotypes in Turner syndrome are mediated in part by one or more autosomal imprinted genes, as well as by X-linked genes.

### Alternative hypotheses for differences between Xmat and Xpat Turner females

Most studies of phenotypic differences between Xmat and Xpat Turner females have interpreted their results in terms of hypothesized effects of one or more X-linked, imprinted genes (e.g. Skuse et al. 1997; Sagi et al. 2007). However there is, as yet, no conclusive evidence for the presence of imprinted genes on the human X-chromosome, despite the discovery of several such genes in mice (Davies et al. 2005; Raefski and O'Neill 2005) and the inferred presence of X-linked imprinted genes in humans from mapping of sex-differential effects on prenatal lethality (Naumova et al. 1998; Green and Keverne 2000).

An alternative, though nonexclusive, hypothesis for differences between Xmat and Xpat Turner females is confounding of parental origin of the X chromosome with the form of the karyotype and the degree of mosaicism in Turner syndrome, such that Xmat and Xpat females tend to exhibit a different karyotype, a differing degree of mosaicism, or both (Box 1). By this hypothesis, Turner females with the Xmat intact, which tend to exhibit a more male-typical Turner-syndrome phenotype for some traits (Table 1), are presumed to have developed under a lower degree of cryptic or documented mosaicism (which leads to a relatively female-typical phenotype) (e.g. Henn and Zang 1997; Haerkamp et al. 1999; Hanson et al. 2001; El-Mansouri et al. 2007), or under the influence of specific karyotypes that produce a more female-typical phenotype, such as karyotypes that lack Y-chromosome material. Data on the frequency of different karyotypes and mosaicism in Turner females with an intact Xmat versus Xpat are now available from four studies, which allows such alternative hypotheses to be evaluated.

First, in Bondy et al. (2007) (Table 1), 61 (46%) of 133 Xmat females were not pure 45,X karyotypes or mosaics (for diverse karyotypes including 46,XX), compared to 33 (66%) of 50 Xpat females (Fisher's exact test,  $P = 0.011$ ). Considering monosomic and mosaic females only, 44 (38%) of 116 Xmat females were mosaics,

**Box 1.** Alternative, nonexclusive hypotheses for the presence of differences between Xmat and Xpat females that parallel the differences between males and females.


**(1) Genomic imprinting.** For X-linked imprinted genes, gene dosages are expected to differ between males and females, with the nature of the difference depending upon the direction of imprinting and whether or not the gene is X-inactivated (Skuse 1999). Males exclusively bear the maternally inherited X, so Turner females with an intact maternally inherited X are expected to exhibit relatively male-typical phenotypes for traits mediated by X-linked imprinted genes. By contrast, females bear one paternally inherited X, and one maternally inherited X.

**(2) Mosaicism and karyotype.** For X-linked genes that are not inactivated, males express one copy, and 46,XX females express two copies. To the extent that human sexual dimorphism is mediated by dosages of such X-linked genes, Turner females are expected to exhibit some degree of male-typical traits. Turner females with a monosomic 45,X karyotype are thus expected to bear traits relatively more typical of males than Turner females with other karyotypes. To the extent that the 45,X karyotype differentially involves the maternally inherited X, due to the nature of the cytogenetic mechanisms whereby it becomes the sole or primary cell line (e.g. Fig. 1), Turner females with an intact maternal X are expected to be more likely to exhibit relatively male-typical traits, compared to Turner females with an intact paternal X. Females with a maternally inherited X are also more likely to bear Y-chromosomal material, but there is no evidence that Y-linked genes mediate the phenotype in Turner syndrome except in some cases of 45,X/46,XY mosaicism, which is rare.

compared to 28 (62%) of 45 Xpat females (Fisher's exact test,  $P = 0.0046$ ). A lower level of mosaicism in Xmat than Xpat Turner females is also suggested by some hypotheses for the generation of Turner syndrome chromosomal anomalies, which posit that deletions (and some other alterations) of all or part of the X chromosome are relatively frequent in the rapidly replicating paternal germ line (Kelly et al. 1992; Jacobs et al. 1997; Uematsu et al. 2002), that mosaicism or a karyotype other than 45,X early in development may be a prerequisite for viable embryonic development (Hecht and Macfarlane 1969; Hook and Warburton 1983), and that abnormal X chromosomes may be differentially lost in development, such that all or most females karyotyped after birth as 45,X are either cryptic mosaics (with a second cell line present but undetected), or exhibited a mosaic karyotype, or a partial second X, earlier in development (Held et al. 1992; Amiel et al. 1996) (Fig. 1). For example, Kelly et al. (1992) provide experimental evidence that mosaicism may be present in fetuses with Turner syndrome, but be lost prior to birth, resulting in 45,X. Mosaicism involving 45,X/46,XX can also be generated via postzygotic nondisjunction, a mechanism that can also explain the presence of mosaicism of the form 45,X/47,XXX in some Turner females.

Second, Sagi et al. (2007) found that 83% (55 of 66) of monosomic 45,X females were Xmat rather than Xpat, whereas 36% (five of 14) females with the isodicentric karyotype 46,XiXq bore Xmat as their intact X chromosome (Fisher's exact test,  $P = 0.0007$ ). These authors did not find significant differences in any phenotypic trait between females with monosomic 45,X versus isodicentric 46,XiXq karyotypes, but evidence for differences between these two karyotypes has been reported in other studies for IQ (Messina et al. 2007) and height (Cohen et al. 1995).

Third, the study population of Hamelin et al. (2006) (Fig. 1) included 7 (20%) of 35 Xmat females that were



**Figure 1** One scenario for the generation and development of monosomy 45,X with the intact chromosome maternally inherited, in Turner syndrome. This series of events is compatible with data showing a high incidence of 45,X in aborted fetuses, which apparently exhibited this karyotype at fertilization (Hook and Warburton 1983), and with data showing changes in karyotype over time, with differential loss of abnormal X chromosomes in some cases (Held et al. 1992; Kelly et al. 1992; Amiel et al. 1996). Turner females may also be born with a mosaic karyotype, depending upon the rate of loss of the abnormal X chromosome. Deletions of Xp, and some other cytogenetic changes involving the X, may be relatively more common in the rapidly dividing paternal germ line (Uematsu et al. 2002). The 45,X karyotype is much more common in females with a maternally inherited X than with a paternally inherited X (Uematsu et al. 2002; Bondy et al. 2007; Sagi et al. 2007).

either not pure 45,X or mosaics (with 46,XX mosaics not included), and 8 (42%) of 19 Xpat females mosaic or otherwise not 45,X (Fisher's exact test,  $P = 0.080$ ). However, this marginally nonsignificant difference was due to a significantly higher incidence of the nonmosaic 46,XiXq karyotype in Xpat females; when this karyotype category is excluded, the frequency of mosaics is essentially the same in both Xmat (15%) and Xpat (18%) females for their sample.

Fourth, Uematsu et al. (2002), Table III) assembled data from 21 earlier studies, and found that most (459, 75% of 614) pure 45,X females bore an X of maternal origin, while the 46,XiXq karyotype was about equally distributed between Xmat ( $n = 60$ ) and Xpat ( $n = 71$ ) females. Thus, considering these two karyotypes, pure 45,X karyotypes were significantly more frequent in Xmat females (88%) than in Xpat females (69%) (Fisher's exact test,  $P < 0.001$ ). These results are consistent with the data from Sagi et al. (2007) described above, with data from Hamelin et al. (2006) who found a higher incidence of isodicentric chromosomes in Xpat females, and with the results of Bondy et al. (2007) given that in their sample, the karyotype 45,X/46,XiXq comprised 13% of the Xmat females, but 26% of the Xpat females.

Uematsu et al. (2002) also showed that 19 of 20 females with Y-chromosome material bore an intact maternal X, which is consistent with simple expectations from Mendelian inheritance. Overall, Y-chromosomal material has been reported in about 10–20% of Turner syndrome cases (Gravholt et al. 2000; Hanson et al. 2001; Alvarez-Nava et al. 2003). In 45,X/46,XY mosaics, the phenotype can vary from Turner-like female, to intermediate in sexual development, to male, depending upon the presence or absence of the male-determining SRY gene and the degree of mosaicism (Robinson et al. 1999; Telvi et al. 1999). However, in Turner syndrome cases, the presence and form of Y material is apparently not associated with phenotype (Telvi et al. 1999).

Taken together, these data indicate that parental origin of the X chromosome can be confounded with karyotype in three ways: (i) females with an isodicentric karyotype (46,XiXq or 45,X/46,XiXq) are relatively more likely, or similarly likely, to bear an intact paternal than maternal X chromosome, (ii) Xmat females appear less likely than Xpat females to exhibit mosaicism when karyotyped and (iii) Xmat females are more likely than Xpat females to bear Y-chromosomal material. Based on available evidence from karyotype-phenotype correlations, the second difference may parsimoniously account, at least in part, for parallel patterns in phenotypic variation between Xmat versus Xpat Turner females, compared to males versus 46,XX females. However, it is important to note that these parallel patterns are by no means consistently

supported for each phenotype examined, and that more data are needed on mosaicism in relation to parental origin of the X for robust evaluation of this hypothesis.

Most generally, separating the confounded effects of parental origin of the X, karyotype and mosaicism requires fine-scale genotype–phenotype correlations with larger samples than have been used thus far in most studies. Similar considerations should also apply to Klinefelter syndrome, for which phenotypic differences between XmatXmatY and XmatXpatY males have been described (Stemkens et al. 2006; Wikström et al. 2006). Thus, about 8–20% of Klinefelter patients are 46,XY/47,XXX mosaics with relatively moderate phenotypes (Ratcliffe et al. 1986; Bojesen et al. 2003; Abdelmoula et al. 2004), and only XmatXmatY males may develop as a result of postzygotic errors in mitosis (Thomas and Hassold 2003). Future studies of the causes of phenotypic variation among and between individuals with different sex chromosome aneuploidies might usefully focus on traits, such as fingerprint ridge counts (Penrose 1968) and enamel and dentin thickness (Alvesalo 1997; Lähdesmäki and Alvesalo 2006), for which male–female differences appear to mirror differences between Turner females and Klinefelter males, and for which effects of parental origin of the X have yet to be investigated.

## Conclusions

The development of human sexual dimorphism is mediated by four main causes: (i) hormonal differences that follow from activation of the SRY male-determining gene, (ii) other effects of genes on the Y, (iii) dosage effects of the 15–20% of X-linked genes that are not inactivated (Carrel and Willard 2005), and (iv) hypothetically, by genes that are X-linked and imprinted (Skuse 1999, 2000, 2005; Arnold et al. 2004; Davies and Wilkinson 2006; Xu and Disteche 2006; Blecher and Erickson 2007). I have shown in this paper that the differences between Turner syndrome females with an intact maternally inherited versus paternally inherited X chromosome broadly parallel the differences between females who are monosomic 45,X versus other karyotypes, and the differences between normal XY males and XX females. A simple explanation for these patterns, which is supported by data showing relatively severe and relatively male-typical phenotypes in pure X monosomy, is that in Turner syndrome the maternally inherited X is more-frequently monosomic than the paternally inherited X. To the extent that noninactivated X-linked genes, differentially expressed between XY males and XX females, explain variation in Turner syndrome phenotypes, they are also implicated in the development and evolution of human sex differences;

similarly, to the extent that X-linked imprinted genes exist and influence Turner syndrome phenotypes, the patterns described here would strongly implicate such genes as an important mechanism of sexual differentiation. Determining the roles of X-linked imprinting, karyotype, and mosaicism in Turner syndrome may thus help in deciphering not just the genetic aetiology of this condition, but also the genetic and epigenetic basis of human sexual dimorphism (Davies and Wilkinson 2006; Skuse 2006).

### Acknowledgements

I am grateful to T. Day and four anonymous reviewers for helpful comments, and I thank the Natural Sciences and Engineering Research Council of Canada and the Killam Trust for support.

### Literature cited

Abdallah, B. M., C. H. Jensen, G. Gutierrez, R. G. Q. Leslie, T. G. Jensen, and M. Kassem. 2004. Regulation of human skeletal stem cells differentiation by Dlk1/Pref-1. *Journal of Bone and Mineral Research* **19**:841–852.

Abdallah, B. M., P. Boissy, Q. Tan, J. Dahlgaard, G. A. Traustadottir, K. Kupisiewicz, J. Laborda *et al.* 2007. dlk1/FA1 regulates the function of human bone marrow mesenchymal stem cells by modulating gene expression of pro-inflammatory cytokines and immune response-related factors. *Journal of Biological Chemistry* **282**:7339–7351.

Abdelmoula, N. B., A. Amouri, M. Portnoi, A. Saad, T. Boudawara, M. N. Mhiri, A. Bahloul, and T. Rebai. 2004. Cytogenetics and fluorescence in situ hybridization assessment of sex-chromosome mosaicism in Klinefelter's syndrome. *Annales De Génétique* **47**:163–175.

Alvarez-Nava, F., M. Soto, M. A. Sánchez, E. Fernández, and R. Lanes. 2003. Molecular analysis in Turner syndrome. *Journal of Pediatrics* **142**:336–340.

Alvesalo, L. 1997. Sex chromosomes and human growth. A dental approach. *Human Genetics* **101**:1–5.

Amiel, A., D. Kidron, I. Kedar, E. Gaber, O. Reish, and M. D. Feigin. 1996. Are all phenotypically-normal Turner syndrome fetuses mosaics? *Prenatal Diagnosis* **16**:791–795.

Ansell, P. J., Y. Zhou, B. Schjeide, A. Kerner, J. Zhao, X. Zhang, and A. Klibanski. 2007. Regulation of growth hormone expression by Delta-like protein 1 (Dlk1). *Molecular and Cellular Endocrinology* **271**:55–63.

Arnold, A. P., J. Xu, W. Grisham, X. Chen, Y. Kim, and Y. Itoh. 2004. Minireview: sex chromosomes and brain sexual differentiation. *Endocrinology* **145**:1057–1062.

Baron-Cohen, S. 2003. *The Essential Difference: The Truth About the Male and Female Brain*. Basic Books, New York.

Barrenäs, M. L., O. Nylén, and C. Hanson. 1999. The influence of karyotype on the auricle, otitis media and hearing in Turner syndrome. *Hearing Research* **138**:163–170.

Barrenäs, M., K. Landin-Wilhelmsen, and C. Hanson. 2000. Ear and hearing in relation to genotype and growth in Turner syndrome. *Hearing Research* **144**:21–28.

Berg, J. S., N. Brunetti-Pierri, S. U. Peters, S. L. Kang, C. Fong, J. Salamone, D. Freedenberg *et al.* 2007. Speech delay and autism spectrum behaviors are frequently associated with duplication of the 7q11.23 Williams-Beuren syndrome region. *Genetics in Medicine* **9**:427–441.

Bishop, D. V., E. Canning, K. Elgar, E. Morris, P. A. Jacobs, and D. H. Skuse. 2000. Distinctive patterns of memory function in subgroups of females with Turner syndrome: evidence for imprinted loci on the X-chromosome affecting neurodevelopment. *Neuropsychologia* **38**:712–721.

Blecher, S. R., and R. P. Erickson. 2007. Genetics of sexual development: a new paradigm. *American Journal of Medical Genetics. Part A* **143**:3054–3068.

Bojesen, A., S. Juul, and C. H. Gravholt. 2003. Prenatal and postnatal prevalence of Klinefelter syndrome: a national registry study. *Journal of Clinical Endocrinology and Metabolism* **88**:622–626.

Bondy, C. A. 2006. Turner's syndrome and X chromosome-based differences in disease susceptibility. *Gender Medicine* **3**:18–30.

Bondy, C. A., L. A. Matura, N. Wooten, J. Troendle, A. R. Zinn, and V. K. Bakalov. 2007. The physical phenotype of girls and women with Turner syndrome is not X-imprinted. *Human Genetics* **121**:469–474.

Brock, J. (2007) Language abilities in Williams syndrome: a critical review. *Development and Psychopathology* **19**:97–127.

Brown, W. E., S. R. Kesler, S. Eliez, I. S. Warsofsky, M. Haberrecht, A. Patwardhan, J. L. Ross *et al.* 2002. Brain development in Turner syndrome: a magnetic resonance imaging study. *Psychiatry Research* **116**:187–196.

Burman, P., A. G. Johansson, A. Siegbahn, B. Vessby, and F. A. Karlsson. 1997. Growth hormone (GH)-deficient men are more responsive to GH replacement therapy than women. *Journal of Clinical Endocrinology and Metabolism* **82**:550–555.

Carrel, L., and H. F. Willard. 2005. X-inactivation profile reveals extensive variability in X-linked gene expression in females. *Nature* **434**:400–404.

Chen, X., P. S. Sachdev, W. Wen, and K. J. Anstey. 2007. Sex differences in regional gray matter in healthy individuals aged 44–48 years: a voxel-based morphometric study. *NeuroImage* **36**:691–699.

Chu, C. E., M. D. Donaldson, C. J. Kelnar, P. J. Smail, S. A. Greene, W. F. Paterson, and J. M. Connor. 1994. Possible role of imprinting in the Turner phenotype. *Journal of Medical Genetics* **31**:840–842.

Cohen, A., R. Kauli, A. Pertzelan, A. Lavagetto, Y. Roitmano, C. Romano, and Z. Laron. 1995. Final height of girls with

Turner's syndrome: correlation with karyotype and parental height. *Acta Paediatrica* **84**:550–554.

Crespi, B., and C. Badcock. 2008. Psychosis and autism as diametrical disorders of the social brain. *Behavioral and Brain Sciences* (in press).

Cutter, W. J., E. M. Daly, D. M. Robertson, X. A. Chitnis, T. A. van Amelsvoort, A. Simmons, V. W. Ng *et al.* 2006. Influence of X chromosome and hormones on human brain development: a magnetic resonance imaging and proton magnetic resonance spectroscopy study of Turner syndrome. *Biological Psychiatry* **59**:273–283.

Davies, W., and L. S. Wilkinson. 2006. It is not all hormones: alternative explanations for sexual differentiation of the brain. *Brain Research* **1126**:36–45.

Davies, W., A. Isles, R. Smith, D. Karunadasa, D. Burrmann, T. Humby, O. Ojarikre *et al.* 2005. Xlr3b is a new imprinted candidate for X-linked parent-of-origin effects on cognitive function in mice. *Nature Genetics* **37**:625–629.

Davies, W., A. R. Isles, P. S. Burgoine, and L. S. Wilkinson. 2006. X-linked imprinting: effects on brain and behaviour. *BioEssays* **28**:35–44.

Devi, A. S., D. A. Metzger, A. A. Luciano, and P. A. Benn. 1998. 45,X/46,XX mosaicism in patients with idiopathic premature ovarian failure. *Fertility and Sterility* **70**:89–93.

El-Mansouri, M., M. Barrenäs, I. Bryman, C. Hanson, C. Larsson, L. Wilhelmsen, and K. Landin-Wilhelmsen. 2007. Chromosomal mosaicism mitigates stigmata and cardiovascular risk factors in Turner syndrome. *Clinical Endocrinology* **66**:744–751.

Feinstein, C. and S. Singh. 2007. Social phenotypes in neurogenetic syndromes. *Child and Adolescent Psychiatry Clinics of North America* **16**:631–647.

Fernández-García, R., S. García-Doval, S. Costoya, and E. Pásaro. 2000. Analysis of sex chromosome aneuploidy in 41 patients with Turner syndrome: a study of 'hidden' mosaicism. *Clinical Genetics* **58**:201–208.

Freedman, D. S., B. A. Bowman, J. D. Otvos, S. R. Srinivasan, and G. S. Berenson. 2000. Levels and correlates of LDL and VLDL particle sizes among children: the Bogalusa heart study. *Atherosclerosis* **152**:441–449.

Freedman, D. S., J. D. Otvos, E. J. Jeyarajah, I. Shalaurova, L. A. Cupples, H. Parise, R. B. D'Agostino, P. W. Wilson and E. J. Schaefer. 2004. Sex and age differences in lipoprotein subclasses measured by nuclear magnetic resonance spectroscopy: the Framington Study. *Clinical Chemistry* **50**:1189–1200.

Geary, D. C. 1998. *Male, Female: The Evolution of Human Sex Differences*. American Psychological Association, Washington, DC.

Geodakian, V. A., and A. L. Sherman. 1970. Congenital heart disease and sex. *Eksperimental'Naia Khirurgiia i Anestesiologiya* **15**:18–23.

Geodakian, V. A., and A. L. Sherman. 1971. Relation of congenital anomalies to sex. *Zhurnal Obshchei Biologii* **32**:417–424.

Good, C. D., I. Johnsrude, J. Ashburner, R. N. Henson, K. J. Friston, and R. S. Frackowiak. 2001. Cerebral asymmetry and the effects of sex and handedness on brain structure: a voxel-based morphometric analysis of 465 normal adult human brains. *NeuroImage* **14**:685–700.

Good, C. D., K. Lawrence, N. S. Thomas, C. J. Price, J. Ashburner, K. J. Friston, R. S. J. Frackowiak *et al.* 2003. Dosage-sensitive X-linked locus influences the development of amygdala and orbitofrontal cortex, and fear recognition in humans. *Brain* **126**:2431–2446.

Götzsche, C. O., B. Krag-Olsen, J. Nielsen, K. E. Sørensen, and B. O. Kristensen. 1994. Prevalence of cardiovascular malformations and association with karyotypes in Turner's syndrome. *Archives of Disease in Childhood* **71**:433–436.

Gravholt, C. H., J. Fedder, R. W. Naeraa, and J. Müller. 2000. Occurrence of gonadoblastoma in females with Turner syndrome and Y chromosome material: a population study. *The Journal of Clinical Endocrinology and Metabolism* **85**:3199–3202.

Gravholt, C. H., B. E. Hjerrild, R. W. Naeraa, F. Engbaek, L. Mosekilde, and J. S. Christiansen. 2005. Effect of growth hormone and 17beta-oestradiol treatment on metabolism and body composition in girls with Turner syndrome. *Clinical Endocrinology* **62**:616–622.

Green, R., and E. B. Keverne. 2000. The disparate maternal aunt-uncle ratio in male transsexuals: an explanation invoking genomic imprinting. *Journal of Theoretical Biology* **202**:55–63.

Haig, D., and R. Wharton. 2003. Prader-Willi syndrome and the evolution of human childhood. *American Journal of Human Biology* **15**:320–329.

Hamelin, C. E., G. Anglin, C. A. Quigley, and C. L. Deal. 2006. Genomic imprinting in Turner syndrome: effects on response to growth hormone and on risk of sensorineural hearing loss. *Journal of Clinical Endocrinology and Metabolism* **91**:3002–3010.

Hanson, L., I. Bryman, M. L. Barrenäs, P. O. Janson, J. Wahlström, K. Albertsson-Wikland, and C. Hanson. 2001. Genetic analysis of mosaicism in 53 women with Turner syndrome. *Hereditas* **134**:153–159.

Haverkamp, F., J. Wölflé, K. Zerres, O. Butenandt, P. Amendt, B. P. Hauffa, E. Weimann *et al.* 1999. Growth retardation in Turner syndrome: aneuploidy, rather than specific gene loss, may explain growth failure. *Journal of Clinical Endocrinology and Metabolism* **84**:4578–4582.

Hecht, F., and J. P. Macfarlane. 1969. Mosaicism in Turner's syndrome reflects the lethality of XO. *Lancet* **2**:1197–1198.

Held, K. R., S. Kerber, E. Kaminsky, S. Singh, P. Goetz, E. Seemanova, and H. W. Goedde. 1992. Mosaicism in 45,X Turner syndrome: does survival in early pregnancy depend on the presence of two sex chromosomes? *Human Genetics* **88**:288–294.

Henn, W., and K. D. Zang. 1997. Mosaicism in Turner's syndrome. *Nature* **390**:569–569.

Henry, K. R.. 2004. Males lose hearing earlier in mouse models of late-onset age-related hearing loss; females lose hearing earlier in mouse models of early-onset hearing loss. *Hearing Research* **190**:141–148.

Hermens, D. F., M. R. Kohn, S. D. Clarke, E. Gordon, and L. M. Williams. 2005. Sex differences in adolescent ADHD: findings from concurrent EEG and EDA. *Clinical Neurophysiology* **116**:1455–1463.

Hook, E. B., and D. Warburton. 1983. The distribution of chromosomal genotypes associated with Turner's syndrome: livebirth prevalence rates and evidence for diminished fetal mortality and severity in genotypes associated with structural X abnormalities or mosaicism. *Human Genetics* **64**:24–27.

Iwasa, Y., and A. Pomiankowski. 1999. Sex specific X chromosome expression caused by genomic imprinting. *Journal of Theoretical Biology* **197**:487–495.

Jacobs, P., P. Dalton, R. James, K. Mosse, M. Power, D. Robinson, and D. Skuse. 1997. Turner syndrome: a cytogenetic and molecular study. *Annals of Human Genetics* **61**:471–483.

Jolliffe, C. J., and I. Janssen. 2006. Distribution of lipoproteins by age and gender in adolescents. *Circulation* **114**:1056–1062.

Kelly, T. E., J. E. Ferguson, and W. Golden. 1992. Survival of fetuses with 45,X: an instructive case and an hypothesis. *American Journal of Medical Genetics* **42**:825–826.

Kesler, S. R.. 2007. Turner syndrome. *Child and Adolescent Psychiatric Clinics of North America* **16**:709–722.

Kesler, S. R., C. M. Blasey, W. E. Brown, J. Yankowitz, S. M. Zeng, B. G. Bender, and A. L. Reiss. 2003. Effects of X-monosomy and X-linked imprinting on superior temporal gyrus morphology in Turner syndrome. *Biological Psychiatry* **54**:636–646.

Kesler, S. R., A. Garrett, B. Bender, J. Yankowitz, S. M. Zeng, and A. L. Reiss. 2004. Amygdala and hippocampal volumes in Turner syndrome: a high-resolution MRI study of X-monosomy. *Neuropsychologia* **42**:1971–1978.

King, K. A., T. Makishima, C. K. Zalewski, V. K. Bakalov, A. J. Griffith, C. A. Bondy, and C. C. Brewer. 2007. Analysis of auditory phenotype and karyotype in 200 females with Turner syndrome. *Ear and Hearing* **28**:831–841.

Kochi, C., C. A. Longui, S. H. V. Lemos-Marini, G. Guerra-Junior, M. B. Melo, L. E. P. Calliari, and O. Monte. 2007. The influence of parental origin of X chromosome genes on the stature of patients with 45 X Turner syndrome. *Genetics and Molecular Research* **6**:1–7.

Kotzot, D. 2004. Maternal uniparental disomy 14 dissection of the phenotype with respect to rare autosomal recessively inherited traits, trisomy mosaicism, and genomic imprinting. *Annales De Génétique* **47**:251–260.

Lähdesmäki, R., and L. Alvesalo. 2006. Root growth in the permanent teeth of 45,X/46,XX females. *European Journal of Orthodontics* **28**:339–344.

Lary, J. M., and L. J. Paulozzi. 2001. Sex differences in the prevalence of human birth defects: a population-based study. *Teratology* **64**:237–251.

Leppig, K. A., and C. M. Disteche. 2001. Ring X and other structural X chromosome abnormalities: X inactivation and phenotype. *Seminars in Reproductive Medicine* **19**:147–157.

Lorda-Sánchez, I., F. Binkert, M. Maechler, and A. Schinzel. 1992. Molecular study of 45,X conceptuses: correlation with clinical findings. *American Journal of Medical Genetics* **42**:487–490.

Loschiavo, M., Q. K. Nguyen, A. R. Duselis, and P. B. Vrana. 2007. Mapping and identification of candidate loci responsible for *Peromyscus* hybrid overgrowth. *Mammalian Genome* **18**:75–85.

Lynn, P. M. Y., and W. Davies. 2007. The 39,XO mouse as a model for the neurobiology of Turner syndrome and sex-biased neuropsychiatric disorders. *Behavioural Brain Research* **179**:173–182.

Mathur, A., L. Stekol, D. Schatz, N. K. McLaren, M. L. Scott, and B. Lippe. 1991. The parental origin of the single X chromosome in Turner syndrome: lack of correlation with parental age or clinical phenotype. *American Journal of Human Genetics* **48**:682–686.

Messina, M. F., G. Zirilli, R. Civa, I. Rulli, G. Salzano, T. Aversa, and M. Valenzise. 2007. Neurocognitive profile in Turner's syndrome is not affected by growth impairment. *Journal of Pediatric Endocrinology & Metabolism* **20**:677–684.

Meyer-Lindenberg, A., C. B. Mervis, and K. F. Berman. 2006. Neural mechanisms in Williams syndrome: a unique window to genetic influences on cognition and behaviour. *Nature Reviews Neuroscience* **7**:380–393.

Migeon, B. R., P. Jeppesen, B. S. Torchia, S. Fu, M. A. Dunn, J. Axelman, B. J. Schmeckpeper *et al.* 1996. Lack of X inactivation associated with maternal X isodisomy: evidence for a counting mechanism prior to X inactivation during human embryogenesis. *American Journal of Human Genetics* **58**:161–170.

Mills, W., and T. Moore. 2006. Evolution of mammalian X chromosome-linked imprinting. *Cytogenetic and Genome Research* **113**:336–344.

Money, J. 1993. Specific neuro-cognitive impairments associated with Turner (45,X) and Klinefelter (47,XXY) syndromes: a review. *Social Biology* **40**:147–151.

Morimoto, N., T. Tanaka, H. Taiji, R. Horikawa, Y. Naiki, Y. Morimoto, and N. Kawashiro. 2006. Hearing loss in Turner syndrome. *Journal of Pediatrics* **149**:697–701.

Mukhopadhyay, N., and D. E. Weeks. 2003. Linkage analysis of adult height with parent-of-origin effects in the Framingham Heart Study. *BMC Genetics* **4**(Suppl. 1):S76.

Murphy, D. G., C. DeCarli, E. Daly, J. V. Haxby, G. Allen, B. J. White, A. R. McIntosh *et al.* 1993. X-chromosome effects on female brain: a magnetic resonance imaging study of Turner's syndrome. *Lancet* **342**:1197–1200.

Murphy, D. G., G. Allen, J. V. Haxby, K. A. Largay, E. Daly, B. J. White, C. M. Powell, and M. B. Schapiro. 1994. The effects of sex steroids, and the X chromosome, on female brain function: a study of the neuropsychology of adult Turner syndrome. *Neuropsychologia* **32**:1309–1323.

Murphy, D. G., M. J. Mentis, P. Pietrini, C. Grady, E. Daly, J. V. Haxby, M. De La Granja *et al.* 1997. A PET study of Turner's syndrome: effects of sex steroids and the X chromosome on brain. *Biological Psychiatry* **41**:285–298.

Naumova, A. K., L. Olien, L. M. Bird, M. Smith, A. E. Verner, M. Leppert, K. Morgan, and C. Sapienza. 1998. Genetic mapping of X-linked loci involved in skewing of X chromosome inactivation in the human. *European Journal of Human Genetics* **6**:552–562.

Nielsen, J. 1976. Cell selection in vivo in normal/aneuploid chromosome abnormalities. *Human Genetics* **32**:203–206.

Nielsen, R., C. Bustamante, A. G. Clark, S. Glanowski, T. B. Sackton, M. J. Hubisz, A. Fledel-Alon *et al.* 2005. A scan for positively selected genes in the genomes of humans and chimpanzees. *PLoS Biology* **3**:e170.

Oliver, C., K. Horsler, K. Berg, G. Bellamy, K. Dick, and E. Griffiths. 2007. Genomic imprinting and the expression of affect in Angelman syndrome: what's in the smile? *Journal of Child Psychology and Psychiatry, and Allied Disciplines* **48**:571–579.

Pan, L., C. Ober, and M. Abney. 2007. Heritability estimation of sex-specific effects on human quantitative traits. *Genetic Epidemiology* **31**:338–347.

Penrose, L. S.. 1968. Medical significance of finger-prints and related phenomena. *British Medical Journal* **2**:321–325.

Prandstraller, D., L. Mazzanti, F. M. Picchio, C. Magnani, R. Bergamaschi, A. Perri, E. Tsingos, and E. Cacciari. 1999. Turner's syndrome: cardiologic profile according to the different chromosomal patterns and long-term clinical follow-up of 136 nonpresselected patients. *Pediatric Cardiology* **20**:108–112.

Raefski, A. S., and M. J. O'Neill. 2005. Identification of a cluster of X-linked imprinted genes in mice. *Nature Genetics* **37**:620–624.

Ratcliffe, S. G., L. Murray, and P. Teague. 1986. Edinburgh study of growth and development of children with sex chromosome abnormalities. III. *Birth Defects Original Article Series* **22**:73–118.

Reish, O., R. Gal, E. Gaber, C. Sher, T. Bistritzer, and A. Amiel. 2002. Asynchronous replication of biallelically expressed loci: a new phenomenon in Turner syndrome. *Genetics in Medicine* **4**:439–443.

Robinson, D. O., P. Dalton, P. A. Jacobs, K. Mosse, M. M. Power, D. H. Skuse, and J. A. Crolla. 1999. A molecular and FISH analysis of structurally abnormal Y chromosomes in patients with Turner syndrome. *Journal of Medical Genetics* **36**:279–284.

Ross, J. L. 2005. Effects of growth hormone on cognitive function. *Hormone Research* **64**(Suppl. 3):89–94.

Ross, J. L., J. G. Hall, and E. G. Pfendner. 1991. The contribution of imprinting to the phenotype in Turner syndrome. *American Journal of Human Genetics* **49**:s19.

Russell, H. F., D. Wallis, M. M. Mazzocco, T. Moshang, E. Zackai, A. R. Zinn, J. L. Ross, and M. Muenke. 2006. Increased prevalence of ADHD in Turner syndrome with no evidence of imprinting effects. *Journal of Pediatric Psychology* **31**:945–955.

Sagi, L., N. Zuckerman-Levin, A. Gawlik, L. Ghizzoni, A. Buyukgeviz, Y. Rakover, T. Bistritzer *et al.* 2007. Clinical significance of the parental origin of the X chromosome in Turner syndrome. *Journal of Clinical Endocrinology and Metabolism* **92**:846–852.

Salerno, M. C., and J. C. Job. 1987. Height in Turner's syndrome: correlation with parents' height. *Archives Françaises de Pédiatrie* **44**:863–865.

Shaw, G. M., S. L. Carmichael, Z. Kaidarov, and J. A. Harris. 2003. Differential risks to males and females for congenital malformations among 2.5 million California births, 1989–1997. *Birth Defects Research. Part A, Clinical and Molecular Teratology* **67**:953–958.

Simpson, J. L., F. de la Cruz, R. S. Swerdlow, C. Samango-Sprouse, N. E. Skakkebaek, J. M. Graham, T. Hassold *et al.* 2003. Klinefelter syndrome: expanding the phenotype and identifying new research directions. *Genetics in Medicine* **5**:460–468.

Skuse, D. H. 1999. Genomic imprinting of the X chromosome: a novel mechanism for the evolution of sexual dimorphism. *Journal of Laboratory and Clinical Medicine* **133**:23–32.

Skuse, D. H. 2000. Imprinting, the X-chromosome, and the male brain: explaining sex differences in the liability to autism. *Pediatric Research* **47**:9–16.

Skuse, D. H. 2005. X-linked genes and mental functioning. *Human Molecular Genetics* **14** (Spec No. 1):r27–r32.

Skuse, D. H. 2006. Sexual dimorphism in cognition and behaviour: the role of X-linked genes. *European Journal of Endocrinology* **155**(Suppl. 1):s99–s106.

Skuse, D. H., R. S. James, D. V. Bishop, B. Coppin, P. Dalton, G. Aamodt-Leeper, M. Bacarese-Hamilton *et al.* 1997. Evidence from Turner's syndrome of an imprinted X-linked locus affecting cognitive function. *Nature* **387**:705–708.

Stemkens, D., T. Roza, L. Verrij, H. Swaab, M. K. van Werkhoven, B. Z. Alizadeh, R. J. Sinke, and J. C. Giltay. 2006. Is there an influence of X-chromosomal imprinting on the phenotype in Klinefelter syndrome? A clinical and molecular genetic study of 61 cases. *Clinical Genetics* **70**:43–48.

Sybert, V. P., and E. McCauley. 2004. Turner's syndrome. *New England Journal of Medicine* **351**:1227–1238.

Tassabehji, M. 2003. Williams-Beuren syndrome: a challenge for genotype-phenotype correlations. *Human Molecular Genetics* **12** (Spec No. 2):R229–R237.

Telvi, L., A. Lebbar, O. Del Pino, J. P. Barbet, and J. L. Chaussain. 1999. 45,X/46,XY mosaicism: report of 27 cases. *Pediatrics* **104**:304–308.

Temple, C. M., and R. A. Carney. 1993. Intellectual functioning of children with Turner syndrome: a comparison of behavioural phenotypes. *Developmental Medicine and Child Neurology* **35**:691–698.

Temple, I. K., V. Shrubb, M. Lever, H. Bullman, and D. J. G. Mackay. 2007. Isolated imprinting mutation of the DLK1/GTL2 locus associated with a clinical presentation of maternal uniparental disomy of chromosome 14. *Journal of Medical Genetics* **44**:637–640.

Thangavel, C., and B. H. Shapiro. 2007. A molecular basis for the sexually dimorphic response to growth hormone. *Endocrinology* **148**:2894–2903.

Thomas, N. S., and T. J. Hassold. 2003. Aberrant recombination and the origin of Klinefelter syndrome. *Human Reproduction Update* **9**:309–317.

Tsezou, A., C. Hadjithanasiou, D. Gourgiotis, A. Galla, E. Kavazarakis, A. Pasparaki, M. Kapsetaki *et al.* 1999. Molecular genetics of Turner syndrome: correlation with clinical phenotype and response to growth hormone therapy. *Clinical Genetics* **56**:441–446.

Uematsu, A., T. Yorifuji, J. Muroi, M. Kawai, M. Mamada, M. Kaji, C. Yamanaka *et al.* 2002. Parental origin of normal X chromosomes in Turner syndrome patients with various karyotypes: implications for the mechanism leading to generation of a 45,X karyotype. *American Journal of Medical Genetics* **111**:134–139.

Vallender, E. J., and B. T. Lahn. 2004. How mammalian sex chromosomes acquired their peculiar gene content. *BioEssays* **26**:159–169.

Van, P. L., V. K. Bakalov, and C. A. Bondy. 2006a. Monosomy for the X-chromosome is associated with an atherogenic lipid profile. *Journal of Clinical Endocrinology and Metabolism* **91**:2867–2870.

Van, P. L., V. K. Bakalov, A. R. Zinn, and C. A. Bondy. 2006b. Maternal X chromosome, visceral adiposity, and lipid profile. *Journal of the American Medical Association* **295**:1373–1374.

Vawter, M. P., P. D. Harvey, and L. E. Delisi. 2007. Dysregulation of X-linked gene expression in Klinefelter's syndrome and association with verbal cognition. *American Journal of Medical Genetics. Part B. Neuropsychiatric Genetics* **144B**:728–734.

Vrana, P. B., J. A. Fossella, P. Matteson, T. del Rio, M. J. O'Neill, and S. M. Tilghman. 2000. Genetic and epigenetic incompatibilities underlie hybrid dysgenesis in *Peromyscus*. *Nature Genetics* **25**:120–124.

Wikström, A. M., J. N. Painter, T. Raivio, K. Aittomäki, and L. Dunkel. 2006. Genetic features of the X chromosome affect pubertal development and testicular degeneration in adolescent boys with Klinefelter syndrome. *Clinical Endocrinology* **65**:92–97.

Wilke, M., I. Krägeloh-Mann, and S. K. Holland. 2007. Global and local development of gray and white matter volume in normal children and adolescents. *Experimental Brain Research* **178**:296–307.

Wolff, D. J., S. Schwartz, and L. Carrel. 2000. Molecular determination of X inactivation pattern correlates with phenotype in women with a structurally abnormal X chromosome. *Genetics in Medicine* **2**:136–141.

Xu, J., and C. M. Disteche. 2006. Sex differences in brain expression of X- and Y-linked genes. *Brain Research* **1126**:50–55.