

Genetic variation in nuclear and mitochondrial markers supports a large sex difference in lifetime reproductive skew in a lekking species

Yvonne I. Verkuil^{1,2}, Cedric Juillet², David B. Lank³, Fredrik Widemo⁴ & Theunis Piersma^{1,5}

¹Animal Ecology Group, Centre for Ecological and Evolutionary Studies, University of Groningen, PO Box 11103, Groningen 9700 CC, the Netherlands

²Department of Natural History, Royal Ontario Museum, 100 Queen's Park Crescent, Toronto, M5S 2C6 Ontario, Canada

³Department of Biological Sciences, Simon Fraser University, Burnaby, V5A 1S6 British Columbia, Canada

⁴Department of Fish, Wildlife and Environmental Studies, Swedish University of Agricultural Sciences, Umeå SE- 901 83, Sweden

⁵Department of Marine Ecology, NIOZ Royal Netherlands Institute for Sea Research, PO Box 59, Den Burg, Texel 1790 AB, the Netherlands

Keywords

Effective population size, *Philomachus pugnax*, ruff, sexual selection, shorebirds.

Correspondence

Yvonne I. Verkuil, Marine Evolution and Conservation, Centre for Ecological and Evolutionary Studies, University of Groningen, PO Box 11103, 9700 CC Groningen, the Netherlands.

Tel: 31 50 3632226;

Fax: 31 50 3635205;

E-mail: yvonne_verkuil@hotmail.com

Funding Information

CJ was supported by a Rebanks Postdoctoral Fellowship. DBL was supported by operating grants from the Natural Sciences and Engineering Research Council of Canada. YIV was supported by a start-up grant to TP from the University of Groningen, by NWO travel grants (R84-606, R87-307, R84-643), by the GUF/Gratama Foundation and by two grants of the Schure-Beijerinck Popping Foundation.

Received: 4 July 2014; Revised: 15 July 2014;

Accepted: 17 July 2014

Ecology and Evolution 2014 4(18): 3626–3632

doi: 10.1002/ece3.1188

Introduction

In a lek system, a small number of males obtain most of the copulations, and therefore the distribution of mating success is strongly positively skewed with a mode near zero (Höglund and Alatalo 1995). An issue of interest in studies of lek mating systems is the extent to which seasonal or apparent skews in mating success ultimately

Abstract

Sex differences in skews of vertebrate lifetime reproductive success are difficult to measure directly. Evolutionary histories of differential skew should be detectable in the genome. For example, male-biased skew should reduce variation in the biparentally inherited genome relative to the maternally inherited genome. We tested this approach in lek-breeding ruff (Class Aves, *Philomachus pugnax*) by comparing genetic variation of nuclear microsatellites (θ_n ; biparental) versus mitochondrial D-loop sequences (θ_m ; maternal), and conversion to comparable nuclear (N_e) and female (N_{ef}) effective population size using published ranges of mutation rates for each marker (μ). We provide a Bayesian method to calculate N_e ($\theta_n = 4N_e\mu_n$) and N_{ef} ($\theta_m = 2N_{ef}\mu_m$) using 95% credible intervals (CI) of θ_n and θ_m as informative priors, and accounting for uncertainty in μ . In 96 male ruffs from one population, N_e was 97% (79–100%) lower than expected under random mating in an ideal population, where $N_e:N_{ef} = 2$. This substantially lower autosomal variation represents the first genomic support of strong male reproductive skew in a lekking species.

produce a comparable skew in lifetime reproductive success (LRS) (Mackenzie et al. 1995; Kokko et al. 1998, 1999). Variance in LRS may be smaller than expected because mating skews at leks are not necessarily mirrored by a skew in paternity because of off-lek copulations. This is the case in the lekking Houbara bustard (*Chlamydotis undulata undulata*) (Lesobre et al. 2010). In addition, mating skews can be highly age dependent (e.g., Gibson

and Guinness 1980; McDonald 1989), multiple paternity of offspring can reduce the mating variance in polygynous species (Webster et al. 1995), and trade-offs against survivorship may occur.

Directly measuring the skew in LRS requires comprehensive paternity analyses for at least one generation, which is difficult or impossible in many study systems (but see e.g., DuVal 2012). However, an evolutionary history of sex differences in skew in LRS should be visible in the genome. Strong sexual selection should result in a strongly reduced effective population size for the selected sex; for male-lekking species, reduced neutral genetic diversity should occur in paternally only inherited genes (Chesser and Baker 1996). In a recent paper, approximately a 10% reduction in neutral genetic variation on the Z chromosome was found in two polygynous shorebird species (Corl and Ellegren 2012). Because lekking is generally associated with extreme polygyny, an even stronger reduction in variation is expected in lekking species. In the lekking Gunnison sage-grouse (*Centrocercus minimus*), the simulated reduction in effective population size due to male mating skew was at least 23%, while accounting for the variance in seasonal reproductive success in females, which was almost as high as in males due to a high rate of nest failure (Stiver et al. 2008). In the lek-breeding European treefrog (*Hyla arborea*), no sex differences in effective population size were found; in this species, a negative effect of the mating system was obscured by the much stronger effect of delayed maturity in both sexes (Broquet et al. 2009).

Here, we revisit the empirical support for the predicted reduction in paternally transmitted genetic variation in a lekking species (for predictions see: Chesser and Baker 1996; Charlesworth 2009; Corl and Ellegren 2012). The test case is the ruff (*Philomachus pugnax*), which has a well-described lek mating system. The species has (1) highly ornamented males that are much larger than the nonornamented females, (2) three genetically based alternative male morphs, and (3) strong annual reproductive skews in males (Hogan-Warburg 1966; van Rhijn 1991; Lank et al. 1995, 2013; Widemo 1998; Jukema and Piersma 2006), all of which points toward strong sexual selection on males (Shuster and Wade 2003). However, in studies of direct paternity of ruffs, many offspring could not be assigned to fathers (Lank et al. 2002; Thuman and Griffith 2005), indicating that copulations took place with nonsampled males at other leks or off-lek (Lank and Smith 1987; van Rhijn 1991), potentially reducing the extent of variance in mating success relative to that assumed from counting copulations at leks (see Widemo and Owens 1995). Potential negative trade-offs between mating success and longevity are undocumented. Therefore, the magnitude of skew in the lifetime reproductive success of males remains unknown.

We developed a method that can be generally applied to detect sex differences in skews in lifetime reproductive success in birds, when, or assuming that ecological knowledge is available to cautiously explore other deviations from equilibrium conditions, such as sex differences in dispersal, recruitment, survival, and selection. Our Bayesian approach compares genetic variation of biparentally inherited autosomal genes with genetic variation of a maternally only inherited mitochondrial gene and estimates the likelihood to reject the random mating hypothesis under which autosomal variation is larger than mitochondrial variation. The Bayesian framework accounts for variation and uncertainty in mutation rates using a range of published mutation rates for each of the genetic markers.

Materials and Methods

Sampling

Ruffs have slight population structure across a breeding range that extends from Western Europe to eastern Siberia (Verkuil et al. 2012). To avoid potential effects of population substructure, we confined the present analysis to samples collected at one breeding site, in Sweden. Blood samples were taken from the brachial vein of 96 territorial male ruffs captured with cannon nets between 1990 and 2001 on Gotland, Sweden (57°10'N, 18°20'E) (Widemo 1997; Thuman 2003). Blood samples were stored in 95% ethanol, and DNA was isolated by standard phenol–chloroform or Chelex extractions and stored at –20°C.

Choice of genetic markers

For this study, highly variable markers were used that are generally applied in population genetic studies: microsatellites or simple sequence repeats (SSRs), and sequences of the mitochondrial D-loop or control region. These mutate fast and are therefore adequate archives of recent genetic evolution (Avise 2004). Importantly, data on genetic variation collected with these markers are available for many study organisms. The disadvantage is that marker-specific mutation rates (μ) have to be used to convert genetic variation to comparable units, and estimates of μ often have large uncertainties. Our method explores the use of these markers while accounting for the uncertainty in μ .

Biparental loci

The biparentally transmitted genes used to genotype the 96 males were seven polymorphic autosomal microsatellites loci (nDNA): ruff1, ruff6, ruff10, ruff12, and ruff50 are published loci cloned from ruffs (Thuman et al. 2002), and SNIPE B2 (Sæther et al. 2007) and M2 (L.

Wennerberg, pers. comm.). PCR profiles and details on fragment analysis, and the assignment of alleles can be found in Thuman et al. (2002) and Sæther et al. (2007). Standard diversity indices were obtained with the program Migrate (Beerli 2006).

Maternal locus

The maternally inherited locus used in this study is the control region (CR) of the mitochondrial genome (mtDNA). A 512 nt segment of domain I and II of the CR was amplified and sequenced with forward primer L141 (5'-TCCATT AATCTACAACCGGGCT) and reverse primer PropR (5'-A ATACCAGCTTGGGAGTTGG). Ruffs have a duplicated CR with high sequence similarity between copies (Verkuil et al. 2010). By anchoring the primer PropR in the tRNA^{Pro}, only CR1 was targeted. The amplification profile used was 2 min denaturation at 95°C, followed by 36 cycles of 94°C for 45 s, 53–57°C for 45 s, 72°C for 1.30 min, followed by a final 7 min elongation at 72°C. PCR products were gel-purified and prepared for sequencing using Big-Dye Terminal Cycle Sequencing reagents according to the manufacturer's instructions (Applied Biosystems, Foster City, CA, USA) and sequenced on an ABI 3100 automated sequencer. Sequences were edited and aligned in MEGA 3.1 (Kumar et al. 2004), and standard diversity indices were obtained with ARLEQUIN, version 3.11 (Excoffier and Schneider 2005).

Estimation of Θ with coalescent analysis

To obtain the effective population size from the genetic data, the following equations need to be applied,

$$\theta_n = 4N_e\mu_n, \text{ for nDNA,} \quad (1)$$

$$\theta_m = 2N_{ef}\mu_m, \text{ for mtDNA,} \quad (2)$$

where θ_n and θ_m are the respective nuclear and mitochondrial genetic variance parameters (i.e., sequence diversity that evolved under a species-specific effective population size and mutation rate), N_e and N_{ef} are biparental and female-transmitted only effective population sizes, and μ_n and μ_m the marker-scaled mutation rates (Hartl and Clark 1997: Pp. 121–127). The factor four in equation 1 accounts for the ploidy difference with mtDNA and the estimation over two branches. The factor two in equation 2 accounts for the estimation of θ over two branches of the ancestral tree.

Estimates of the variance parameters θ_n and θ_m were obtained by coalescent analysis (Beerli and Felsenstein 1999), using the program Migrate-n version 3.5.1 (Beerli 2006). The genotypes and haplotypes were considered to come from a genetically uniform population, because we

used males from a single breeding population. For microsatellites, the Bayesian Brownian motion model was used, which is a continuous modification of the discrete stepwise mutation model (Kuhner 2006). After initial runs (with various random seed numbers) to determine likely priors for θ_n leading to unimodal posterior distributions, one final long run was executed with a uniform prior distribution for θ of [0;100]. After a burn-in of 20,000 steps, the parameter space was sampled by one adaptively heated chain, sampling 200,000 trees, with an increment of 100; a total of 300 replicates were run, meaning that in total 6 billion parameter values were visited. Settings were adjusted to allow substitution rates to vary among loci (P. Beerli, pers. comm.).

For mtDNA sequences, Bayesian interference was also used. The substitution rate was assumed to be constant between sites and the transition/transversion bias was set to $R = 33$ (R was 33 in a larger sample of all breeding birds, see Verkuil et al. (2012)). After initial runs to test for convergence under different priors, a uniform prior distribution of [0.0;0.1] was set for theta. One final run was performed with a burn-in of 10,000 and a static heating scheme of four temperatures, sampling 10 million trees with an increment of 100; consequently, 1 billion parameter values were visited.

Modeling $N_e:N_{ef}$ under uncertainty in both mutation rates and genetic variance parameters

The quantity $N_e:N_{ef}$ is provided by equations 1 and 2,

$$N_e:N_{ef} = [\theta_n/(4\mu_n)]/[\theta_m/(2\mu_m)] \quad (3)$$

However, the uncertainty in the parameters θ and μ can substantially affect the estimation of N_e and N_{ef} from genetic data. To estimate $N_e:N_{ef}$, we used OpenBUGS, a Bayesian analyses dedicated software that makes the MCMC method easily accessible (Spiegelhalter et al. 2010). For calculation of N_e , a generation time of 3 years was assumed (Colwell 2010). In the computation of $N_e:N_{ef}$, we incorporated the uncertainty of published mutation rates (μ_n and μ_m) and the genetic variance parameters (θ_n and θ_m), estimated from the data. Instead of using single fixed values, we defined two uniform prior distributions, one for θ_n and one for θ_m . The uniform distributions were defined over a range of values that included 95% of their respective posterior distribution obtained by coalescent analysis of the genetic data ($U[10.0;16.5]$ and $U[0.01;0.05]$, respectively, see Results).

For the mutation rates μ_n and μ_m , we defined, a priori, the range of possible mutation rates according to the literature ($[1 \times 10^{-5}; 1 \times 10^{-3}]$ and $[1 \times 10^{-8}; 1 \times 10^{-7}]$, respectively). This means that the high variability of

mutation rates, as reported in the literature, is the starting point of the Bayesian analysis. To determine the ranges of mutation rates, we used the following information. In vertebrates, the typical mutation rate for microsatellites is 1×10^{-4} substitutions/locus/generation (Ellegren 2000; Whittaker et al. 2003). Birds have fewer microsatellites than mammals, but characteristics such as length, allele dispersion, and range of allele sizes do not vary between birds and mammals, indicating that mutation rates might be comparable (Neff and Gross 2001). Recently, fast microsatellite mutation rates have been reported in one Falconiformes and three Passeriformes, ranging between 3.0×10^{-3} and 1.3×10^{-2} (Brohede et al. 2002; Beck et al. 2003; Ortego et al. 2008; Masters et al. 2011); the variation in mutation rates between these four studies strongly correlated with median allele spans ([24–140] bp). Our microsatellites had a median allele span of 36 bp (Verkuil et al. 2012). We therefore assumed an upper boundary for μ_n of 1×10^{-3} . To stay conservative, a lower boundary of 1×10^{-5} was assumed.

For mtDNA, a μ_m of 2–15% per Ma was assumed, which translates into [2×10^{-8} ; 1.5×10^{-7}] substitutions/locus/year (using control region sequence of 512 nt). This range includes observed values for shorebirds for domain I of the control region (Wenink and Baker 1996) and also the lower mutation rate in domain II (Buehler and Baker 2005). In the results, the term “CI” corresponds to the credible interval of estimates.

Results

Genotypic data

Ninety-six individuals were successfully genotyped at seven microsatellite loci. None of the loci had null-alleles, significant linkage disequilibrium, or significant deviations from Hardy–Weinberg equilibrium (see Thuman et al. 2002). The number of alleles per locus was 9, 14, 10, 5, 12, 20, and 15, respectively. The expected heterozygosity (H_{exp}) was 0.695 (0.419–0.845).

Eighteen individuals were successfully sequenced for the 512-bp segment of the mitochondrial control region. The segment had sixteen polymorphic sites (15 transitions and 1 transversion), and across the samples 14 different haplotypes were detected (GenBank under Accession Nos HQ171099–HQ171161). The gene diversity (H_s) was 0.954 ± 0.039 , and the average nucleotide diversity was 0.007 ± 0.004 .

Estimates of theta

The coalescent run to estimate theta from microsatellites had an effective MCMC sample size of 609 million and

returned an unimodal posterior distribution with a global estimate of θ_n of 13.23 (95% CI 10.2–16.6). For mtDNA, the effective MCMC sample size was 5.5 million, with resulted in a unimodal posterior distribution, and a most probable estimate of θ_m of 0.0238 (95% CI 0.012–0.048).

Ratio of biparental to maternal effective population size

Including the 95% CIs of the estimates of θ_n and θ_m , the ratio $N_e:N_{ef}$ was 0.068 (SD = 0.168, 95% CI 0.004–0.425), assuming prior uniform distributions for μ_m and μ_n as given in the Methods. This means that in ruffs, nuclear genetic variation is 97% (range: 79–100%) lower than expected under random mating in an ideal population, where $N_e:N_{ef} = 2$.

Discussion

The 97% reduction in nuclear genetic variation is empirical support for the assumed stronger interindividual variance in lifetime reproductive success (LRS) of male ruffs relative to females. This result confirms the intuition of previous field researchers who observed the large investments by some males in dominating the lek (Hogan-Warburg 1966; van Rhijn 1983, 1991). Males may display so intensively that they refrain from eating and drinking, and lose mass throughout the breeding season (Widemo 1998; Bachman and Widemo 1999). Our study suggests that for the winners, male competition indeed does pay off in a high LRS. Annual and lifetime reproductive skew probably exists within all three male morphs (independents, satellites and faeders, see Jukema and Piersma 2006; Lank et al. 2013), but the skew among independents will drive the reduction of genetic variation, because they comprise ca. 80–85% of the population (Lank et al. 1995; Widemo 1998). Strong skews are hypothesized to facilitate invasion of alternative reproductive strategies (Shuster and Wade 2003), as clearly has occurred in this case. The results show that in ruffs sex-specific fitness variation (sexual selection) has been strong and stable long enough for the nuclear genetic variation to be much lower than the mitochondrial genetic variation inherited through the female line.

Our approach to evaluate the degree of difference in genetic variation between the nuclear and mitochondrial genome through estimating the effective population size (N_e) can be applied more generally. The degree of difference in N_e between the two genomes will summarize longer-term mating skews between the sexes in any species. However, knowledge of the ecology of the species is a prerequisite, because sex differences in dispersal, recruitment, survival, and selection may obscure the effect of the mating

system (see Chesson and Baker 1996; Barton 2000; Waples 2010). We will use ruffs to illustrate the possible effects of sex differences in ecology on the reduction in N_e .

Stronger male than female dispersal will reduce N_e . For example in an extreme case, 90% of the dispersal could take place through males, which would reduce N_e by as much as 65% (expected $N_e:N_{ef}$ changes from 2 to 0.7, see the mammal model of Chesson and Baker 1996). Theoretically, lekking bird species are assumed to have female-biased dispersal (Greenwood and Harvey 1982; Chesson and Baker 1996), but empirical data are inconclusive on this prediction (see Lebigre et al. 2008; Martín et al. 2008). In ruffs, female natal philopatry is extremely low (Andersen 1948, 1951); adults of both sexes show large geographic flexibility in migration routes (Rakhimberdiev et al. 2011). At the same time, adult territorial males in our study population show substantial lek site fidelity between years (Widemo 1997) and were never observed to shift to another breeding area once established (Lif 2002; F. Widemo, unpublished data). Therefore, it is unlikely that ruffs have strongly male-biased dispersal.

Adult sex ratio can also influence N_e . Ruffs have an assumed skewed adult sex ratio of ca. 40% males to 60% females (OAG Münster 1996; Zwarts et al. 2009), which may result from a sex bias in recruitment (Jaatinen et al. 2010), as found in other polygynous birds (Donald 2007; see Kokko and Jennions 2008). This sex ratio bias would lead to a reduction in N_e of 35%, even under random mating (expected $N_e:N_{ef}$ changes from 2 to 1.3).

In principle, $N_e:N_{ef}$ can also be influenced by differential selection regimes in the two genomes and/or the Hill-Robertson interference; mtDNA does generally not recombine and is therefore susceptible to genetic sweeps reducing variation (Marais 2007). This would lead to an underestimate of N_{ef} . Also, if genetic diversity was not at mutation/drift equilibrium (e.g., the species is recovering from a severe population bottleneck), microsatellites would have higher diversity than mitochondrial genome because mutation rate rather than N_e would control the speed of recovery. As we found relatively high variation in mtDNA, in ruffs neither process can be likely responsible for the observed low $N_e:N_{ef}$ ratio.

We conclude that in ruffs, the observed reduction in N_e of 97% (range: 79–100%) is stronger than can be accounted for by ecological differences between males and females, and primarily results from the larger variance in LRS in males than in females, which is congruent with the predicted $N_e:N_{ef}$ under the lek model of Chesson and Baker (1996). This is not the first study assessing the reduction of genetic variation in a lekking species (e.g., Stiver et al. 2008; Broquet et al. 2009; Corl and Ellegren 2012), but it is the first to detect a reduction. Ruffs were included in a comparison of genetic diversity at autoso-

mal and Z-linked introns of monogamous and polygynous shorebird species (Corl and Ellegren 2012), but had few variable sites compared with the other species, and the authors of this earlier study considered the results inconclusive for ruff. We showed that a long-term mating skew can be genetically assessed in ruffs using microsatellites and the mitochondrial control region, which have a higher mutation rates than the nuclear introns used in the earlier study. The uncertainty in mutation rate of these markers should be accounted for with methodology such as we presented. Using our approach, existing and future genotyping datasets from other species may be used to characterize their mating systems. The method will be most useful if information on other potential biasing ecological factors is available, but mating patterns themselves cannot be directly assessed.

Acknowledgments

We want to thank Allan Baker and Oliver Haddrath for excellent training, and for help in the laboratory. We thank Kathrin Thuman for collecting and sharing samples. CJ was supported by a Rebanks Postdoctoral Fellowship. DBL was supported by operating grants from the Natural Sciences and Engineering Research Council of Canada. YIV was supported by a start-up grant to TP from the University of Groningen, by NWO travel grants (R84-606, R87-307, R84-643), by the GUF/Gratama Foundation and by two grants of the Schure-Beijerinck Popping Foundation. Allan Baker, Debbie Buehler and the Groningen Shorebird Lunch group (Jesse Conklin, Jeroen Reneerkens, Lucie Schmaltz, Nathan Senner, and Mo Verhoeven) participated in helpful discussions. Hanna Kokko, Jesse Conklin, Robert Gibson, and three anonymous reviewers provided constructive feedback.

Data Accessibility

DNA sequences: GenBank Accession Nos: HQ171099-HQ171161.

Conflict of Interest

None declared.

References

- Andersen, F. S. 1948. Contributions to the biology of the ruff (*Philomachus pugnax* (L)) II Dansk. Ornitologisk Forenings Tidsskrift 42:125–148.
- Andersen, F. S. 1951. Contributions to the biology of the ruff (*Philomachus pugnax* (L)) III Dansk. Ornitologisk Forenings Tidsskrift 45:145–173.

Avise, J. C. 2004. Molecular markers, natural history, and evolution. Sinauer Associates Inc., Sunderland, Massachusetts.

Bachman, G., and F. Widemo. 1999. Relationships between body composition, body size and alternative reproductive tactics in a lekking sandpiper, the ruff (*Philomachus pugnax*). *Funct. Ecol.* 13:411–416.

Barton, N. H. 2000. Genetic hitchhiking. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 355:1553–1562.

Beck, N. R., M. C. Double, and A. Cockburn. 2003. Microsatellite evolution at two hypervariable loci revealed by extensive avian pedigrees. *Mol. Biol. Evol.* 20:54–61.

Beerli, P. 2006. Comparison of Bayesian and maximum-likelihood inference of population genetic parameters. *Bioinformatics* 22:341–345.

Beerli, P., and J. Felsenstein. 1999. Maximum-likelihood estimation of migration rates and effective population numbers in two populations using a coalescent approach. *Genetics* 152:763–773.

Brohede, J., C. R. Primmer, A. Møller, and H. Ellegren. 2002. Heterogeneity in the rate and pattern of germline mutation at individual microsatellite loci. *Nucleic Acids Res.* 30:1997–2003.

Broquet, T., J. Jaquière, and N. Perrin. 2009. Opportunity for sexual selection and effective population size in the lek-breeding European treefrog (*Hyla arborea*). *Evolution* 63:674–683.

Buehler, D. M., and A. J. Baker. 2005. Population divergence time and historical demography in red knots and dunlins. *Condor* 107:497–513.

Charlesworth, B. 2009. Effective population size and patterns of molecular evolution and variation. *Nat. Rev. Genet.* 10:195–205.

Chesser, R. K., and R. J. Baker. 1996. Effective sizes and dynamics of uniparentally and diparentally inherited genes. *Genetics* 144:1225–1235.

Colwell, M. A. 2010. Shorebird ecology, conservation and management. University of California Press, Berkeley and Los Angeles, USA.

Corl, A., and H. Ellegren. 2012. The genomic signature of sexual selection in the genetic diversity of the sex chromosome and autosomes. *Evolution* 66:2138–2149.

Donald, P. F. 2007. Adult sex ratios in wild bird populations. *Ibis* 149:671–692.

DuVal, E. H. 2012. Variation in annual and lifetime reproductive success of lance-tailed manakins: alpha experience mitigates effects of senescence on siring success. *Proc. R. Soc. Lond. B Biol. Sci.* 279:1551–1559.

Ellegren, H. 2000. Microsatellite mutations in the germline: implications for evolutionary inference. *Trends Genet.* 16:551–558.

Excoffier, L. G. L., and S. Schneider. 2005. Arlequin ver. 3.0: an integrated software package for population genetics data analysis. *Evol. Bioinform. Online* 1:47–50.

Gibson, R. M., and F. E. Guinness. 1980. Differential reproduction among red deer *Cervus elaphus* stags on Rhum. *J. Anim. Ecol.* 49:199–208.

Greenwood, P. J., and P. H. Harvey. 1982. The natal and breeding dispersal of birds. *Annu. Rev. Ecol. Syst.* 13:1–21.

Hartl, D. L., and A. G. Clark. 1997. Principles of population genetics, 4th edn. Sinauer Associates, Inc., Sunderland, USA.

Hogan-Warburg, A. J. 1966. Social behavior of ruff *Philomachus pugnax*. *Ardea* 54:109–229.

Höglund, J., and R. V. Alatalo. 1995. Leks. Princeton University Press, Princeton, USA.

Jaatinen, K., A. Lehtinen, and D. B. Lank. 2010. Female-biased sex ratios and the proportion of cryptic male morphs of migrant juvenile ruffs (*Philomachus pugnax*) in Finland. *Ornis Fennica* 87:125–134.

Jukema, J., and T. Piersma. 2006. Permanent female mimics in a lekking shorebird. *Biol. Lett.* 2:161–164.

Kokko, H., and M. D. Jennions. 2008. Parental investment, sexual selection and sex ratios. *J. Evol. Biol.* 21:919–948.

Kokko, H., W. J. Sutherland, J. Lindström, J. D. Reynolds, and A. Mackenzie. 1998. Individual mating success, lek stability, and the neglected limitations of statistical power. *Anim. Behav.* 56:755–762.

Kokko, H., A. Mackenzie, J. D. Reynolds, J. Lindström, and W. J. Sutherland. 1999. Measures of inequality are not equal. *Am. Nat.* 154:358–382.

Kuhner, M. K. 2006. LAMARC 2.0: maximum likelihood and Bayesian estimation of population parameters. *Bioinformatics* 22:768–770.

Kumar, S., K. Tamura, and M. Nei. 2004. MEGA3: integrated software for molecular evolutionary genetics analysis and sequence alignment. *Brief. Bioinform.* 5:150–163.

Lank, D. B., and C. M. Smith. 1987. Conditional lekking in ruff (*Philomachus pugnax*). *Behav. Ecol. Sociobiol.* 20:137–145.

Lank, D. B., C. M. Smith, O. Hanotte, T. Burke, and F. Cooke. 1995. Genetic polymorphism for alternative mating behavior in lekking male ruff *Philomachus pugnax*. *Nature* 378:59–62.

Lank, D. B., C. M. Smith, O. Hanotte, A. Ohtonen, S. Bailey, and T. Burke. 2002. High frequency of polyandry in a lek mating system. *Behav. Ecol.* 13:209–215.

Lank, D. B., L. L. Farrell, T. Burke, T. Piersma, and S. B. McRae. 2013. A dominant allele controls development into female mimic male and diminutive female ruffs. *Biol. Lett.* 9:20130653.

Lebigre, C., R. V. Alatalo, H. E. Forss, and H. Siitari. 2008. Low levels of relatedness on Black Grouse leks despite male philopatry. *Mol. Ecol.* 17:4512–4521.

Lesobre, L., F. Lacroix, E. Le Nuz, Y. Hingrat, T. Chalah, and M. Saint Jaime. 2010. Absence of male reproductive skew, along with high frequency of polyandry and conspecific brood parasitism in the lekking Houbara bustard *Chlamydotis undulata undulata*. *J. Avian Biol.* 41:117–126.

Lif, M. 2002. Dispersal, recruitment and survival in the fragmented population of ruff, *Philomachus pugnax*, on Gotland. Master Thesis, Uppsala University.

Mackenzie, A., J. D. Reynolds, V. J. Brown, and W. J. Sutherland. 1995. Variation in male mating success on leks. *Am. Nat.* 145:633–652.

Marais, G. A. B. 2007. Sex chromosomes and mitochondrial DNA polymorphism in birds: the Hill-Robertson effects extend from nucleus to mitochondria. *Heredity* 99:357–358.

Martín, C. A., J. C. Alonso, J. A. Alonso, C. Palacín, M. Magaña, and B. Martín. 2008. Natal dispersal in great bustards: the effect of sex, local population size and spatial isolation. *J. Anim. Ecol.* 77:326–334.

Masters, B. S., L. S. Johnson, B. G. P. Johnson, J. L. Brubaker, S. K. Sakaluk, and C. F. Thompson. 2011. Evidence for heterozygote instability in microsatellite loci in house wrens. *Biol. Lett.* 7:127–130.

McDonald, D. B. 1989. Cooperation under sexual selection: age-graded changes in a lekking bird. *Am. Nat.* 134:709–730.

Münster, O. A. G. 1996. Do females really outnumber males in ruff *Philomachus pugnax* wintering in Africa? *J. Ornithol.* 137:91–100.

Neff, B. D., and M. R. Gross. 2001. Microsatellite evolution in vertebrates: inference from AC dinucleotide repeats. *Evolution* 55:1717–1733.

Ortego, J., J. M. Aparicio, P. J. Cordero, and G. Calabuig. 2008. Characteristics of loci and individuals are associated with mutation rates in lesser kestrels (*Falco naumanni*). *Mutat. Res.* 648:82–86.

Rakhimberdiev, E. N., Y. I. Verkuil, A. A. Saveliev, R. A. Väistönen, J. V. Karagicheva, M. Y. Soloviev, et al. 2011. A global population redistribution in a migrant shorebird detected with continent-wide qualitative breeding survey data. *Divers. Distrib.* 17:144–151.

van Rhijn, J. G. 1983. On the maintenance and origin of alternative strategies in the ruff *Philomachus pugnax*. *Ibis* 125:482–498.

van Rhijn, J. G. 1991. The Ruff. Individuality in a gregarious wading bird. Poyser, London.

Sæther, S. A., P. Fiske, J. A. Kållås, A. Kuresoo, L. Luigujoe, S. B. Piertney, et al. 2007. Inferring local adaptation from Q_{ST} – F_{ST} comparisons: neutral genetic and quantitative trait variation in European populations of great snipe. *J. Evol. Biol.* 20:1563–1576.

Shuster, S. M., and M. J. Wade. 2003. Mating Systems and Strategies. Princeton University Press, Princeton.

Spiegelhalter, D., A. Thomas, N. Best, and D. Lunn. 2010. OpenBUGS user manual Version 312. Technical report. Medical Research Council Biostatistics Unit, Cambridge, UK. Available at: <http://www.openbugs.net> (accessed 21 April 2014).

Stiver, J. R., A. D. Apa, T. E. Remington, and R. M. Gibson. 2008. Polygyny and female breeding failure reduce effective population size in the lekking Gunnison sage grouse. *Biol. Conserv.* 141:472–481.

Thuman, K. A. 2003. Female reproductive strategies in the Ruff (*Philomachus pugnax*). PhD thesis, Uppsala University.

Thuman, K. A., and S. C. Griffith. 2005. Genetic similarity and the nonrandom distribution of paternity in a genetically highly polyandrous shorebird. *Anim. Behav.* 69:765–770.

Thuman, K. A., F. Widemo, and S. B. Piertney. 2002. Characterization of polymorphic microsatellite DNA markers in the ruff (*Philomachus pugnax*). *Mol. Ecol. Notes* 2:276–277.

Verkuil, Y. I., T. Piersma, and A. J. Baker. 2010. A novel mitochondrial gene order in shorebirds (Scolopacidae, Charadriiformes). *Mol. Phylogenet. Evol.* 57:411–416.

Verkuil, Y. I., T. Piersma, J. Jukema, J. C. E. W. Hooijmeijer, L. Zwarts, and A. J. Baker. 2012. The interplay between habitat availability and population differentiation: a case study on genetic and morphological structure in an inland wader (Charadriiformes). *Biol. J. Linn. Soc.* 106:641–656.

Waples, R. S. 2010. Spatial-temporal stratifications in natural populations and how they affect understanding and estimation of effective population size. *Mol. Ecol.* 10:785–796.

Webster, M. S., S. Pruitt-Jones, D. F. Westneat, and S. J. Arnold. 1995. Measuring the effects of pairing success, extra-pair copulations and mate quality on the opportunity for sexual selection. *Evolution* 49:1147–1157.

Wenink, P. W., and A. J. Baker. 1996. Mitochondrial DNA lineages in composite flocks of migratory and wintering dunlins (*Calidris alpina*). *Auk* 113:744–756.

Whittaker, J. C., R. M. Harbord, N. Boxall, I. Mackay, G. Dawson, and R. M. Sibley. 2003. Likelihood-based estimation of microsatellite mutation rates. *Genetics* 164:781–787.

Widemo, F. 1997. The social implications of traditional use of lek sites in the ruff *Philomachus pugnax*. *Behav. Ecol.* 8:211–217.

Widemo, F. 1998. Alternative reproductive strategies in the ruff, *Philomachus pugnax*: a mixed ESS? *Anim. Behav.* 56:329–336.

Widemo, F., and I. P. F. Owens. 1995. Lek size, male mating skew and the evolution of lekking. *Nature* 373:148–151.

Zwarts, L., R. Bijlsma, J. van der Kamp, and E. Wymenga. 2009. Living on the edge. Wetlands and birds in changing Sahel. KNNV Publishing, Zeist, The Netherlands.