

Classical Mechanics

Lecture 14

Today's Concepts:

- a) Rotational Motion
- b) Moment of Inertia

Your Comments

Will we have to memorize all the different formulas for moment of inertia?

What is a moment of inertia?

Can we go over the equations more thoroughly such as kinetic energy?

Will we be required to memorize all the fractions of different objects' moment of inertia, or are we expected to be able to derive the integral?

A review of all the concepts would be nice, especially the moment of inertia. And some homework problems as well please

But the bugs on the first question were a bit too realistic looking. How about a cute, non threatening example next time?

It was ALL so very confusing!! :'(

"oooh pick me pick me pick me!"-Donkey from Shrek. Actually, Where did the constants of proportionality come from when going from a solid cylinder to a cylindrical shell? also, can we do more find the moment of inertia examples?



What is a Radian?

- A) A measure of angle
- B) The distance from the centre of a circle to its circumference
- C) A brand of space heater
- D) Something like a jelly fish



What are the units of a Radian?

A) m

B) m^{-1}

C) m/degree

D) degree/m


E) **none**

Review




About how large is a radian in degrees?

- A) 0.6°
- B) 6.0°
- C) 60°
- D) 360°
- E) 600°



Summary of Rotations



Arc Length

$$s = R\theta$$

where θ is measured in radians

Tangential Speed

$$v = R\omega$$

Tangential Acceleration

$$a = R\alpha$$

For constant angular acceleration

$$\theta = \theta_o + \omega_o t + \frac{1}{2} \alpha t^2$$

$$\omega = \omega_o + \alpha t$$

$$\omega^2 - \omega_o^2 = 2\alpha(\theta - \theta_o)$$

Rotational Kinetic Energy

$$K_{\text{system}} = \frac{1}{2} I \omega^2$$

Angular velocity ω is measured in radians/sec

Frequency f is measured in revolutions/sec

1 revolution = 2π radians

Period $T = 1/f$



$$\omega = \frac{2\pi}{T}$$

Another Summary


$$\alpha = \frac{d\omega}{dt}$$

For constant angular acceleration

$$\theta = \theta_o + \omega_o t + \frac{1}{2} \alpha t^2$$

$$\omega = \omega_o + \alpha t$$

$$\omega^2 - \omega_o^2 = 2\alpha(\theta - \theta_o)$$



Constant α does not mean constant ω

Clicker Question




A disk spins at 2 revolutions/sec.

What is its period?

A) $T = 2 \text{ sec}$

B) $T = 2\pi \text{ sec}$

C) $T = \frac{1}{2} \text{ sec}$



Clicker Question




A disk spins at 2 revolutions/sec.

What is its angular velocity?

A) $\omega = 2\pi$ rad/sec

B) $\omega = \frac{\pi}{2}$ rad/sec

C) $\omega = 4\pi$ rad/sec




CheckPoint

A wheel which is initially at rest starts to turn with a constant angular acceleration. After 4 seconds it has made 4 complete revolutions.

How many revolutions has it made after 8 seconds?

- A) 8 B) 12 C) 16



Only about half got this right
so let's try again...


CheckPoint Response




After 4 seconds it has made 4 complete revolutions.

How many revolutions has it made after 8 seconds?

- A) 8 B) 12 C) 16



Spinning Wheel: Question 1 (N = 93)



A) Since it made 4 revolutions in 4 seconds, its angular velocity is 1 revolution per second. Therefore, in 8 seconds, it will have made 8 revolutions.

B) it makes 4 in the first 4 seconds and then 4 + 4 in the second 4 seconds.
 $4 + 4 + 4 = 12$

C) The number of revolutions is proportional to time squared.

Calculating Moment of Inertia


Moment of Inertia

For Discrete Distributions

$$I \equiv \sum m_i r_i^2$$

For Continuous Distributions

$$I = \int r^2 dm$$



Moment of Inertia

$$I \equiv \sum m_i r_i^2$$

$$I_y = I_x = 4ML^2$$

$$I_z = 8ML^2$$

Depends on
rotation axis

CheckPoint


A triangular shape is made from identical balls and identical rigid, massless rods as shown. The moment of inertia about the a , b , and c axes is I_a , I_b , and I_c respectively.

Which of the following orderings is correct?

A) $I_a > I_b > I_c$

B) $I_a > I_c > I_b$

C) $I_b > I_a > I_c$



Only about half got this right
so let's try again...


CheckPoint Response



A) $I_a > I_b > I_c$

B) $I_a > I_c > I_b$


C) $I_b > I_a > I_c$



A) $I_a = 8mr^2$ $I_b = 3mr^2$ $I_c = 2mr^2$

B) $I_a = 8mr^2$ $I_b = 3mr^2$ $I_c = 4mr^2$

Triangle of Spheres: Question 1 (N = 92)



Calculation Moment of Inertia

Moment of Inertia

For Discrete Distributions

$$I \equiv \sum m_i r_i^2$$

For Continuous Distributions

$$I = \int r^2 dm$$

Solid Cylinder

$$I = \frac{1}{2} MR^2$$



Cylindrical Shell

$$I = MR^2$$



Solid Sphere

$$I = \frac{2}{5} MR^2$$



Spherical Shell

$$I = \frac{2}{3} MR^2$$



Bigger when
the mass is
further out

Clicker Question




In both cases shown below a hula hoop with mass M and radius R is spun with the same angular velocity about a vertical axis through its center. In **Case 1** the plane of the hoop is parallel to the floor and in **Case 2** it is perpendicular.


In which case does the spinning hoop have the most kinetic energy?

- A) Case 1 B) Case 2 C) Same

Case 1



Case 2



Only about half got this right so let's try again...


Clicker Question



In which case does the spinning hoop have the most kinetic energy?

- A) Case 1 B) Case 2 C) Same


Case 1



Case 2

$$K = \frac{1}{2} I \omega^2$$

Rotating Hoop: Question 1 (N = 90)



- A) In case one, more mass is located away from its axis, so it has larger moment of inertia. Therefore it has more kinetic energy.
- C) The radius, angular velocities and masses are the same so the inertia is the same which means the kinetic energy is the same for both.

Clicker Question




A mass M is uniformly distributed over the length L of a thin rod. The mass inside a short element dx is given by:

A) $M dx$

B) $\frac{dx}{M}$

C) $\frac{M}{L} dx$

D) $\frac{L}{M} dx$



analyze the dimensions!

Clicker Question




A mass M is uniformly distributed over the length L of a thin rod. The contribution to the rod's moment of inertia provided by element dx is given by:

A) $x^2 \frac{M}{L} dx$

B) $\frac{1}{x^2} \frac{M}{L} dx$

C) $\frac{M}{L} dx^2$



Clicker Question




A disk has a radius R . The area of a thin ring inside the disk with radius r and thickness dr is:


A) $\pi r^2 dr$

B) $2\pi r dr$

C) $4\pi r^3 dr$

analyze the dimensions!





- (i) $\theta = \theta_0 + \omega_0 t + \frac{1}{2} \alpha t^2$
- (ii) $\omega = \omega_0 + \alpha t$
- (iii) $\omega^2 = \omega_0^2 + 2\alpha \Delta\theta$

A disk with mass $m = 11.2$ kg and radius $R = 0.34$ m begins at rest and accelerates uniformly for $t = 17.4$ s, to a final angular speed of $\omega = 29$ rad/s.

1) What is the angular acceleration of the disk?

 rad/s²

Using (ii)


$$\alpha = \frac{\omega - \omega_0}{t}$$

2) What is the angular displacement over the 17.4 s?

 rad

Using (i)

$$\theta = \frac{1}{2} \alpha t^2$$



$$(iv) \quad I_{DISK} = \frac{1}{2} MR^2$$

$$(v) \quad K = \frac{1}{2} I \omega^2$$

A disk with mass $m = 11.2$ kg and radius $R = 0.34$ m begins at rest and accelerates uniformly for $t = 17.4$ s, to a final angular speed of $\omega = 29$ rad/s.

3) What is the moment of inertia of the disk?


kg-m²

Use (iv)

4)  What is the change in rotational energy of the disk?

J

Use (v)



$$(vi) \quad d = R\theta$$

$$(vii) \quad v = R\omega$$

$$(viii) \quad a_T = R\alpha$$


$$(ix) \quad a_c = \frac{v^2}{R} = \omega^2 R$$

A disk with mass $m = 11.2$ kg and radius $R = 0.34$ m begins at rest and accelerates uniformly for $t = 17.4$ s, to a final angular speed of $\omega = 29$ rad/s.

5) What is the tangential component of the acceleration of a point on the rim of the disk when the disk has accelerated to half its final angular speed?


 m/s²

Use (viii)

6)  What is the radial component of the acceleration of a point on the rim of the disk when the disk has accelerated to half its final angular speed?

 m/s²

Use (ix)



$$(vi) \quad d = R\theta$$

$$(vii) \quad v = R\omega$$

$$(viii) \quad a_T = R\alpha$$


$$(ix) \quad a_c = \frac{v^2}{R} = \omega^2 R$$

A disk with mass $m = 11.2$ kg and radius $R = 0.34$ m begins at rest and accelerates uniformly for $t = 17.4$ s, to a final angular speed of $\omega = 29$ rad/s.

7) What is the final speed of a point on the disk half-way between the center of the disk and the rim?

 m/s

Use (vii)

8)  What is the total distance a point **on the rim** of the disk travels during the 17.4 seconds?

 m

Use (vi)