


Today: Continuation of Ch 4

First: Recall from last time....



p 180 Note the pairs of graphs

Can you see any dependence of Abrasion Loss on Tensile Strength and Hardness?

P 181 Note role of variables – 1 dependent, 2 independent

Coordinate system for panels (i,j) i is number of panel left to right

j is number of panel bottom to top

(just as in usual coordinate system)

p 182 concept of slice of one variable (hardness) to examine relationship between two other variables (abrasion loss and tensile strength). Conditioning plot = "coplot"

p 183 Does the relationship between abrasion loss and tensile strength depend on hardness? i.e. Do Tensile Strength and Hardness interact in determining Abrasion Loss?

pp 184-5 Putting several slices together. Panels show response to increasing hardness, from lower left to upper right. (1,1), (2,1), (3,1), (1,2), (2,2), (3,2) is the order of the plots corresponding to the increasing slices.

Note robust fitting of loess curves to each slice. Reveals possible interaction. What if highest hardness range were missing?


pp 186-7 This time condition on Tensile Strength. Slight interaction less obvious.

p188 Ethanol Data:

Row	NOX	CR	ER	ER="richness, or fuel/air" CR="compression ratio" NOX=nitrogen oxides
1	3.741	12.0	0.907	
2	2.295	12.0	0.761	
3	1.498	12.0	1.108	
4	2.881	12.0	1.016	
5	0.760	12.0	1.189	
6	3.120	9.0	1.001	
7	0.638	9.0	1.231	
8	1.170	9.0	1.123	
9	2.358	12.0	1.042	
10	0.606	12.0	1.215	
11	3.669	12.0	0.930	
12	1.000	12.0	1.152	
13	0.981	15.0	1.138	
14	1.192	18.0	0.601	
15	0.926	7.5	0.696	
16	1.590	12.0	0.686	
17	1.806	12.0	1.072	
18	1.962	15.0	1.074	
19	4.028	15.0	0.934	
20	3.148	9.0	0.808	
21	1.836	9.0	1.071	
22	2.845	7.5	1.009	
23	1.013	7.5	1.142	
24	0.414	18.0	1.229	
25	0.812	18.0	1.175	


26	0.374	15.0	0.568
27	3.623	15.0	0.977
28	1.869	7.5	0.767
29	2.836	7.5	1.006
30	3.567	9.0	0.893
31	0.866	15.0	1.152
32	1.369	15.0	0.693
33	0.542	15.0	1.232
34	2.739	15.0	1.036
35	1.200	15.0	1.125
36	1.719	9.0	1.081
37	3.423	9.0	0.868
38	1.634	7.5	0.762
39	1.021	7.5	1.144
40	2.157	7.5	1.045
41	3.361	18.0	0.797
42	1.390	18.0	1.115
43	1.947	18.0	1.070
44	0.962	18.0	1.219
45	0.571	9.0	0.637
46	2.219	9.0	0.733
47	1.419	9.0	0.715
48	3.519	9.0	0.872
49	1.732	7.5	0.765
50	3.206	7.5	0.878
51	2.471	7.5	0.811
52	1.777	15.0	0.676
53	2.571	18.0	1.045
54	3.952	18.0	0.968
55	3.931	15.0	0.846
56	1.587	15.0	0.684
57	1.397	7.5	0.729
58	3.536	7.5	0.911
59	2.202	7.5	0.808
60	0.756	7.5	1.168
61	1.620	7.5	0.749
62	3.656	7.5	0.892
63	2.964	7.5	1.002
64	3.760	18.0	0.812
65	0.672	18.0	1.230

66	3.677	18.0	0.804
67	3.517	12.0	0.813
68	3.290	12.0	1.002
69	1.139	9.0	0.696
70	0.727	9.0	1.199
71	2.581	9.0	1.030
72	0.923	15.0	0.602
73	1.527	15.0	0.694
74	3.388	15.0	0.816
75	2.085	15.0	1.037
76	0.966	15.0	1.181
77	3.488	7.5	0.899
78	0.754	7.5	1.227
79	0.797	9.0	1.180
80	2.064	7.5	0.795
81	3.732	18.0	0.990
82	0.586	18.0	1.201
83	0.561	7.5	0.629
84	0.563	9.0	0.608
85	0.678	12.0	0.584
86	0.370	15.0	0.562
87	0.530	18.0	0.535
88	1.900	18.0	0.655



Understand the graph

Note that labeling could be improved.



NOX is dependent, ER and CR are independent.


EQ – richness of mixture

CR – max volume of cylinder/min volume of cylinder

Fig 4.6 : 9 slices of ER, $f=.25$ (overlap), NOX vs CR examined. Big interaction here. What if only top four or bottom five panels? Interaction?

Fig 4.7: 5 slices (not really slices in usual sense) NOX vs ER examined. Max increases.

```
> attach(ethanol.df)
> a=coplot(NOX~CR|ER)
```



Continuation of Ch 4 next time