

[12] 1. In parts (a)–(d) below, algebraic simplification is *not required*:

(a) Calculate $\frac{d}{dx}(e^x \tan x)$.

(b) Find $f'(x)$, given $f(x) = \left(x^2 + \sqrt{\frac{x - \pi}{7}}\right)^{2001}$.

(c) Given $g(t) = \sin(2 \ln t)$, find $g''(1)$.

(d) Suppose $u = \frac{\sin x^2}{1 + \cos^2 x}$. Find $\frac{du}{dx}$.

[6] 2. Find all points on the curve $y = \sin^{-1}(x)$ where the tangent line is parallel to the line

$$2x - \sqrt{3}y = 100.$$

[6] 3. Use the definition of the derivative as a limit to find $f'(3)$ for $f(x) = x^{-2}$.
[No marks will be given for an answer obtained using only differentiation rules.]

[4] 4. Find the positive constant k for which $y = k\sqrt{5x+1}$ satisfies the equation $y \frac{dy}{dx} = 1$.

[8] 5. Consider the curve $3^x - 2^y = 1$.

(a) Find the equation of the tangent line at the point $(2, 3)$.

(b) By expressing $\frac{dy}{dx}$ as a function of x , or otherwise, find $\lim_{x \rightarrow \infty} \frac{dy}{dx}$.

[8] 6. A ladder 5 metres long is leaning against a high vertical wall when its base begins to slip horizontally away from the wall. The distance s from the base to the wall (measured in metres) satisfies the differential equation

$$\frac{ds}{dt} = 1 + e^{-s}.$$

(Time is measured in seconds.)

Consider the area of the right-angled triangle formed by the ladder, the wall, and the ground. At the instant when the top of the ladder is 3 metres above the ground, ...

(a) is this area increasing or decreasing?

(b) what is the area's exact rate of change? (Give units with your answer.)

- [8] 7. Let $I(x)$ be the amount of light that gets through an x millimeter thick layer of tinted glass. Then $\frac{dI}{dx} = -kI$ for some positive constant k .

Suppose that a 1 mm layer of the glass allows 60% of the incident light to get through. How thick a layer of the glass should be used so that 1% of the incident light gets through?

- [6] 8. Suppose the function $y = y(t)$ satisfies this differential equation for some $c \geq 0$:

$$y''(t) + cy'(t) + y(t) = 0.$$

Use y to define $E(t) = (y(t))^2 + (y'(t))^2$. Prove that whenever $t_1 < t_2$, we have $E(t_1) \geq E(t_2)$. [Note: It is possible to present a convincing proof without solving the differential equation.]

- [6] 9. A moving particle's displacement s is given by $s(t) = e^{-t} \sin t$ at all times $t > 0$.

- (a) For which values of t in the interval $0 < t < \pi$ is the particle's velocity positive?
- (b) For which values of t in the interval $0 < t < \pi$ is the particle's acceleration positive?

- [8] 10. A certain function f obeys $f''(x) < 0$ for all x , and $f(2) = 2$. In seeking a zero of f with Newton's Method, the starting point $x_0 = 2$ gives the next guess $x_1 = 1$.

- (a) Find $f'(2)$.
- (b) With the aid of a suitable sketch, explain why f must have a zero at some point x satisfying $1 < x < 2$.

Your answer should apply to every function f with the properties described above.

- [10] 11. Find the dimensions of the circular cylinder of greatest volume that can be inscribed in a cone of base radius R and height H , if the base of the cylinder lies in the base of the cone. Express your answer in terms of R and H .

- [9] 12. A certain function f is given. Its second derivative, $f''(x)$, is defined and continuous for all real x . Furthermore, f satisfies $f(-2) = 0$, $f'(-2) = 0$, $f(0) = -2$, $f(4) = 0$. For $y = f(x)$,


$$\begin{aligned} x < -2 &\implies y' < 0, y'' > 0, \\ -2 < x < 0 &\implies y' < 0, y'' < 0, \\ 0 < x < 2 &\implies y' < 0, y'' > 0, \\ 2 < x &\implies y' > 0, y'' > 0. \end{aligned}$$

Sketch the curve $y = f(x)$, paying particular attention to slope and concavity. Label any local maxima and minima and points of inflection on your sketch.

[9] **13.** Let $f(x) = 3x^4 + 4(k-2)x^3 - 3kx^2$.

The curve $y = f(x)$ is shown for the case $k = 5/4$.

- (a) Find all real numbers k for which f has a local maximum at the point $x = 0$.
- (b) Let x_L and x_R denote the x -coordinates of the local minimum points for f , as illustrated in the sketch provided. Assuming $k > 0$, express the separation $s = |x_R - x_L|$ as a function of k .



- (c) Among all $k > 0$, which choice minimizes the separation s described in part (b)? Why?

UBC-SFU-UVic-UNBC

Calculus Examination

7 June 2001

Name: _____ Signature: _____

School: _____ Candidate Number: _____

Rules and Instructions

1. *Show all your work!* Full marks are given only when the answer is correct, and is supported with a written derivation that is orderly, logical, and complete. Part marks are available in every question.
2. Calculators are optional, not required. Correct answers that are “calculator ready,” like $3 + \ln 7$ or $e^{\sqrt{2}}$, are fully acceptable.
3. Any calculator acceptable for the Provincial Examination in Principles of Mathematics 12 may be used.
4. No notes, books, or other aids are allowed. In particular, *all calculator memories must be empty when the exam begins.*
5. If you need more space to solve a problem on page n , work on the back of page $n - 1$.
6. CAUTION - Candidates guilty of any of the following or similar practices shall be dismissed from the examination immediately and assigned a grade of 0:
 - (a) Using any books, papers or memoranda.
 - (b) Speaking or communicating with other candidates.
 - (c) Exposing written papers to the view of other candidates.
7. Do not write in the grade box shown to the right.

1		12
2		6
3		6
4		4
5		8
6		8
7		8
8		6
9		6
10		8
11		10
12		9
13		9
Total		100