

Physics 101

Lecture 30.

Wed. March. 19. 2003.


Topics covered:

Streamline flow of fluid.

- equation of continuity.
- incompressible fluid (Liquid)
- Bernoulli's equation.
- Torricelli's law.

Ref: 15-6, 7, 8.

- 30-2
~~29-5~~
- Fluid Dynamics. (Assuming No turbulence)
i.e., streamline flow.



Basic Principle: in Δt ,
Amount of mass in = Amount of mass out.

$$\rho_1 A_1 v_1 \Delta t = \rho_2 A_2 v_2$$

Equation of continuity

$$\rho_1 A_1 v_1 = \rho_2 A_2 v_2$$

Same density — for incompressible fluid (liquid).

$$A_1 v_1 = A_2 v_2$$

$$v_2 = \frac{A_1}{A_2} v_1$$

To make v_2 large at the output,

we can reduce A_2 .


e.g.: Nozzle of a fire hose.



Dynamics 2 Pressure can cause motion of a fluid.

Assuming
same density.

In time
interval Δt :



(Ignore drag).

Pressure can do work $\Rightarrow$ Energy is increased.

$$\begin{aligned} \text{Work by Pressure} &= P_1 A_1 \cdot v_1 \Delta t - P_2 v_2 A_2 \cdot \Delta t \\ &= \frac{\Delta m}{\rho} (P_1 - P_2) \end{aligned}$$

$$\text{Gain in M. E.} = \Delta U + \Delta K$$

$$= \Delta m (y_2 - y_1) \cdot g + \frac{1}{2} \cdot \Delta m (v_2^2 - v_1^2)$$

$$\therefore \frac{\Delta m}{\rho} (P_2 - P_1) = \Delta m (y_2 - y_1) g + \frac{1}{2} \Delta m (v_2^2 - v_1^2)$$

∴ Bernoulli's Equation :

$$P_1 + \rho g y_1 + \frac{1}{2} \rho v_1^2 = P_2 + \rho g y_2 + \frac{1}{2} \rho v_2^2$$

OR: $P + \rho g y + \frac{1}{2} \rho v^2 = \text{constant}$

• Special Case

When $y_1 = y_2 = \text{const}$.


B. E. : $P_1 + \frac{1}{2} \rho v_1^2 = P_2 + \frac{1}{2} \rho v_2^2$

OR: $P + \frac{1}{2} \rho v^2 = \text{const}$

There is a tradeoff between P and v^2 .

e.g: Moving fluids have a lower pressure than stationary fluid.


Demo:



e.g.: Torricelli's Law .

Large container with
a small hole near the
bottom .

What's the speed
of the ~~flow~~ fluid
when it flows out
of the small hole ?



$$B. \Sigma q: \quad P_1 + \frac{1}{2} \rho v_1^2 + \rho g y_1 = P_2 + \frac{1}{2} \rho v_2^2 + \rho g y_2 .$$

$$\therefore P_1 = P_2 = P_{at} ,$$

$$v_1 = 0 . \quad \text{since } A_1 \text{ is very large .}$$

$$\therefore \rho g y_1 = \frac{1}{2} \rho v_2^2 + \rho g y_2 .$$

$$\therefore v_2^2 = 2g(y_1 - y_2) = 2gh .$$

$$v_2 = \sqrt{2gh} .$$

Conservation of Mechanical Energy . (just like a free-falling object)